## Appendix

## Acute Flaccid Myelitis Outcome – follow-up of confirmed and probable AFM cases (*completed at 60 days, 6 months and 12 months after onset of limb weakness*)

**34**. Date of follow-up: \_\_\_/\_\_\_/ (*mm/dd/yyyy*)

**35.** Impairment:
 □ None
 □ Minor (any minor involvement)
 □ Significant (≤2 extremities, major involvement)

 □ Severe (≥3 extremities and respiratory involvement)
 □ Death
 □ Unknown

**35a.** Date of death: \_\_ \_/\_\_ \_/ \_\_ \_\_ (*mm/dd/yyyy*)

36. Physical condition (includes cardiovascular, gastrointestinal, urologic, endocrine as well as neurologic disorders):

- i. Medical problems sufficiently stable that medical or nursing monitoring is not required more often than 3-month intervals
- ii. Medical or nurse monitoring is needed more often than 3-month intervals but not each week.
- iii. Medical problems are sufficiently unstable as to require medical and/or nursing attention at least weekly.
- iv. Medical problems require intensive medical and/or nursing attention at least daily (excluding personal care assistance)

**37.** Upper limb functions: Self-care activities (drink/feed, dress upper/lower, brace/prosthesis, groom, wash, perineal care) dependent mainly upon upper limb function:

- i. Age-appropriate independence in self-care without impairment of upper limbs
- ii. Age-appropriate independence in self-care with some impairment of upper limbs
- iii. Dependent upon assistance in self-care with or without impairment of upper limbs.
- iv. Dependent totally in self-care with marked impairment of upper limbs.

**38.** Lower limb functions: Mobility (walk, stairs, wheelchair, transfer chair/toilet/tub or shower) dependent mainly upon lower limb function:

- i. Independent in mobility without impairment of lower limbs
- ii. Independent of mobility with some impairment of lower limbs, such as needing ambulatory aids, a brace or prosthesis
- iii. Dependent upon assistance or supervision in mobility with or without impairment of lower limbs.
- iv. Dependant totally in mobility with marked impairment of lower limbs.

39. Sensory components: Relating to communication (speech and hearing) and vision:

- i. Age-appropriate independence in communication and vision without impairment
- ii. Age-appropriate independence in communication and vision with some impairment such as mild dysarthria, mild aphasia or need for eyeglasses or hearing aid.
- iii. Dependent upon assistance, an interpreter, or supervision in communication or vision
- iv. Dependent totally in communication or vision

40. Excretory functions (bladder and bowel control, age-appropriate):

- i. Complete voluntary control of bladder and bowel sphincters
- ii. Control of sphincters allows normal social activities despite urgency or need for catheter, appliance, suppositories, etc.
- iii. Dependent upon assistance in sphincter management
- iv. Frequent wetting or soiling from bowel or bladder incontinence

## 41. Support factors:

- i. Able to fulfil usual age-appropriate roles and perform customary tasks
- ii. Must make some modifications in usual age-appropriate roles and performance of customary tasks
- iii. Dependent upon assistance, supervision, and encouragement from an adult due to any of the above considerations
- iv. Dependent upon long-term institutional care (chronic hospitalization, residential rehabilitation, etc. Excluding timelimited hospitalization for specific evaluation or treatment)