



Published in final edited form as:

*Equine Vet Educ.* 2023 ; 35(11): 589–593. doi:10.1111/eve.13843.

## On skunk rabies and its prevention in North America

Andres Velasco-Villa

Centers for Disease Control and Prevention, Atlanta, Georgia, USA

The case report by Navarro-López et al. (2023) in this issue describes a thorough differential diagnosis in a horse with acute neurological disease. Rabies was confirmed by direct fluorescent antibody (DFA) test. Rabies virus (RABV) was isolated from brain tissue of the ill mare by intracerebral inoculation in sucking mice. Sequencing and phylogenetic inference allowed the identification of the rabies virus variant (RVV) associated with this case. Thus, Navarro-López et al. reported that the rabid mare got infected with a RVV associated with skunks. This clinical commentary elaborates on the seemingly rare skunk rabies across North America (NA), highlighting its relevance in human and animal health that have remained somehow neglected.

Over the last two centuries, skunks have been one of the most common wildlife rabid animals across North America (Gremillion-Smith & Woolf, 1988; Ma et al., 2023; Oertli et al., 2009). Early reports of skunk rabies transmitted to humans, companion animals and farm animals, date from 1826 in lower California (Johnson, 1971), where most exposures were associated with spotted skunks (*Spilogale* spp.). Skunk rabies across the Central Great Plains mainly involving striped skunks (*Mephitis mephitis*) began to be observed during the second half of the 19th century (Hovey, 1874).

Due to advances in molecular epidemiology, phylogenetics and notable improvements on wildlife rabies surveillance (primarily passive both laboratory-based and epidemiological), it is now known that the skunk rabies epizootic is comprised by regional independent enzootics in which RVVs with particular evolutionary histories predominate (Charlton et al., 1988; Clark et al., 2015; Kuzmina et al., 2013). The north-central skunk (NCSK), the California skunk (CASK), the Baja California Sur Mexico skunk (BCSMXSK), the Northwest Mexico skunk (NWMXSK) and the Yucatan Mexico skunk (YUCMXSK) enzootics maintain circulation of specific RVVs that originated from long-standing dog rabies epizootics prevailing in their respective regions, as shown in Figure 1 (Barton & Wisely, 2012; Davis et al., 2013; Kuzmina et al., 2013; Velasco-Villa et al., 2002, 2005, 2008, 2017). Meanwhile, the South-central skunk (SCSK) and the North-central Mexico skunk (NCMXSK) enzootics maintain RVVs that originated from ancestral bat rabies

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

**Correspondence:** Andres Velasco-Villa, dly3@cdc.gov.

**AUTHOR CONTRIBUTIONS**

This is a sole author contribution.

**CONFLICT OF INTEREST STATEMENT**

No conflict of interest has been declared.

epizootics (see Figure 1) (Barton & Wisely, 2012; Davis et al., 2013; Kuzmina et al., 2013; Velasco-Villa et al., 2002, 2005, 2008, 2017).

Thus far, individuals of all skunk species in North America have been reported to be rabid (*Conepatus leuconotus*, *Conepatus semistriatus*, *Mephitis mephitis*, *Mephitis macroura*, *Spilogale gracillis*, *Spilogale putorius*, *Spilogale leucoparia* and *Spilogale yucatanensis*); however, only *M. mephitis* and some *Spilogale* spp have been recognised as major rabies reservoir hosts across North America (Aranda & López-de Buen, 1999; Dragoo et al., 2004; 2006; Gremillion-Smith & Woolf, 1988; Oertli et al., 2009; Velasco-Villa et al., 2002, 2005).

Albeit their divergent origins, all RRVs circulating in North American skunk populations seem to be virulent and capable of causing rabies in any sympatric mammal that a rabid skunk may encounter (Barton & Wisely, 2012; Charlton et al., 1991; Hill et al., 1993). Rabies is a highly lethal infectious disease for all mammals including the rabies reservoir hosts – i.e. mammal species in the Carnivore and Chiroptera orders that play a central role in maintaining specific RRVs through intraspecific transmission (Smith, 1989). The most common mechanism of rabies transmission is by direct contact through bites of sick animals infected with RABV. However, rabies could be acquired through contact with RABV infected/contaminated saliva or brain tissue from a rabid animal in open wounds or mucosa (Fisher et al., 2018). The incubation period for rabies in skunks varies from 21 to 117 days, which is similar to that observed in most mammals (Borchering et al., 2012; Gremillion-Smith & Woolf, 1988; Oertli et al., 2009). A rabid skunk may stay alive for up to 10 days after the onset of neurological signs (Charlton et al., 1987). However, RABV shedding in saliva can occur up to 6 days before the onset of neurological signs (Charlton et al., 1991); thereby, any skunk bite should be considered as a rabies exposure, particularly if the animal is not available for laboratory diagnosis. The most common rabies signs in rabid skunks are abnormal behaviour (active during daytime), hyperesthesia, hypersensitiveness to light noise or movement, incoordination and extremely aggressive behaviour (Gremillion-Smith & Woolf, 1988; Oertli et al., 2009). Rabid skunk attacks on people or animals have been reported as incursions at animal enclosures, people's homes and/or camping sites (Gremillion-Smith & Woolf, 1988; Hovey, 1874; Johnson, 1971; Oertli et al., 2009).

Skunk rabies occurs throughout the year with disease peaks during spring, when males increase their regular home range for mating. Skunk rabies epizootic cycles have been reported to occur every 4, 6, 8 or 20 years in the US and Canada, depending on the region or the period of observation (Charlton et al., 1991; Gremillion-Smith & Woolf, 1988; Oertli et al., 2009; Pybus, 1988).

Understanding the current geographic distribution of rabies enzootics and the main reservoir host species associated with them allows for a better design of species targeted control, prevention and elimination strategies, as well as for tailored outreach campaigns aimed at the most vulnerable populations (Fehlner-Gardiner, 2018; Pepin et al., 2017; Wohlers et al., 2018). Striped skunks (*M. mephitis*) are considered generalist, while other species such as *Conepatus* spp. and *Spilogale* spp. could be more specialised inhabiting diverse ecosystems, comprising high forests to plains and semi-desertic regions, including transition areas. Unlike most skunk species, striped skunks have successfully adapted across the

urbanisation gradient, including densely populated areas (Harrington et al., 2017; Hass & Dragoo, 2017; Ramey et al., 2013). Skunks may be targeted for removal or exclusion due to the nuisance provoked by their pungent defence scent and the occasional damage they cause to people properties or crops (Harrington et al., 2017). In some instances, skunks may be removed through poisoning or trapping. The latter is not considered practical, nor cost-effective or feasible in rural and farm areas (Harrington et al., 2017). Skunk extermination or extirpation from certain areas often leads to ecological imbalances and the emergence of pests, namely rodents and some insects they predate on (Harrington et al., 2017; Hass & Dragoo, 2017). Measures that combine oral rabies vaccination and humane population control (such as hormone derived contraception or relocation measures) may be more acceptable and effective to mitigate rabies exposures to humans, companion and farm animals (Charlton et al., 1992; Elmore et al., 2017; Feldhamer et al., 2003; Richards et al., 2019).

Skunk rabies rarely affects humans in the US and Canada, with only a few cases reported in Mexico in the last three decades (Aranda et al., 1999; Arechiga-Ceballos et al., 2022). Securing camping tent's entrances and/or keeping doors closed at homes and stables may help prevent incursions of rabid skunks (Gremillion-Smith & Woolf, 1988; Hovey, 1874; Johnson, 1971; Oertli et al., 2009). There is no accurate estimation of the total number of rabies cases due to skunk RVVs or rabid skunk exposures in domesticated and wildlife animals throughout North America (Ma et al., 2023). Nonetheless, it would be recommended to be cautious of its presence in regions known enzootic for skunk rabies (Fehlner-Gardiner, 2018). Economic losses due to skunk rabies in farm animals have not been accurately estimated, but the loss of high-value breeders, racehorses or other farm animals due to skunk rabies have been reported (Aranda et al., 1999; Arechiga-Ceballos et al., 2022; Velasco-Villa et al., 2002). Commonly available parenteral rabies vaccines labelled for use in cattle, horses and other farm animals protect against all known RVVs, including those present in rabid skunks (Green et al., 2011; Wohlers et al., 2018). However, having pre-exposure vaccination (receiving vaccine before an exposure to rabies occurs), in both humans and animals, does not always warrant 100% immunity after severe exposure to a confirmed rabid animal (Larghi, 2004; National Association of State Public Health Veterinarians Committee, 2008). If the viral entry location (bites) is too close to the head or the spine, and a high RABV load has been delivered by a rabid animal, RABV could potentially reach the central nervous system (CNS) before the immune system can stop infection at the periphery (Brunt et al., 2021; Green et al., 2011). Once RABV reaches the CNS, rabies will be almost 100% fatal; thereby, it is highly recommended to consult a veterinarian or a rabies medical expert to help assess exposure levels and decide the pertinence to provide booster vaccinations (Green et al., 2011; National Association of State Public Health Veterinarians Committee, 2008; Rupprecht et al., 2006).

Currently, there are no systematic oral vaccination campaigns to reduce or eliminate the skunk rabies epizootic in North America (Elmore et al., 2017). However, some efforts have been conducted in the US and Canada to reduce or mitigate the associated risks that imply the expansion of skunk rabies outbreaks, or the establishment of new RVVs in certain skunk populations (Elmore et al., 2017; Fehlner-Gardiner et al., 2012; Slate et al., 2005). Examples of outbreaks and the detection of non skunk-associated RVVs in skunks include a skunk

rabies outbreak in Flagstaff Arizona, where skunk to skunk rabies transmission of a RVV maintained by big brown bats lasted over consecutive years, and southern Canada where intraspecific rabies transmission of the raccoon RVV in skunk populations sparked a rabies outbreak among skunks (Elmore et al., 2017).

Since skunks seem to have remarkable ecological plasticity and an outstanding susceptibility to RABV from different origins, it would not be surprising if new skunk rabies enzootic cycles appear (Elmore et al., 2017; Fehlner-Gardiner et al., 2012; Slate et al., 2009). Rabies vaccination should be considered to mitigate the effects of skunk-transmitted rabies in horses and all other rabies susceptible farm animals when pertinent (Green et al., 2011; National Association of State Public Health Veterinarians Committee, 2008; Rupprecht et al., 2006). Feasibility of oral rabies vaccination for skunks continues to be explored, and ongoing research efforts include improvement of the uptake (such as improved vaccine baits that could be more attractive for skunk consumption), as well as formulations that evaluate different vectors, and genetically modified RABVs and their concentrations (Elmore et al., 2017; Fehlner-Gardiner et al., 2012; Slate et al., 2009).

## FUNDING INFORMATION

Centers for Disease Control and Prevention.

## DISCLAIMER

The opinions and conclusions expressed in this report are those of the authors and do not necessarily represent the official position of the CDC or the US Department of Health and Human Services.


## REFERENCES

- Aranda M & López-de Buen L (1999) Rabies in skunks from Mexico. *Journal of Wildlife Diseases*, 35(3), 574–577. Available from: 10.7589/0090-3558-35.3.574 [PubMed: 10479094]
- Aréchiga Ceballos N, Puebla Rodríguez P & Aguilar Setién Á (2022) The new face of human rabies in Mexico, What's next after eradicating rabies in dogs. *Vector Borne and Zoonotic Diseases*, 22(2), 69–75. Available from: 10.1089/vbz.2021.0051 [PubMed: 35175137]
- Barton HD & Wisely SM (2012) Phylogeography of striped skunks (*Mephitis mephitis*) in North America: Pleistocene dispersal and contemporary population structure. *Journal of Mammalogy*, 93(1), 38–51. Available from: 10.1644/10-MAMM-A-270.1
- Borchering RK, Liu H, Steinhaus MC, Gardner CL & Kuang Y (2012) A simple spatiotemporal rabies model for skunk and bat interaction in Northeast Texas. *Journal of Theoretical Biology*, 314, 16–22. Available from: 10.1016/j.jtbi.2012.08.033 [PubMed: 22981921]
- Brunt S, Solomon H, Brown K & Davis A (2021) Feline and canine rabies in New York state, USA. *Viruses*, 13(3), 450. Available from: 10.3390/v13030450 [PubMed: 33802123]
- Charlton KM, Artois M, Prevec L, Campbell JB, Casey GA, Wandeler AI et al. (1992) Oral rabies vaccination of skunks and foxes with a recombinant human adenovirus vaccine. *Archives of Virology*, 123(1–2), 169–179. Available from: 10.1007/BF01317147 [PubMed: 1550495]
- Charlton KM, Casey GA & Campbell JB (1987) Experimental rabies in skunks: immune response and salivary gland infection. *Comparative Immunology, Microbiology and Infectious Diseases*, 10(3–4), 227–235. Available from: 10.1016/0147-9571(87)90033-6 [PubMed: 3427891]
- Charlton KM, Webster WA & Casey GA (1991) Skunk rabies. In: Baer GM (Ed.) *The natural history of rabies*, 2nd edition. Boca Raton (FL): CRC Press, Inc., pp. 307–324
- Charlton KM, Webster WA, Casey GA & Rupprecht CE (1988) Skunk rabies. *Reviews of Infectious Diseases*, 10(S4), 626–628. Available from: 10.1093/clinids/10.Supplement\_4.S626

- Clark R, Taylor A, Garcia F, Krone T & Brown HE (2015) Recognizing the role of skunks in human and animal rabies exposures in the southwest. *Vector Borne and Zoonotic Diseases*, 15(8), 494–501. Available from: 10.1089/vbz.2014.1719 [PubMed: 26273811]
- Davis R, Nadin-Davis SA, Moore M & Hanlon C (2013) Genetic characterization and phylogenetic analysis of skunk-associated rabies viruses in North America with special emphasis on the central plains. *Virus Research*, 174, 27–36. Available from: 10.1016/j.virusres.2013.02.008 [PubMed: 23524137]
- Dragoo JW, Matthes DK, Aragon A, Hass CC & Terry LY (2004) Identification of skunk species submitted for rabies testing in the desert southwest. *Journal of Wildlife Diseases*, 40(2), 371–376. Available from: 10.7589/0090-3558-40.2.371 [PubMed: 15362845]
- Elmore SA, Chipman RB, Slate D, Huyvaert KP, VerCauteren KC & Gilbert AT (2017) Management and modeling approaches for controlling raccoon rabies: the road to elimination. *PLoS Neglected Tropical Diseases*, 11(3), e0005249. Available from: 10.1371/journal.pntd.0005249 [PubMed: 28301480]
- Fehlner-Gardiner C. (2018) Rabies control in North America – past, present and future. *Revue Scientifique et Technique*, 37(2), 421–437. English. Available from: 10.20506/rst.37.2.2812 [PubMed: 30747137]
- Fehlner-Gardiner C, Rudd R, Donovan D, Slate D, Kempf L & Badcock J (2012) Comparing ONRAB<sup>®</sup> and RABORAL V-RG<sup>®</sup> oral rabies vaccine field performance in raccoons and striped skunks, New Brunswick, Canada and Maine, USA. *Journal of Wildlife Diseases*, 48, 157–167 [PubMed: 22247384]
- Feldhamer GA, Thompson BC & Chapman JA (2003) *Wild mammals of North America: biology, management, and conservation*. Baltimore, MD: Johns Hopkins University Press. Xiii. 1216.
- Fisher CR, Streicker DG & Schnell MJ. (2018) The spread and evolution of rabies virus: conquering new frontiers. *Nature Reviews Microbiology*, 16(4), 241–255. Available from: 10.1038/nrmicro [PubMed: 29479072]
- Green AL, Carpenter LR & Dunn JR (2011) Rabies epidemiology, risk assessment, and pre- and post exposure vaccination. *The Veterinary Clinics of North America. Exotic Animal Practice*, 14(3), 507–518, vii. Available from: 10.1016/j.cvex.2011.05.012 [PubMed: 21872785]
- Gremillion-Smith C & Woolf A (1988) Epizootiology of skunk rabies in North America. *Journal of Wildlife Diseases*, 24, 620–626. Available from: 10.7589/0090-3558-24.4.620 [PubMed: 3193556]
- Harrington LA, Marino J & King CM (2017) People and wild musteloids. In: Macdonald DW, Newman C & Harrington LA (Eds.) *Biology and conservation of Musteloids*. online edn. Oxford, UK: Oxford University Press, pp. 189–215.
- Hass CC & Dragoo JW (2017) Competition and coexistence in sympatric skunks. In: Macdonald DW, Newman C & Harrington LA (Eds.) *Biology and conservation of Musteloids*. Oxford, UK: Oxford University Press, pp. 464–477.
- Hill RE, Smith KE, Beran GW & Beard PD (1993) Further-studies on the susceptibility of raccoons (*Procyon-Lotor*) to a rabies virus of skunk origin and comparative susceptibility of striped skunks (*mephitis-mephitis*). *Journal of Wildlife Diseases*, 29, 475–477. [PubMed: 8355351]
- Hovey HC (1874) Rabies mephitica. *The American Journal of Science and Arts*, 7, 477–483.
- Johnson HN (1971) General epizootiology of rabies. In: Nagano Y & Davenport FM (Eds.) *Rabies. Proceedings of Working Conference on Rabies Sponsored by Japan-United States Cooperative Medical Science, Program*. Tokyo, Japan: University of Tokyo Press pp. 237–251.
- Kuzmina NA, Lemey P, Kuzmin IV, Mayes BC, Ellison JA, Orciari LA et al. (2013) The phylogeography and spatiotemporal spread of south-central skunk rabies virus. *PLoS One*, 8(12), e82348. Available from: 10.1371/journal.pone.0082348 [PubMed: 24312657]
- Larghi OP (2004) Perspectives for rabies control and eradication from domestic species in developing countries. *Developmental Biology (Basel)*, 119, 205–212.
- Ma X, Bonaparte S, Corbett P, Orciari LA, Gigante CM, Kirby JD et al. (2023) Rabies surveillance in the United States during 2021. *Journal of the American Veterinary Medical Association* (published online ahead of print 2023), 1–9. Available from: 10.2460/javma.23.02.0081

- National Association of State Public Health Veterinarians Committee. (2008) Compendium of animal rabies prevention and control, 2008. *Journal of the American Veterinary Medical Association*, 232(10), 1478–1486. Available from: 10.2460/javma.232.10.1478 [PubMed: 18479237]
- Navarro-López R, Pérez-de la Rosa JD, Solís-Hernández M, Cañez-de la Fuente JF, Villarreal-Silva M & Gómez-Romero N (2023) Skunk-associated rabies case in equine from Northern Mexico. *Equine Veterinary Education*, 1–5. Available from: 10.1111/eve.13788
- Oertli EH, Wilson PJ, Hunt PR, Sidwa TJ & Rohde RE (2009) Epidemiology of rabies in skunks in Texas. *Journal of the American Veterinary Medical Association*, 234, 616–620. Available from: 10.2460/javma.234.5.616 [PubMed: 19250039]
- Pepin KM, Davis AJ, Streicker DG, Fischer JW, VerCauteren KC & Gilbert AT (2017) Predicting spatial spread of rabies in skunk populations using surveillance data reported by the public. *PLoS Neglected Tropical Diseases*, 11(7), e0005822. Available from: 10.1371/journal.pntd.0005822 [PubMed: 28759576]
- Pybus MJ (1988) Rabies and rabies control in striped skunks (*Mephitis mephitis*) in three prairie regions of western North America. *Journal of Wildlife Diseases*, 24(3), 434–449. Available from: 10.7589/0090-3558-24.3.434 [PubMed: 3045346]
- Ramey CA, Mills KW, Fischer JW, McLean RG, Fagerstone KA & Engeman RM (2013) Graphically characterizing the movement of a rabid striped skunk epizootic across the landscape in northwestern Wyoming. *EcoHealth*, 10(3), 246–256. Available from: 10.1007/s10393-013-0853-3 [PubMed: 23812724]
- Richards S, Rusk R & Douma DA (2019) One health approach to rabies management in Manitoba Canada. *The Canadian Veterinary Journal*, 60(7), 737–743. [PubMed: 31281191]
- Rupprecht CE, Hanlon CA & Slate D (2006) Control and prevention of rabies in animals: paradigm shifts. *Developmental Biology (Basel)*, 125, 103–111.
- Slate D, Algeo TP, Nelson KM, Chipman RB, Donovan D, Blanton JD et al. (2009) Oral rabies vaccination in north america: opportunities, complexities, and challenges. *PLoS Neglected Tropical Diseases*, 3(12), e549. Available from: 10.1371/journal.pntd.0000549 [PubMed: 20027214]
- Slate D, Rupprecht CE, Rooney JA, Donovan D, Lein DH & Chipman RB (2005) Status of oral rabies vaccination in wild carnivores in the United States. *Virus Research*, 111(1), 68–76. Available from: 10.1016/j.virusres.2005.03.012 [PubMed: 15896404]
- Smith JS (1989) Rabies virus epitopic variation: use in ecologic studies. *Advances in Virus Research*, 36, 215–253. [PubMed: 2472046]
- Velasco-Villa A, Gómez-Sierra M, Hernández-Rodríguez G, Juárez-Islas V, Meléndez-Félix A, Vargas-Pino F et al. (2002) Antigenic diversity and distribution of rabies virus in Mexico. *Journal of Clinical Microbiology*, 40(3), 951–958. Available from: 10.1128/JCM.40.3.951-958.2002 [PubMed: 11880422]
- Velasco-Villa A, Mauldin MR, Shi M, Escobar LE, Gallardo-Romero NF, Damon I et al. (2017) The history of rabies in the Western hemisphere. *Antiviral Research*, 146, 221–232. Available from: 10.1016/j.antiviral.2017.03.013 [PubMed: 28365457]
- Velasco-Villa A, Orciari LA, Souza V, Juárez-Islas V, Gomez-Sierra M, Castillo A et al. (2005) Molecular epizootiology of rabies associated with terrestrial carnivores in Mexico. *Virus Research*, 111, 13–27. Available from: 10.1016/j.virusres.2005.03.007 [PubMed: 15896399]
- Velasco-Villa A, Reeder SA, Orciari LA, Yager PA, Franka R, Blanton JD et al. (2008) Enzootic rabies elimination from dogs and reemergence in wild terrestrial carnivores, United States. *Emerging Infectious Diseases*, 14(12), 1849–1854. Available from: 10.3201/eid1412.080876 [PubMed: 19046506]
- Wohlers A, Lankau EW, Oertli EH & Maki J (2018) Challenges to controlling rabies in skunk populations using oral rabies vaccination: a review. *Zoonoses and Public Health*, 65(4), 373–385. Available from: 10.1111/zph.12471 [PubMed: 29633545]





**FIGURE 1.** Geographic distribution of major skunk rabies epizootics in North America with most likely putative reservoir host species.