



Outbreak History

AT A GLANCE

Ebola disease was first identified in 1976 after an outbreak in what is now the Democratic Republic of Congo. Expand the sections below to read a brief summary of all known cases and outbreaks, organized by year.

Cases and Outbreaks of Ebola Disease by Year

COLLAPSE ALL —

2022

Uganda

November

- Species: *Sudan ebolavirus*
- Reported number of cases: **164**
- Reported number of deaths and percentage of fatal cases: **55 (34%)**

On September 20, 2022, the Ministry of Health confirmed an outbreak of Ebola (Sudan virus) in Mubende District, in western Uganda. This announcement came after a patient with a suspected viral hemorrhagic fever (VHF) was identified and isolated at Mubende Regional Referral Hospital. A sample from the patient was sent to the VHF laboratory at the Uganda Virus Research Institute, where Ebola disease (species Sudan ebolavirus) was confirmed. This marks the sixth Ebola outbreak in Uganda. Five of the six have been caused by the species Sudan ebolavirus. The outbreak was declared over on January 11, 2023, with 142 confirmed cases (and 22 probable) and 55 confirmed deaths.

Democratic Republic of the Congo

August–September

- Species: *Zaire ebolavirus*
- Reported number of cases: **1**
- Reported number of deaths and percentage of fatal cases: **1 (100%)**

On August 22, 2022, the Ministry of Health announced an outbreak of Ebola virus disease (EVD) in North Kivu Province. This announcement came after a fatal case was discovered in Beni Health Zone, one of the epicenters of DRC's 10th outbreak in 2018-2020. A specimen was collected and tested positive at the Beni Referral Hospital laboratory and later confirmed at the DRC's national laboratory in Goma – the Institut National de la Recherche Biomédicale (INRB). The outbreak was declared over on September 27, 2022, with only one confirmed case.

Democratic Republic of the Congo

April–July

- Species: *Zaire ebolavirus*
- Reported number of cases: **5**

- Reported number of deaths and percentage of fatal cases: **5 (100%)**

On April 23, the Ministry of Health of the Democratic Republic of the Congo (DRC) declared an outbreak of EVD in Mbandaka city, Equateur Province. This was the 14th EVD outbreak in DRC and marked the third in a series of outbreaks in Equateur province since 2018. Sequencing data from the first confirmed case in this outbreak indicated that this was a new spillover event from an animal to a person and was not directly linked to previous outbreaks. The following four cases were epidemiologically linked, or had known contact with an EVD patient, and were reported from Mbandaka and Wangata health zones. The outbreak was declared over on July 4, 2022.

2021

Democratic Republic of the Congo

October–December

- Species: *Zaire ebolavirus*
- Reported number of cases: **11**
- Reported number of deaths and percentage of fatal cases: **9 (82%)**

On October 8, the Ministry of Health in DRC announced an outbreak of EVD in Beni Health Zone, North Kivu Province. Subsequent cases were confirmed in the same health zone and three probable cases were retrospectively identified from September 2021. Sequencing data showed a link to the 2018–2020 outbreak in the same region, suggesting this outbreak was likely caused by a persistent infection in an EVD survivor. This outbreak, the 13th EVD outbreak in DRC, was declared officially over on December 16, 2021, 42 days after the last confirmed patient was released from treatment.

Democratic Republic of the Congo

February–May

- Species: *Zaire ebolavirus*
- Reported number of cases: **12**
- Reported number of deaths and percentage of fatal cases: **6 (50%)**

On February 7, 2021, the Ministry of Health (MOH) in the Democratic Republic of the Congo (DRC) announced that a case of Ebola virus disease (EVD) had been confirmed in Biena Health Zone, North Kivu Province. Subsequent cases were confirmed. North Kivu was previously affected by EVD during the 2018–2020 Ebola outbreak, the largest in DRC's history, which was declared over on June 25, 2020. Sequencing of samples suggests that cases in this outbreak were linked to cases in the area during the 2018–2020 outbreak and likely resulted from persistent infection in a survivor that led to either a relapse or sexual transmission of the virus. The outbreak was declared over on May 3, 2021.

Guinea

- Species: *Zaire ebolavirus*
- Reported number of cases: **23**
- Reported number of deaths and percentage of fatal cases: **12 (52.2%)**

On February 14, 2021, the Ministry of Health (MOH) in Guinea announced that cases of Ebola virus disease (EVD) had been confirmed in N'Zérékoré Prefecture, a forested rural region in southeast Guinea. These are the first cases of EVD confirmed in Guinea since the 2014–2016 West Africa outbreak, the largest in history, was declared over. Sequencing of samples from the outbreak was conducted and compared to sequences from cases during the 2014–2016 West Africa outbreak. While researchers cannot definitively determine the cause of the outbreak, findings strongly support the conclusion that the outbreak was likely caused by a persistent source of infection (i.e. a survivor) and not a new introduction of the virus from the animal reservoir. The outbreak was declared over on June 19, 2021.

2020

Democratic Republic of the Congo

- Species: *Zaire ebolavirus*
- Reported number of cases: 130
- Reported number of deaths and percentage of fatal cases: **55 (42.3%)**

The DRC government declared a new Ebola outbreak in Mbandaka, Équateur Province of western DRC on June 1, 2020. International partners, including CDC, provided technical assistance to the DRC government to support response efforts. This was DRC's 11th Ebola outbreak and distinct from the 10th Ebola outbreak in eastern DRC, which was still ongoing when this one began.

Laboratory sequencing suggests that most cases in this outbreak were likely the result of a new spillover event (i.e. a new introduction of the virus into the community from an animal reservoir) followed by person-to-person spread. Sequencing efforts also identified a few cases which appeared to be linked to the prior Équateur Province outbreak in 2018, possibly due to sexual transmission or relapse of a survivor.

On November 18, 2020, the DRC Ministry of Health and WHO announced the outbreak was over.

2018

Democratic Republic of the Congo

- Species: *Zaire ebolavirus*
- Reported number of cases: 3,470*
- Reported number of deaths and percentage of fatal cases: **2,287* (66%)**

The DRC government declared its 10th Ebola outbreak on August 1, 2018, in North Kivu province of eastern DRC. Cases were also reported in Ituri and South Kivu provinces, and in Uganda. CDC assisted the DRC government, neighboring countries, and local and international partners to coordinate activities and provide technical guidance related to laboratory testing, contact tracing, infection control, border health screening, data management, risk communication and health education, vaccination, and logistics.

The outbreak was declared over by the World Health Organization (WHO) on June 25, 2020.

*In 2019, four cases confirmed in Uganda were attributed to cross-border movement from DRC and recorded in both countries. These cases died in DRC and are reported in the DRC death count.

Democratic Republic of the Congo

- Species: *Zaire ebolavirus*
- Reported number of cases: **54**
- Reported number of deaths and percentage of fatal cases: **33 (61%)**

The DRC government declared the outbreak in the Bikoro region of Équateur Province in the northwestern part of the country on May 8 after two cases were confirmed by laboratory testing at the Institut National de Recherche Biomédicale in Kinshasa. CDC assisted the DRC government and local and international partners, including the World Health Organization (WHO), as they pursued priority areas of support, including establishing an outbreak response platform; implementing surge support for deployment of personnel, supplies, laboratory materials, operational support, logistics, and transportation; and identifying communication needs to support the partners and the response. On July 24, 2018, WHO declared the end of the ninth outbreak of Ebola in the Democratic Republic of the Congo.

2017

Democratic Republic of the Congo

- Species: *Zaire ebolavirus*
- Reported number of cases: **8**
- Reported number of deaths and percentage of fatal cases: **4 (50%)**

On May 11, 2017, the Ministry of Public Health of the Democratic Republic of the Congo notified international public health agencies of a cluster of suspected cases of Ebola Virus Disease (EVD) in the Likati health zone of the province of Bas Uélé. The first report mentioned eight suspected cases, including two deaths, with a third death reported on May 12. Testing of samples was conducted by the Institut National de Recherche Biomedicale (INRB) in Kinshasa, with two samples testing positive for Ebola Zaire. Teams from international agencies, including CDC, WHO, MSF (Doctors without Borders), and others, supported the Ministry of Public Health's epidemiologic, diagnostic, clinical, and communications efforts to respond to the outbreak. The response faced challenging logistical obstacles, including the remoteness of the area and limited services. Mobile diagnostic laboratories provided testing of samples in the affected areas. Following a period of 42 days since the second negative laboratory diagnostic test of the last confirmed patient, WHO declared an end to the outbreak on July 2, 2017.

2014

Democratic Republic of the Congo

- Species: *Zaire ebolavirus*
- Reported number of cases: **69**
- Reported number of deaths and percentage of fatal cases: **49 (71%)**

The outbreak occurred in multiple villages in the vicinity of Boende town but was limited to the Équateur province in the western part of the country. The Ebola virus variant that caused this outbreak was closely related to the one that caused the 1995 outbreak in Kikwit, indicating that this outbreak was not related to the large outbreak happening at the same time in West Africa.

Guinea, Liberia, Sierra Leone (West African Epidemic)

- Species: *Zaire ebolavirus*
- Reported number cases: **28,610**
- Reported number of deaths and percentage of fatal cases: **11,308 (39%)**

The West African Ebola epidemic was the largest in history. It started with cases of EVD in the forested rural region of southeastern Guinea, reported by WHO on March 23, 2014. Soon, cases were discovered in Liberia and Sierra Leone, which border Guinea. Identification of cases was difficult because of weak surveillance and fragile public health infrastructure. Poor infection control measures and strained health care systems also contributed to the devastation of this outbreak. By August 2014, WHO declared the situation a Public Health Emergency of International Concern (PHEIC). CDC collaborated with other U.S. government agencies, Ministries of Health of the affected countries, WHO, and national and international partners in the intensive effort to end the epidemic. The outbreak was declared over in June 2016.

Italy

- Species: *Zaire ebolavirus*
- Reported number of cases: **1**
- Reported number of deaths: **0**

During the West African Ebola epidemic, an Italian healthcare worker who had volunteered in an Ebola Treatment Unit in Sierra Leone developed symptoms of EVD 72 hours after returning to Rome. The patient had no symptoms while in flight, so other airline passengers were not at risk. The healthcare workers treating the patient were adequately protected and carefully monitored. There were no additional cases in Italy.

Mali

- Species: *Zaire ebolavirus*
- Reported number of cases: **8**
- Reported number of deaths and percentage of fatal cases: **6 (75%)**

During the West African Ebola epidemic, an infected traveler from Guinea brought EVD into Mali. Immediate identification and monitoring of all the contacts of infected people successfully prevented a broader outbreak.

Nigeria

- Species: *Zaire ebolavirus*
- Reported number of cases: **20**
- Reported number of deaths and percentage of fatal cases: **8 (40%)**

During the West African Ebola epidemic, an infected person traveling from Monrovia, Liberia brought EVD into Lagos, Nigeria. Unprotected responders were subsequently infected. Fearing a rapid spread within Africa's most populated city, emergency response efforts focused on quickly establishing an Ebola Treatment Unit, training caregivers, and identifying all contacts for each case of EVD. As a result of this rapid response, the outbreak was confined to two cities, and additional spread of the Ebola virus through Nigeria and into other areas in Africa was prevented.

Senegal

- Species: *Zaire ebolavirus*
- Reported number of cases: **1**
- Reported number of fatal cases: **0**

During the West African Ebola epidemic, an infected traveler brought EVD into Senegal. Immediate identification and monitoring of all the contacts of the infected person successfully prevented a broader outbreak and no additional cases occurred in Senegal.

Spain

- Species: *Zaire ebolavirus*
- Reported number of cases: **1**
- Reported number of deaths: **0**

This was the first case of human-to-human transmission outside of Africa during the West African Ebola epidemic. A healthcare worker in Spain was infected with Ebola virus while treating a patient recently evacuated from Sierra Leone. The patient later died. The healthcare worker was monitored for signs of EVD and treated in isolation at the start of a fever. Contacts of the healthcare worker and other hospital staff were also monitored. The healthcare worker recovered and there were no other cases reported in Spain.

United Kingdom

- Species: *Zaire ebolavirus*
- Reported number of cases: **1**
- Reported number of deaths and percentage of fatal cases: **0 (0%)**

During the West Africa epidemic, a healthcare worker returning to the UK after volunteering in an Ebola treatment center in Sierra Leone became symptomatic upon arrival in the UK. The case was treated in strict isolation and a range of public health measures were implemented by UK authorities. This was the first case of Zaire ebolavirus to be detected in the UK.

United States

- Species: *Zaire ebolavirus*
- Reported number of cases: **4**
- Reported number of deaths and percentage of fatal cases: **1 (25%)**

During the West African Ebola outbreak, 11 people were treated for EVD in the U.S., two of whom died. The majority were infected with the Ebola virus outside of the U.S. and either medically evacuated into the U.S. for treatment or entered the country as a regular airline passenger. Two nurses who cared for a sick Ebola patient contracted EVD, marking the first known transmission of EVD in the United States. Both recovered. CDC collaborated with U.S. Customs and Border Protection, Department of Homeland Security, and state and local public health departments to screen travelers returning from Ebola-affected countries, provide safe transport for patients being assessed for EVD, and strengthen preparedness and infection control in hospitals.

2012

Uganda

- Species: *Sudan ebolavirus*
- Reported number of cases: **6***
- Reported number of deaths and percentage of fatal cases: **3* (50%)**

The outbreak occurred in November in the Luwero, Jinja, and Nakasongola districts. Through a collaboration established in 2010, CDC provided assistance to the Uganda Ministry of Health and Uganda Virus Research Institute (UVRI) laboratory throughout the outbreak.

* Numbers reflect laboratory-confirmed cases only

Democratic Republic of the Congo

- Species: *Bundibugyo ebolavirus*
- Reported number of cases: **38***
- Reported number of deaths and percentage of fatal cases: **13* (34%)**

The outbreak occurred in the Orientale province in the northeast of the country. CDC and the Public Health Agency of Canada (PHAC) provided laboratory support through a field laboratory in Isiro Health Zone, as well as through the CDC/UVRI (Uganda Virus Research Institute) laboratory in Uganda. This outbreak in DRC had no epidemiologic link to the Ebola outbreak occurring in the Kibaale district of Uganda at the same time.

* Numbers reflect laboratory-confirmed cases only

Uganda

- Species: *Sudan ebolavirus*
- Reported number of cases: **11***
- Reported number of deaths and percentage of fatal cases: **4* (36%)**

The outbreak occurred in June in Kibaale District. Laboratory tests of blood samples were conducted by Uganda Virus Research Institute (UVRI) and CDC

* Numbers reflect laboratory-confirmed cases only

2011

Uganda

- Species: *Sudan ebolavirus*
- Reported number of cases: **1**
- Reported number of deaths and percentage of fatal cases: **1 (100%)**

The Uganda Ministry of Health informed the public that a patient with suspected EVD died on May 6, 2011, in the Luwero district. Quick diagnosis of Ebola virus from a blood sample was provided by the newly established CDC Viral Hemorrhagic Fever laboratory at the Uganda Virus Research Institute (UVRI). Factors that helped stop the spread of the Ebola virus and limit the size of this outbreak included: 1) a high suspicion of hemorrhagic fever by clinical staff at the outset, 2) the correct use of personal protective equipment and barrier methods to protect hospital staff, and 3) the ability to rapidly confirm Ebola virus through laboratory testing in-country.

2008

Democratic Republic of the Congo

- Species: *Zaire ebolavirus*
- Reported number of cases: **32**
- Reported number of deaths and percentage of fatal cases: **15 (47%)**

The outbreak occurred in the Mweka and Luebo health zones in the Kasai Occidental province. A number of international partners were involved in the response to this outbreak.

The Philippines

- Species: *Reston ebolavirus* (does not cause illness in people)
- Reported number of cases: **6 (asymptomatic)**
- Reported number of deaths: **0**

This was the first known occurrence of Ebola-Reston virus in pigs. The virus strain was similar to earlier strains. Six workers from a pig farm and slaughterhouse developed antibodies against the virus but did not become sick.

2007

Uganda

- Species: *Bundibugyo ebolavirus*
- Reported number of cases: **131**
- Reported number of deaths and percentage fatal cases: **42 (32%)**

The outbreak occurred in the Bundibugyo district. This is the first reported occurrence of a new Ebola virus strain. This novel strain appeared similar to other related viruses. However, the case fatality rate was below 50% in this outbreak, compared to 80–90% found in

earlier outbreaks caused by other Ebola virus strains, like Zaire ebolavirus.

The Democratic Republic of the Congo

- Species: *Zaire ebolavirus*
- Reported number of cases: **264**
- Reported number of deaths and percentage of fatal cases: **187 (71%)**

The outbreak was declared in mid-September in Luebo and Mweke health zones in the Kasai Occidental Province. Radio broadcasts were used to deliver accurate and timely messages to the local population about EVD spread and prevention. The last confirmed case was on October 4 and the outbreak was declared over November 20.

2005

The Republic of the Congo

- Species: *Zaire ebolavirus*
- Reported number of cases: **12**
- Reported number of deaths and percentage of fatal cases: **10 (83%)**

Two hunters (index patients) died in Etoumbi Medical Center in April 2005. A response team led by the Ministry of Health was rapidly sent to the site. Most cases were hunters, patient caretakers, or funeral attendees.

2004

Russia

- Species: *Zaire ebolavirus*
- Reported number of cases: **1**
- Reported number of deaths and percentage of fatal cases: **1(100%)**

A Russian laboratory worker was injected with the virus accidentally while working on an Ebola vaccine and later died.

Sudan

- Species: *Sudan ebolavirus*
- Reported number of cases: **17**
- Reported number of deaths and percentage of fatal cases: **7 (41%)**

The outbreak occurred in Yambio County at the same time as an outbreak of measles in the same area. Several suspected cases of EVD were later reclassified as measles cases.

2003

The Republic of the Congo

November–December

- Species: *Zaire ebolavirus*
- Reported number of cases: **35**
- Reported number of deaths and percentage of fatal cases: **29 (83%)**

The outbreak occurred in Mbomo district in the Cuvette Ouest Département found in the western part of the country. Community mobilization activities were carried out with meetings held specifically for women in local villages, as they are usually the primary caretaker for patients in their families and communities.

The Republic of the Congo

January–April

- Species: *Zaire ebolavirus*
- Reported number of cases: **143**
- Reported number of deaths and percentage of fatal cases: **128 (89%)**

The outbreak occurred in Mbomo and Kéllé districts in the Cuvette Ouest Département in the western part of the country. Introduction of the virus into the population occurred after hunters reported close contact with wildlife that was killed or found dead. Direct contact with an infected person, particularly a family member, was the main mode of transmission, with very little spread of the virus within the healthcare setting.

2001

The Republic of the Congo

- Species: *Zaire ebolavirus*
- Reported number of cases: **59**
- Reported number of deaths and percentage of fatal cases: **44 (75%)**

The first time EVD was reported in the Republic of the Congo. The outbreak occurred over the border of Gabon and the Republic of the Congo. In the Republic of the Congo, the affected areas included Mbomo and Kéllé districts in the Cuvette Ouest Département located in the western part of the country. The first human cases were associated with hunting and contact with wildlife in the surrounding area.

Gabon

- Species: *Zaire ebolavirus*
- Reported number of cases: **65**
- Reported number of deaths and percentage of fatal cases: **53 (81%)**

The outbreak occurred over the border of Gabon and the Republic of the Congo. In Gabon, the affected areas included La Zadié, Ivindo, and Mpassa districts. An abnormal amount of animals, mostly nonhuman primates, were found dead in the area. The first human cases were associated with hunting and contact with wildlife in the surrounding area.

2000

Uganda

- Species: *Sudan ebolavirus*

- Reported number of cases: **425**
- Reported number of deaths and percentage of fatal cases: **224 (53%)**

The outbreak started in Gulu and later spread to Masindi and Mbarara districts. Community action and local government support were critical in controlling this outbreak. As gleaned from previous epidemics, providing correct and timely messaging about the disease was important to limit the spread of harmful rumors.

1996

Russia

- Species: *Zaire ebolavirus*
- Reported number of cases: **1**
- Reported number of deaths and percentage of fatal cases: **1 (100%)**

A Russian laboratory worker was infected with the Ebola virus while working on an experimental treatment for Ebola.

The Philippines

- Species: *Reston ebolavirus (does not cause illness in people)*
- Reported number of cases: **0**
- Reported number of deaths and percentage of fatal cases: **0**

Ebola-Reston virus was identified in a monkey export facility in the Philippines. No human infections were identified.

United States

- Species: *Reston ebolavirus (does not cause illness in people)*
- Reported number of cases: **0**
- Reported number of deaths and percentage of fatal cases: **0**

Ebola-Reston virus was introduced into a quarantine facility in Texas by monkeys imported from the Philippines. No human infections were identified.

South Africa

- Species: *Zaire ebolavirus*
- Reported number of cases: **2**
- Reported number of deaths and percentage of fatal cases: **1 (50%)**

A medical professional traveled from Gabon to Johannesburg, South Africa, after becoming exposed to the virus while treating Ebola-infected patients. He was hospitalized and recovered. A nurse treating him became infected and died.

Gabon

- Species: *Zaire ebolavirus*
- Reported number of cases: **60**
- Reported number of deaths and percentage of fatal cases: **45 (75%)**

The outbreak occurred in Booué in the fall. The first case was a hunter living in a logging camp who spread the virus to others. There were reports of several dead chimpanzees in the area. Testing of a skin sample obtained from one of the chimpanzees confirmed the animal was infected with Ebola virus.

Gabon

- Species: *Zaire ebolavirus*
- Reported number of cases: **31**
- Reported number of deaths and percentage of fatal cases: **21 (68%)**

The outbreak occurred in the spring in the village of Mayibout 2, located 0.62 miles (1 Km) from Mayibout 1 near the Ivindo River, about 40 km from the site of the 1994 outbreak. The body of a dead chimpanzee found in the forest was consumed by hunters. Eighteen people involved in the butchering of the animal became ill and other cases occurred in family members.

1995

The Democratic Republic of the Congo

- Species: *Zaire ebolavirus*
- Reported number of cases: **315**
- Reported number of deaths and percentage of fatal cases: **254 (81%)**

The outbreak occurred in Kikwit and surrounding areas and began with a charcoal maker in the forested areas near the city. The epidemic spread through families and hospitals. Transmission in the healthcare setting was halted almost immediately once proper protective measures were taken, such as the use of face masks, gloves, and gowns for healthcare personnel.

1994

Cote d'Ivoire

- Species: *Tai Forest ebolavirus*
- Reported number of cases: **1**
- Reported number of deaths: **0**

High mortality in the chimpanzee population in the Tai Forest was reported. A scientist became ill after conducting an autopsy on a wild chimpanzee. Diagnostic testing suggested a new strain of Ebola. The patient was treated in Switzerland and recovered.

Gabon

- Species: *Zaire ebolavirus*
- Reported number of cases: **51**
- Reported number of deaths and percentage of fatal cases: **31 (61%)**

The outbreak occurred in several gold mining villages deep in the rainforest around Makakou. It was initially believed to be yellow fever, but some of the characteristics of the outbreak were not typical for yellow fever. In 1995, researchers retrospectively discovered that Ebola virus was involved at the same time.

1992

Italy

- Species: *Reston ebolavirus* (does not cause illness in people)
- Reported number of cases: **0**
- Reported number of deaths and percentage of fatal cases: **0**

Ebola-Reston virus was introduced into quarantine facilities in Sienna by monkeys imported from the same export facility in the Philippines that was involved in the episodes in the United States. People were not infected.

1989

The Philippines

- Species: *Reston ebolavirus* (does not cause illness in people)
- Reported number of cases: **3 (asymptomatic)**
- Reported number of deaths: **0**

High mortality among *Cynomolgus* macaques was discovered in a primate facility responsible for exporting animals to the United States. Three workers in the animal facility developed antibodies but never experienced symptoms of Ebola Virus Disease.

United States

- Species: *Reston ebolavirus* (does not cause illness in people)
- Reported number of cases: **4 (asymptomatic)**
- Reported number and percentage of deaths among cases: **0**

Ebola-Reston virus was introduced into primate-holding facilities in Reston, Virginia; Philadelphia, Pennsylvania; and Alice, Texas by monkeys imported from the Philippines. Four people developed antibodies but never experienced symptoms of Ebola virus disease.

1979

Sudan

- Species: *Sudan ebolavirus*
- Reported number of cases: **34**
- Reported number of deaths and percentage of fatal cases: **22 (65%)**

The outbreak occurred in the towns of Nzara and Yambio, the same area affected by the 1976 Sudan epidemic. The index patient in this outbreak worked at the textile factory implicated as the source of the first outbreak in Sudan. Absentee and illness records for employees indicated that the factory was not the source of infection for this outbreak.

1977

The Democratic Republic of the Congo

- Species: *Zaire ebolavirus*
- Reported number of cases: **1**
- Reported number of deaths and percentage of fatal cases: **1 (100%)**

Case was noted retrospectively in the village of Tandala. This case had no known connection to the original Ebola outbreak in 1976, suggesting Ebola virus is enzootic in the area.

1976

United Kingdom

- Species: *Zaire ebolavirus*
- Reported number of cases: **1**
- Reported number of deaths: **0**

This case was a laboratory infection by accidental inoculation from a contaminated needle.

Sudan

- Species: *Sudan ebolavirus*
- Reported number of cases: **284**
- Reported number of deaths and percentage of fatal cases: **151 (53%)**

The outbreak occurred in the towns of Nzara, Maridi, and surrounding areas. The outbreak is believed to have started with workers in a cotton factory where 37% of workers in the cloth room were infected. The virus spread mainly through close personal contact within hospitals. Many healthcare personnel were infected.

The Democratic Republic of the Congo

- Species: *Zaire ebolavirus*
- Reported number of cases: **318**
- Reported number of deaths and percentage of fatal cases: **280 (88%)**

This outbreak was the first recognition of Ebola Virus Disease. It occurred in the Équateur province, with most cases occurring within 70 km of Yambuku village. The index case was treated at the Yambuku Mission Hospital with an injection for possible malaria. Subsequent transmission followed through use of contaminated needles and syringes at the hospital and clinics in the area and close personal contact. There were only 38 serologically confirmed survivors.

References

EXPAND ALL +

Sources

Albarino CG, Shoemaker T, Khristova ML, et al. [Genomic analysis of filoviruses associated with four viral hemorrhagic fever outbreaks in Uganda and the Democratic Republic of the Congo in 2012](#) [↗](#). *Virology*. 2013;442(2):97–100.

Baron RC, McCormick JB, and Zubeir OA. [Ebola virus disease in southern Sudan: hospital dissemination and intrafamilial spread \[PDF-864KB\]](#) [\[PDF\]](#) [↗](#). *Bulletin of the World Health Organization*. 1983;61(6):997–1003

[Déclaration de son Excellence Monsieur le Ministre de la Santé Publique annonçant la fin de l'épidémie de FHV à virus Ebola dans les zones de santé de Mweka, Luebo et Bulape dans la Province du Kasai Occidental \[PDF- 579KB\]](#) [\[PDF\]](#). Mardi, le 20 novembre 2007. Dr Victor Makwenge Kaput, Ministre de la Santé Publique.

Emond RT, Evans B, Bowen ET, et al. [A case of Ebola virus infection \[PDF – 817KB\]](#) [\[PDF\]](#) [↗](#). *British Medical Journal*. 1977;2(6086):541–544.

Formenty P, Libama F, Epelboin A, et al. Outbreak of Ebola hemorrhagic fever in the Republic of the Congo, 2003: a new strategy? *Médecine Tropicale (Marseille)*. 2003;63(3):291–295.

Georges AJ, Leroy EM, Renaud AA, et al. [Ebola hemorrhagic fever outbreaks in Gabon, 1994–1997: epidemiologic and health control issues \[PDF -299KB\]](#) [↗](#). *Journal of Infectious Diseases*. 1999;179:S65–75.

Hayes CG, Burans JP, Ksiazek TG, et al. Outbreak of fatal illness among captive macaques in the Philippines caused by an Ebola-related filovirus. *American Journal of Tropical Medicine and Hygiene*. 1992;46(6):664–671.

Heymann DL, Weisfeld JS, Webb PA, et al. [Ebola hemorrhagic fever: Tandala, Zaire, 1977–1978](#) [↗](#). *Journal of Infectious Diseases*. 1980;142(3):372–376.

Jahrling PB, Geisbert TW, Dalgard DW, et al. [Preliminary report: isolation of Ebola virus from monkeys imported to USA](#) [↗](#). *Lancet*. 1990;335(8688):502–505.

Khan AS, Tshioko FK, Heymann DL, et al. [The Reemergence of Ebola Hemorrhagic Fever, Democratic Republic of the Congo, 1995 \[PDF – 361KB\]](#) [↗](#). *Journal of Infectious Diseases*. 1999;179:S76–S86.

Le Guenno B, Formenty P, Wyers M, et al. [Isolation and partial characterisation of a new strain of Ebola virus](#) [↗](#). *Lancet*. 1995;345:1271–1274.

MacNeil A, Farnon EC, Morgan OW, et al. [Filovirus Outbreak Detection and Surveillance: Lessons from Bundibugyo \[PDF – 203KB\]](#) [\[PDF\]](#). *Journal of Infectious Diseases*. 2011;204:S761–S767.

Maganga GD, Kapetshi J., Berthet N, et al. [Ebola virus disease in the Democratic Republic of Congo \[PDF – 1.91MB\]](#) [↗](#). *New England Journal of Medicine*. 2014; 371: 2083–91.

Milleliri JM, Tévi-Benissan C, Baize S, et al. [Les épidémies de fièvre hémorragique due au virus Ebola au Gabon \(1994–2002\): Aspects épidémiologiques et réflexions sur les mesures de contrôle](#) [\[PDF\]](#) [↗](#). *Bull Soc Pathol Exot*, 2004, 97, 3, 199–205.

Miranda ME, White ME, Dayrit MM, Hayes CG, Ksiazek TG, and Burans JP. [Seroepidemiological study of filovirus related to Ebola in the Philippine](#) [↗](#)s. *Lancet*. 1991;337:425–426.

Okware SI, Omaswa FG, Zaramba S, et al. [An outbreak of Ebola in Uganda](#) [↗](#). *Tropical Medicine and International Health*. 2002;7(12):1068–1075.

Rollin PE, Williams J, Bressler D, et al. [Isolated cases of Ebola \(subtype Reston\) virus among quarantined non-human primates recently imported from the Philippines to the United States](#) [↗](#). *Journal of Infectious Diseases*. 1999;179 (suppl 1):S108–S114.

Shoemaker T, MacNeil A, Balinandi S, et al. [Reemerging Sudan Ebola Virus Disease in Uganda, 2011](#). *Emerging Infectious Diseases*. 2012;18(9):1480–1483.

World Health Organization. [Ebola Virus Disease—Italy](#). *Disease Outbreak News* [↗](#). 13 May 2015.



World Health Organization. [Ebola Virus Disease—Spain](#) [↗](#). 4 October 2014.

World Health Organization. [WHO Congratulates Spain on Ending Ebola Transmission](#) [↗](#). 2 December 2014.

World Health Organization. [Ebola Virus Disease—United Kingdom](#) [↗](#). *Disease Outbreak News*. 30 December 2014.

World Health Organization. [End of the Ebola Outbreak in the Democratic Republic of the Congo](#) [↗](#). *Global Alert and Response*. 17 February 2009.


World Health Organization. [Ebola Reston in pigs and humans, Philippines \[PDF – 240KB\]](#) [\[PDF\]](#) [↗](#). *Weekly Epidemiological Record*. 2009;84(7):49–50.


World Health Organization. [Ebola virus haemorrhagic fever, Democratic Republic of the Congo—Update. \[PDF -373KB\]](#)   *Weekly Epidemiological Record*. 2007;82(40):345–346.


World Health Organization. [Outbreak of Ebola haemorrhagic fever in Yambio, south Sudan, April–June 2000 \[159 KB, 8 pages\]](#)  . *Weekly Epidemiological Record*. 2005;80(43):370–375.



World Health Organization. [Ebola haemorrhagic fever in the Republic of the Congo—Update 6](#) . *Weekly Epidemiological Record*. 6 January 2004.

World Health Organization. [Outbreak\(s\) of Ebola haemorrhagic fever, Congo and Gabon, October 2001–July 200 \[518 KB, 12 pages\]](#)  . *Weekly Epidemiological Report*. 2003;78(26):223–225.

World Health Organization. [Ebola haemorrhagic fever—South Africa \[PDF-469KB\]](#)  . *Weekly Epidemiological Record*. 1996;71(47):359.

World Health Organization. [Viral haemorrhagic fever in imported monkeys \[PDF-799KB\]](#) . *Weekly Epidemiological Record*. 1992;67(24):183.

World Health Organization. [Ebola haemorrhagic fever in Sudan, 1976. Report of a WHO/International Study Team \[PDF – 5.91MB\]](#)  . *Bulletin of the World Health Organization*. 1978;56(2):247–270.

World Health Organization. [Ebola haemorrhagic fever in Zaire, 1976 \[PDF – 3.05MB\]](#)  . Report of an International Commission. *Bulletin of the World Health Organization*. 1978;56(2):271–293.

READ NEXT

[How Ebola Disease Spreads](#)



MAY 6, 2024

⊖ SOURCES

SHARE

CONTENT SOURCE:

[National Center for Emerging and Zoonotic Infectious Diseases \(NCEZID\)](#)

How helpful was this page?



Not helpful

Very helpful

RELATED PAGES

[Ebola Disease Basics](#)

[Signs and Symptoms of Ebola Disease](#)

[How Ebola Disease Spreads](#)

[Ebola Disease Outbreak Locations](#)