

SUPPLEMENTARY TABLE 1. Self-reported substance use and sexual behavior characteristics in the 3 months preceding emergency department visit among 13 male participants with an mpox-compatible rash who identified as gay, bisexual, or queer— 13 U.S. sites, June–December 2023

Characteristic	n/N* (%)
Alcohol and substance use	
Alcohol bingeing	1/13 (7.7)
Smoked cigarettes, vaped, or chewed tobacco	5/13 (38.5)
Used cannabis or tetrahydrocannabinol	8/13 (61.5)
Injected drugs	2/13 (15.4)
Non-injection stimulant use	3/13 (23.1)
Non-injection opioid use	1/13 (7.7)
Amyl nitrate (“popper”) use	3/13 (23.1)
None of the above	2/13 (15.4)
Living with a male who has sex with men or is transgender who has sex with men	5/13 (38.5)
Sexual activity in the last 3 months	8/13 (61.5)
Number of sexual partners	
1	1/8 (12.5)
2	2/8 (25.0)
3	1/8 (12.5)
5	2/8 (25.0)
Declined to answer	2/8 (25.0)
Had sex with someone who had similar symptoms or rash	
Yes	2/8 (25.0)
No	4/8 (50.0)
Don’t know	2/8 (25.0)
Condom use	
Never	3/8 (37.5)
Sometimes	4/8 (50.0)
Always	1/8 (12.5)
Engaged in any anal sex	5/8 (62.5)
Engaged in any oral sex	7/8 (87.5)
Sexual history	
Group sex	3/8 (37.5)
Sex parties	2/8 (25.0)
Met sex partners through dating apps	5/8 (62.5)
Given someone money for sex	0/8 (0.0)
Got paid for sex	1/8 (12.5)
Sex after taking drugs	2/8 (25.0)
Sex at a music festival or rave	0/8 (0.0)
Sex with anonymous partner	3/8 (37.5)
Shared sex toys	1/8 (12.5)
Traveled and had sex with new partner	0/8 (0.0)
Sex with visiting partner from another country	0/8 (0.0)
Sex with someone from out of town	2/8 (25.0)
None of the above	1/8 (12.5)