



Weekly U.S. Influenza Surveillance Report

Updated June 28, 2024

FLUVIEW



A Weekly Influenza Surveillance Report Prepared by the Influenza Division

Key Updates for Week 25, ending June 22, 2024

Seasonal influenza activity remains low nationally.

Viruses

Clinical Lab

1.1% (Trend →)
positive for influenza
this week

[\(/flu/weekly/index.htm#ClinicalLaboratories\)](/flu/weekly/index.htm#ClinicalLaboratories)

Public Health Lab

Influenza A(H1N1)pdm09,
A(H3N2), and B viruses were all
co-circulating this week.
[\(/flu/weekly/index.htm#PublicHealthLaboratories\)](/flu/weekly/index.htm#PublicHealthLaboratories)

Virus Characterization

Genetic and antigenic
characterization and antiviral
susceptibility are summarized in
this report.
[\(/flu/weekly/index.htm#VirusCharacterization\)](/flu/weekly/index.htm#VirusCharacterization)

Illness

Outpatient Respiratory Illness

1.5% (Trend →)
of visits to a health care provider this week were for
respiratory illness
(below baseline).
[\(/flu/weekly/index.htm#ILINet\)](/flu/weekly/index.htm#ILINet)

Outpatient Respiratory Illness: Map

This week no jurisdictions experienced moderate,
high, or very high activity.
[\(/flu/weekly/index.htm#ORIAM\)](/flu/weekly/index.htm#ORIAM)

FluSurv-NET

0.2 per 100,000

weekly hospitalization rate.

[\(/flu/weekly/index.htm#FluSurvNet\)](/flu/weekly/index.htm#FluSurvNet)

NCHS Mortality

0.1% (Trend →)

of deaths attributed to influenza this week.

[\(/flu/weekly/index.htm#NCHSMortality\)](/flu/weekly/index.htm#NCHSMortality)

Pediatric Deaths

3

influenza-associated deaths were reported this week
for a total of 181 deaths this season.

[\(/flu/weekly/index.htm#PedMortality\)](/flu/weekly/index.htm#PedMortality)

All data are preliminary and may change as more reports are received.

Directional arrows indicate changes between the current week and the previous week. [Additional information \(/flu/weekly/index.htm#Trends\)](/flu/weekly/index.htm#Trends) on the arrows can be found at the bottom of this page.

A description of the CDC influenza surveillance system, including methodology and detailed descriptions of each data component is available on the [surveillance methods \(/flu/weekly/overview.htm\)](/flu/weekly/overview.htm) page.

Additional information on the current and previous influenza seasons for each surveillance component are available on [FluView Interactive \(/flu/weekly/fluviewinteractive.htm\)](/flu/weekly/fluviewinteractive.htm).

Key Points

- Seasonal influenza activity remains low nationally.
- CDC estimates that there have been at least 35 million illnesses, 390,000 hospitalizations, and 25,000 deaths from flu so far this season.
- There are prescription flu antiviral drugs that can treat flu illness; those should be started as early as possible and are especially important for higher risk patients.³
- Seasonal flu viruses are among several viruses contributing to respiratory disease activity. CDC is providing updated, integrated information (<https://www.cdc.gov/respiratory-viruses/index.html>) about COVID-19, flu, and RSV activity on a weekly basis.

U.S. Virologic Surveillance

[\(/https://www.cdc.gov/flu/weekly/overview.htm#LabSurveillance\)](https://www.cdc.gov/flu/weekly/overview.htm#LabSurveillance)

Nationally, the percentage of respiratory specimens testing positive for influenza in clinical laboratories remained stable (change of ≤ 0.5 percentage points) compared to the previous week. Nationally, influenza A(H1N1)pdm09, A(H3N2), and B/Victoria viruses are all co-circulating. However, the distribution of circulating viruses varies by region. For regional and state level data and age group distribution, please visit [FluView Interactive \(/https://gis.cdc.gov/grasp/fluview/fluportaldashboard.html\)](https://gis.cdc.gov/grasp/fluview/fluportaldashboard.html).


Clinical Laboratories

The results of tests performed by clinical laboratories nationwide are summarized below. Data from clinical laboratories (the percentage of specimens tested that are positive for influenza virus) are used to monitor whether influenza activity is increasing or decreasing.

	Week 25	Data Cumulative since October 1, 2023 (Week 40)
No. of specimens tested	31,854	3,457,891
No. of positive specimens (%)	352 (1.1%)	348,428 (10.1%)
<i>Positive specimens by type</i>		
Influenza A	283 (80.4%)	240,427 (69.0%)
Influenza B	69 (19.6%)	107,990 (31.0%)

Season: 2023-24 Surveillance Area: National

Influenza Positive Tests Reported to CDC by Clinical Laboratories, National Summary, 2023-24 Season, week ending Jun 22, 2024



View Additional Graphs and Data (<http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>) | Download Chart Data | Download PowerPoint Presentation


Public Health Laboratories

The results of tests performed by public health laboratories nationwide are summarized below. Data from public health laboratories are used to monitor the proportion of circulating influenza viruses that belong to each influenza subtype/lineage.

	Week 25	Data Cumulative since October 1, 2023 (Week 40)
No. of specimens tested	688	114,661
No. of positive specimens	101	37,649
<i>Positive specimens by type/subtype</i>		
Influenza A	82 (81.2%)	28,641 (76.1%)
Subtyping Performed	56 (68.3%)	24,147 (84.3%)
(H1N1)pdm09	21 (37.5%)	16,175 (67.0%)
H3N2	35 (62.5%)	7,972 (33.0%)
H3N2v	0 (0.0%)	0 (0.0%)
Subtyping not performed	26 (31.7%)	4,494 (15.7%)
Influenza B	19 (18.8%)	9,008 (23.9%)
Lineage testing performed	12 (63.2%)	7,853 (87.2%)
Yamagata lineage	0 (0.0%)	0 (0.0%)
Victoria lineage	12 (100.0%)	7,853 (100.0%)
Lineage not performed	7 (36.8%)	1,155 (12.8%)

Season: 2023-24 Surveillance Area: National

Influenza Positive Tests Reported to CDC by Public Health Laboratories, National Summary, 2023-24 Season, week ending Jun 22, 2024



Check All A (H3) A (H1N1)pdm09 A (Subtyping not Performed) B (Lineage Unspecified) B (Victoria Lineage)

View Additional Graphs and Data (<http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>) | Download Chart Data | Download PowerPoint Presentation

Additional virologic surveillance information for current and past seasons:

[Surveillance Methods \(flu/weekly/overview.htm#LabSurveillance\)](#) | [FluView Interactive: National, Regional, and State Data \(http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html\)](#) or [Age Data \(https://gis.cdc.gov/grasp/fluview/flu_by_age_virus.html\)](#)

Novel Influenza A Virus:

Two human infections with a novel influenza A virus were reported by the Pennsylvania Department of Health. The patients, who are close contacts, were both infected with influenza A(H1N2) variant (A(H1N2)v) viruses. Both patients are ≥ 18 years of age and sought healthcare during the week ending June 22, 2024 (Week 25). One of the patients was hospitalized, and one was not. An investigation by state public health officials found that the patients had attended a livestock auction where swine were present prior to their illness onset. Investigation did not identify illness among additional close contacts of either patient. The investigation is ongoing.

Including these two reports there have been a total of three human infections with variant influenza A viruses reported in the United States in 2024.

When an influenza virus that normally circulates in swine (but not people) is detected in a person, it is called a “variant” influenza virus. Most human infections with variant influenza viruses occur following exposure to swine, but human-to-human transmission can occur. It is important to note that in most cases, variant influenza viruses have not shown the ability to spread easily and sustainably from person to person.

Early identification and investigation of human infections with novel influenza A viruses are critical so that the risk of infection can be understood, and appropriate public health measures can be taken.

Additional information on influenza in swine, variant influenza virus infection in humans, and guidance to interact safely with swine can be found at www.cdc.gov/flu/swineflu/index.htm (<http://www.cdc.gov/flu/swineflu/index.htm>).

Additional information regarding human infections with novel influenza A viruses:

Surveillance Methods (<https://www.cdc.gov/flu/weekly/overview.htm#NovelASurveillance>) | FluView Interactive (http://gis.cdc.gov/grasp/fluview/Novel_Influenza.html)

Influenza Virus Characterization ([/flu/weekly/overview.htm#VirusCharacterization](https://www.cdc.gov/flu/weekly/overview.htm#VirusCharacterization))

CDC performs [genetic](https://www.cdc.gov/flu/about/professionals/genetic-characterization.htm) (<https://www.cdc.gov/flu/about/professionals/genetic-characterization.htm>) and [antigenic](https://www.cdc.gov/flu/about/professionals/antigenic.htm) (<https://www.cdc.gov/flu/about/professionals/antigenic.htm>) characterization of U.S. viruses submitted from state and local public health laboratories according to the Right Size Roadmap submission guidance. These data are used to compare how similar the currently circulating influenza viruses are to the reference viruses representing viruses contained in the current influenza vaccines. The data are also used to monitor evolutionary changes that continually occur in influenza viruses circulating in humans. CDC also tests susceptibility of circulating influenza viruses to antiviral medications including the neuraminidase inhibitors (oseltamivir, zanamivir, and peramivir) and the PA endonuclease inhibitor baloxavir.

CDC has genetically characterized 4,922 influenza viruses collected since October 1, 2023.

Virus Subtype or Lineage	Genetic Characterization				
	Total No. of Subtype/Lineage Tested	HA Clade	Number (% of subtype/lineage tested)	HA Subclade	Number (% of subtype/lineage tested)
A/H1	1,828				
		6B.1A.5a	1,828 (100%)	2a	423 (23.1%)
				2a.1	1,405 (76.9%)
A/H3	1,681				
		3C.2a1b.2a	1,681 (100%)	2a.1b	1 (0.1%)
				2a.3a	1 (0.1%)
				2a.3a.1	1,678 (99.8%)
				2b	1 (0.1%)
B/Victoria	1,413				
		V1A	1,413 (100%)	3a.2	1,413 (100%)
B/Yamagata	0				
		Y3	0	Y3	0 (0%)

CDC [antigenically characterizes](https://www.cdc.gov/flu/about/professionals/antigenic.htm) (<https://www.cdc.gov/flu/about/professionals/antigenic.htm>) influenza viruses by hemagglutination inhibition (HI) (H1N1pdm09, H3N2, B/Victoria, and B/Yamagata viruses) or neutralization-based [HINT](https://pubmed.ncbi.nlm.nih.gov/30804469/) [HINT](https://pubmed.ncbi.nlm.nih.gov/30804469/) <https://pubmed.ncbi.nlm.nih.gov/30804469/>) (H3N2 viruses) using antisera that ferrets make after being infected with reference viruses representing the 2023-2024 Northern Hemisphere recommended cell or recombinant-based vaccine viruses. Antigenic differences between viruses are determined by comparing how well the antibodies made against the vaccine reference viruses recognize the circulating viruses that have been grown in cell culture. Ferret antisera are useful because antibodies raised against a particular virus can often recognize small changes in the surface proteins of other viruses. In HI assays, viruses with similar antigenic properties have antibody titer differences of less than or equal to 4-fold when compared to the reference (vaccine) virus. In HINT, viruses with similar antigenic properties have antibody neutralization titer differences of less than or equal to 8-fold. Viruses selected for antigenic characterization are a subset representing the genetic changes in the surface proteins seen in genetically characterized viruses.

Influenza A Viruses

- **A (H1N1)pdm09:** 458 A(H1N1)pdm09 viruses were antigenically characterized by HI, and 457 (99.8%) were well-recognized (reacting at titers that were within 4-fold of the homologous virus titer) by ferret antisera to cell-grown A/Wisconsin/67/2022-like reference viruses representing the A(H1N1)pdm09 component for the cell- and recombinant-based influenza vaccines.
- **A (H3N2):** 537 A(H3N2) viruses were antigenically characterized by HI or HINT, and 519 (96.6%) were well-recognized (reacting at titers that were within 4-fold of the homologous virus titer in HI or reacting at titers that were less than or equal to 8-fold of the homologous virus in HINT) by ferret antisera to cell-grown A/Darwin/6/2021-like reference viruses representing the A(H3N2) component for the cell- and recombinant-based influenza vaccines.

Influenza B Viruses

- **B/Victoria:** 342 influenza B/Victoria-lineage virus were antigenically characterized by HI, and all were well-recognized (reacting at titers that were within 4-fold of the homologous virus titer) by ferret antisera to cell-grown B/Austria/1359417/2021-like reference viruses representing the B/Victoria component for the cell- and recombinant-based influenza vaccines.
- **B/Yamagata:** No influenza B/Yamagata-lineage viruses were available for antigenic characterization.

Assessment of Virus Susceptibility to Antiviral Medications

CDC assesses susceptibility of influenza viruses to the antiviral medications including the neuraminidase inhibitors (oseltamivir, zanamivir, and peramivir) and the PA endonuclease inhibitor baloxavir using next generation sequence analysis supplemented by laboratory assays. Information about antiviral susceptibility test methods can be found at [U.S. Influenza Surveillance: Purpose and \(https://www.cdc.gov/flu/weekly/overview.htm\) Methods | CDC \(https://www.cdc.gov/flu/weekly/overview.htm\).](https://www.cdc.gov/flu/weekly/overview.htm)

Viruses collected in the U.S. since October 1, 2023, were tested for antiviral susceptibility as follows:

Antiviral Medication			Total Viruses	A/H1	A/H3	B/Victoria
Neuraminidase Inhibitors	Oseltamivir	Viruses Tested	4,843	1,807	1,653	1,383
		Reduced Inhibition	1 (0.02%)	1 (0.1%)	0 (0.00%)	0 (0.00%)
		Highly Reduced Inhibition	4 (0.1%)	4 (0.1%)	0 (0.00%)	0 (0.00%)
	Peramivir	Viruses Tested	4,843	1,807	1,653	1,383
		Reduced Inhibition	3 (0.1%)	0 (0.00%)	0 (0.00%)	3 (0.2%)
		Highly Reduced Inhibition	4 (0.1%)	4 (0.1%)	0 (0.00%)	0 (0.00%)
	Zanamivir	Viruses Tested	4,843	1,807	1,653	1,383
		Reduced Inhibition	1 (0.02%)	0 (0.00%)	0 (0.00%)	1 (0.1%)
		Highly Reduced Inhibition	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
PA Cap-Dependent Endonuclease Inhibitor	Baloxavir	Viruses Tested	4,765	1,764	1,630	1,371
		Decreased Susceptibility	1 (0.02%)	0 (0.0%)	1 (0.1%)	0 (0.0%)

Three A(H1N1)pdm09 viruses had NA-H275Y amino acid substitution and one A(H1N1)pdm09 virus had NA-H275H/Y, conferring highly reduced inhibition by oseltamivir and peramivir. One (H1N1)pdm09 virus had NA-I223V and NA-S247N amino acid substitutions and showed reduced inhibition by oseltamivir. Two B viruses had NA-A245G

amino acid substitution and showed reduced inhibition by peramivir. One B virus had NA-D197N amino acid substitution and showed reduced inhibition by zanamivir and peramivir. One A(H3N2) virus had PA-I38T amino acid substitution and showed reduced susceptibility to baloxavir.

High levels of resistance to the adamantanes (amantadine and rimantadine) persist among influenza A(H1N1)pdm09 and influenza A(H3N2) viruses (the adamantanes are not effective against influenza B viruses). Therefore, use of these antivirals for treatment and prevention of influenza A virus infection is not recommended and data from adamantane resistance testing are not presented.

Outpatient Respiratory Illness Surveillance (<https://www.cdc.gov/flu/weekly/overview.htm#ILINet>)


The U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet) monitors outpatient visits for respiratory illness referred to as influenza-like illness [ILI (fever plus cough or sore throat)], not laboratory-confirmed influenza and will therefore capture respiratory illness visits due to infection with pathogens that can present with similar symptoms, including influenza viruses, SARS-CoV-2, and RSV. It is important to evaluate syndromic surveillance data, including that from ILINet, in the context of other sources of surveillance data to obtain a more complete and accurate picture of influenza, SARS-CoV-2, and other respiratory virus activity. CDC is providing integrated information about COVID-19, influenza, and RSV activity on a [website \(https://www.cdc.gov/respiratory-viruses/index.html\)](https://www.cdc.gov/respiratory-viruses/index.html) that is updated weekly. Information about other respiratory virus activity can be found on [CDC's National Respiratory and Enteric Virus Surveillance System \(NREVSS\) website \(https://www.cdc.gov/surveillance/nrevss/index.html\)](https://www.cdc.gov/surveillance/nrevss/index.html).

Outpatient Respiratory Illness Visits

Nationally, the percentage of visits for respiratory illness that were reported through ILINet remained stable (change of ≤ 0.1 percentage points) compared to the previous week and is below the national baseline. All 10 regions are below their region-specific baselines. Multiple respiratory viruses are co-circulating, and the relative contribution of influenza virus infection to ILI varies by location.

Season: 2023-24 and 5 previous seasons ▾

**Percentage of Outpatient Visits for Respiratory Illness Reported by
The U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet),
Weekly National Summary, 2023-24 Season and Selected Previous Seasons**



- 2023-24 National Baseline
- 2023-24 Season
- 2022-23 Season
- 2021-22 Season
- 2020-21 Season
- 2019-20 Season
- 2018-19 Season

View National and Regional Level Graphs and Data (<https://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>) | Download Chart Data | Download PowerPoint Presentation


Outpatient Respiratory Illness Visits by Age Group

About 70% of ILINet participants provide both the number of patient visits for respiratory illness and the total number of patient visits for the week broken out by age group. Data from this subset of providers are used to calculate the percentages of patient visits for respiratory illness by age group.

The percentage of visits for respiratory illness reported in ILINet decreased in the 0-4 years and 5-24 years age groups and remained stable for all other age groups in Week 25 compared to Week 24.

Season: 2023-24

Percent of Outpatient Visits for Respiratory Illness by Age Group
 Reported by the U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet)
 Weekly National Summary, 2023-24 Influenza Season through the Week Ending June 22, 2024



View FluView Interactive (https://gis.cdc.gov/grasp/fluview/flu_by_age_virus.html) | Download Chart Data | Download PowerPoint Presentation

Outpatient Respiratory Illness Activity Map

Data collected in ILINet are used to produce a measure of ILI activity* ([/flu/weekly/overview.htm#ILINet](#)) by state/jurisdiction and Core Based Statistical Areas (CBSA).


	Number of Jurisdictions		Number of CBSAs	
	Week 25 (Week ending	Week 24 (Week ending	Week 25 (Week ending	Week 24 (Week ending

Activity Level	Jun. 22, 2024)	Jun. 15, 2024)	Jun. 22, 2024)	Jun. 15, 2024)
Very High	0	0	0	0
High	0	0	0	0
Moderate	0	0	1	3
Low	1	1	5	5
Minimal	53	54	676	676
Insufficient Data	1	0	247	245


A Weekly Influenza Surveillance Report Prepared by the Influenza Division

Outpatient Respiratory Illness Activity Map Determined by Data Reported to ILINet

This system monitors visits for respiratory illness that includes fever plus a cough or sore throat, also referred to as ILI, not laboratory confirmed influenza and may capture patient visits due to other respiratory pathogens that cause similar symptoms.



2023-24 Influenza Season Week 25 ending Jun 22, 2024



Season: 2023-24 ▲

Download Image Download Data

(<https://www.cdc.gov/flu/weekly/fluereport.xml>)View Full Screen (<http://gis.cdc.gov/grasp/fluview/main.html>)

*Data collected in ILINet may disproportionately represent certain populations within a jurisdiction or CBSA, and therefore, may not accurately depict the full picture of influenza activity for the entire jurisdiction or CBSA. Differences in the data presented here by CDC and independently by some health departments likely represent differing levels of data completeness with data presented by the health department likely being the more complete.

Additional information about medically attended visits for ILI for current and past seasons:

[Surveillance Methods \(/flu/weekly/overview.htm#ILINet\)](#) | [FluView Interactive: National, Regional, and State Data \(http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html\)](#) or [ILI Activity Map \(https://gis.cdc.gov/grasp/fluview/main.html\)](#)

Hospitalization Surveillance

(<http://www.cdc.gov/flu/weekly/overview.htm#HospitalizationSurv>)

FluSurv-NET

The Influenza Hospitalization Surveillance Network (FluSurv-NET) conducts population-based surveillance for laboratory-confirmed influenza-related hospitalizations in select counties in 14 states and represents approximately 9% of the U.S. population. FluSurv-NET hospitalization data are preliminary. As data are received each week, prior case counts and rates are updated accordingly.


A total of 25,155 laboratory-confirmed influenza-associated hospitalizations were reported by FluSurv-NET sites between October 1, 2023, and June 22, 2024. The weekly hospitalization rate observed in Week 25 was 0.2 per 100,000 population. The peak weekly hospitalization rate observed this season was 9.0 per 100,000 population and occurred during Week 52.

Among 25,155 hospitalizations, 21,273 (84.6%) were associated with influenza A virus, 3,696 (14.7%) with influenza B virus, 52 (0.2%) with influenza A virus and influenza B virus co-infection, and 133 (0.5%) with influenza virus for which the type was not determined. Among those with influenza A subtype information, 4,112 (67.9%) were A(H1N1)pdm09 and 1,944 (32.1%) were A(H3N2).

Age Group: Overall | Display By | Select a Surveillance Area ?

Laboratory-Confirmed Influenza Hospitalizations
Preliminary cumulative rates as of Jun 22, 2024

FluSurv-NET :: Entire Network :: Overall Age Group :: Cumulative Rate



- All Seasons
- 2023-24
- 2022-23
- 2021-22
- 2020-21
- 2019-20
- 2018-19
- 2017-18
- 2016-17
- 2015-16
- 2014-15
- 2013-14
- 2012-13
- 2011-12
- 2010-11
- 2009-10

View National and Regional Level Graphs and Data (<https://gis.cdc.gov/grasp/fluview/FluHospRates.html>) | Download Chart Data | Download PowerPoint Presentation

**In this figure, weekly rates for all seasons prior to the 2023-2024 season reflect end-of-season rates. For the 2023-2024 season, rates for recent hospital admissions are subject to reporting delays and are shown as a dashed line for the current season. As hospitalization data are received each week, prior case counts and rates are updated accordingly.

Additional FluSurv-NET hospitalization surveillance information for current and past seasons and additional age groups:

Surveillance Methods (<https://www.cdc.gov/flu/weekly/overview.htm#FluSurvNet>) | FluView Interactive: Rates by Age, Sex, and Race/Ethnicity (<http://gis.cdc.gov/GRASP/Fluview/FluHospRates.html>) or Data on Patient Characteristics (<http://gis.cdc.gov/grasp/fluview/FluHospChars.html>) | RESP-NET Interactive (<https://www.cdc.gov/surveillance/resp-net/dashboard.html>)

National Healthcare Safety Network (NHSN) Hospitalization Surveillance

Effective May 1, 2024, hospitals are no longer required to report hospital admissions, hospital capacity, or hospital occupancy data to HHS through NHSN. Voluntarily reported NHSN hospital data can found at [Weekly United States Hospitalization Metrics by Jurisdiction](https://data.cdc.gov/Public-Health-Surveillance/Weekly-United-States-Hospitalization-Metrics-by-Ju/aemt-mg7g/about_data) (https://data.cdc.gov/Public-Health-Surveillance/Weekly-United-States-Hospitalization-Metrics-by-Ju/aemt-mg7g/about_data).

Additional NHSN Hospitalization Surveillance information:

[Surveillance Methods](https://www.cdc.gov/flu/weekly/overview.htm#NHSN) (<https://www.cdc.gov/flu/weekly/overview.htm#NHSN>) | [Additional Data](https://data.cdc.gov/Public-Health-Surveillance/Weekly-United-States-Hospitalization-Metrics-by-Ju/aemt-mg7g/about_data) (https://data.cdc.gov/Public-Health-Surveillance/Weekly-United-States-Hospitalization-Metrics-by-Ju/aemt-mg7g/about_data) | [FluView Interactive](http://gis.cdc.gov/grasp/fluview/FluView12.html) (<http://gis.cdc.gov/grasp/fluview/FluView12.html>)

Mortality Surveillance

(<https://www.cdc.gov/flu/weekly/overview.htm#MortalitySurveillance>)

National Center for Health Statistics (NCHS) Mortality Surveillance


Based on NCHS mortality surveillance data available on June 27, 2024, the percentage of deaths that were due to influenza remained stable (< 0.1 percentage point change) compared to the previous week. The data presented are preliminary and may change as more data are received and processed.

Season: 2023-24 Surveillance Area: National Age: All

Pneumonia and Influenza Mortality from the National Center for Health Statistics Mortality Surveillance System

National Summary data through the week ending June 22, 2024

2019-24 Show Number of Influenza Deaths



View Regional and State Level Data (<https://gis.cdc.gov/grasp/fluview/Mortality.html>) | Download Chart Data | Download PowerPoint Presentation

<

Additional pneumonia, influenza and COVID-19 mortality surveillance information for current and past seasons:

Surveillance Methods (<https://www.cdc.gov/flu/weekly/overview.htm#NCHSMortality>) | FluView Interactive

(<https://gis.cdc.gov/grasp/fluview/mortality.html>)


Influenza-Associated Pediatric Mortality

Three influenza-associated pediatric deaths occurring during the 2023-2024 season were reported to CDC during Week 25. Two deaths were associated with influenza A viruses with no subtyping performed. One of the deaths occurred during Week 51 (the week ending December 23, 2023) and the other occurred during Week 14 (the week ending April 6, 2024). The third death occurred during Week 20 (the week ending May 18, 2024) and was associated with an influenza B/Victoria virus.

A total of 181 influenza-associated pediatric deaths occurring during the 2023-2024 season have been reported to CDC.

Influenza-Associated Pediatric Deaths by Week of Death, 2020-21 season to 2023-24 season

Select up to 4 Seasons ▼ Group By Reported Virus Type
 By Week Current Week Previous Weeks All Deaths



[View FluView Interactive \(https://gis.cdc.gov/grasp/fluview/PedFluDeath.html\)](https://gis.cdc.gov/grasp/fluview/PedFluDeath.html) | [Download Chart Data](#) | [Download PowerPoint Presentation](#)

Additional pediatric mortality surveillance information for current and past seasons:

[Surveillance Methods \(https://www.cdc.gov/flu/weekly/overview.htm#PediatricMortality\)](https://www.cdc.gov/flu/weekly/overview.htm#PediatricMortality) | [FluView Interactive \(https://gis.cdc.gov/GRASP/Fluview/PedFluDeath.html\)](https://gis.cdc.gov/GRASP/Fluview/PedFluDeath.html)

Trend Indicators

Increasing: ↑
Decreasing: ↓
Stable: →

Indicators Status by System

Clinical Labs: Up or down arrows indicate a change of greater than or equal to 0.5 percentage points in the percent of specimens positive for influenza compared to the previous week.

Outpatient Respiratory Illness (ILINet): Up or down arrows indicate a change of greater than 0.1 percentage points in the percent of visits due to respiratory illness (ILI) compared to the previous week.

NHSN Hospitalizations: Up or down arrows indicate change of greater than or equal to 5% of the number of patients admitted with laboratory-confirmed influenza compared to the previous week.

NCHS Mortality: Up or down arrows indicate change of greater than 0.1 percentage points of the percent of deaths due to influenza compared to the previous week.

Reference Footnotes

¹U.S. Influenza Surveillance: Purpose and Methods (2023 Oct). Centers for Disease Control and Prevention.
<https://www.cdc.gov/flu/weekly/overview.htm#ILINet> (<https://www.cdc.gov/flu/weekly/overview.htm#ILINet>).

²Grohskopf LA, Blanton LH, Ferdinands JM, Chung JR, Broder KR, Talbot HK. Prevention and Control of Seasonal Influenza with Vaccines: Recommendations of the Advisory Committee on Immunization Practices — United States, 2023–24 Influenza Season. MMWR Recomm Rep 2023;72(No. RR-2):1–25.
DOI: <http://dx.doi.org/10.15585/mmwr.rr7202a1>  (<http://dx.doi.org/10.15585/mmwr.rr7202a1>)

³Influenza Antiviral Medications: Summary for Clinicians (2023 Sept). Centers for Disease Control and Prevention.
<https://www.cdc.gov/flu/professionals/antivirals/summary-clinicians.htm> (<https://www.cdc.gov/flu/professionals/antivirals/summary-clinicians.htm>).

Additional National and International Influenza Surveillance Information

FluView Interactive: FluView includes enhanced web-based interactive applications that can provide dynamic visuals of the influenza data collected and analyzed by CDC. These [FluView Interactive applications](http://www.cdc.gov/flu/weekly/fluviewinteractive.htm) (<http://www.cdc.gov/flu/weekly/fluviewinteractive.htm>) allow people to create customized, visual interpretations of influenza data, as well as make comparisons across flu seasons, regions, age groups and a variety of other demographics.

National Institute for Occupational Safety and Health: Monthly surveillance data on the prevalence of health-related workplace absenteeism among full-time workers in the United States are [available from NIOSH](https://www.cdc.gov/niosh/topics/absences/default.html) (<https://www.cdc.gov/niosh/topics/absences/default.html>).

U.S. State and local influenza surveillance: Select a jurisdiction below to access the latest local influenza information.

[Alabama](http://adph.org/influenza/) (<http://adph.org/influenza/>)

[Alaska](http://dhss.alaska.gov/dph/Epi/id/Pages/influenza/fluinf)

(<http://dhss.alaska.gov/dph/Epi/id/Pages/influenza/fluinf>)

Colorado (<https://www.mass.gov/info-details/influenza-reporting>)

Connecticut (<https://portal.ct.gov/DPH/Epidemiology-Emerging-Infections/Influenza-Surveillance-and-Statist>)

Georgia (<https://dph.georgia.gov/flu-activity-georgia>)

Hawaii
(<http://health.hawaii.gov/docd/resources/reports/influenza-reports/>)

Iowa (<https://idph.iowa.gov/influenza/reports>)

Kansas (<http://www.kdheks.gov/flu/surveillance.htm>)

Maryland (<https://phpa.health.maryland.gov/influenza/fluwatch/>)

Massachusetts (<https://www.mass.gov/info-details/influenza-reporting>)

Missouri
(<http://health.mo.gov/living/healthcondiseases/communicable/influenza/reports.php>)

Montana
(<https://dphhs.mt.gov/publichealth/cdepi/diseases/Pan-Respiratory/index>)

New Jersey (<http://www.nj.gov/health/cd/topics/flu.shtml>)

New Mexico (<https://nmhealth.org/about/erd/ideb/isp>)

Ohio (<http://www.flu.ohio.gov>)

Oklahoma (<https://oklahoma.gov/health/health-education/acute-disease-service/disease-information/influenza-home-page.html>)

South Carolina
(<http://www.scdhec.gov/Health/DiseasesandConditions/InfectiousDiseases/Flu/FluData/>)

South Dakota (<https://doh.sd.gov/health-data-reports/data-dashboards/influenza-dashboard/>)

Vermont (<http://www.healthvermont.gov/immunizations-infectious-disease/influenza/flu-activity-and-surveillance>)

Virginia
(<http://www.vdh.virginia.gov/epidemiology/influenza-flu-virginia/influenza-surveillance/>)

Wyoming (<https://health.wyo.gov/publichealth/infectious-disease-epidemiology-unit/disease/influenza/>)

New York City
(<http://www1.nyc.gov/site/doh/providers/health-topics/flu-alerts.page>)

World Health Organization:

Additional influenza surveillance information from participating WHO member nations is available through FluNet [🔗](https://www.who.int/tools/flunet) (<https://www.who.int/tools/flunet>) and the Global Epidemiology Reports. [🔗](https://www.who.int/teams/global-influenza-programme/surveillance-and-monitoring/influenza-surveillance-outputs) (<https://www.who.int/teams/global-influenza-programme/surveillance-and-monitoring/influenza-surveillance-outputs>)

WHO Collaborating Centers for Influenza:

Australia [🔗](http://www.influenzacentre.org/Surveillance_Samples_Received.html) (http://www.influenzacentre.org/Surveillance_Samples_Received.html), China [🔗](http://www.chinaivdc.cn/cnic/) (<http://www.chinaivdc.cn/cnic/>), Japan [🔗](http://idsc.nih.go.jp/index.html) (<http://idsc.nih.go.jp/index.html>), the United Kingdom [🔗](https://www.crick.ac.uk/research/worldwide-influenza-centre) (<https://www.crick.ac.uk/research/worldwide-influenza-centre>), and the United States (<http://www.cdc.gov/flu/>) (CDC in Atlanta, Georgia)

Europe:

The most up-to-date influenza information from Europe is available from [WHO/Europe and the European Centre for Disease Prevention and Control](http://www.flunewseurope.org/) [ⓘ](http://www.flunewseurope.org/) (<http://www.flunewseurope.org/>).

Public Health Agency of Canada:

The most up-to-date influenza information from Canada is available in [Canada's weekly FluWatch report](http://www.phac-aspc.gc.ca/fluwatch/) [ⓘ](http://www.phac-aspc.gc.ca/fluwatch/) (<http://www.phac-aspc.gc.ca/fluwatch/>).

Public Health England:

The most up-to-date influenza information from the United Kingdom is available from [Public Health England](http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/SeasonalInfluenza/) [ⓘ](http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/SeasonalInfluenza/) (<http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/SeasonalInfluenza/>).

Any links provided to non-Federal organizations are provided solely as a service to our users. These links do not constitute an endorsement of these organizations or their programs by CDC or the Federal Government, and none should be inferred. CDC is not responsible for the content of the individual organization web pages found at these links.

A description of the CDC influenza surveillance system, including methodology and detailed descriptions of each data component is available on the [surveillance methods](http://www.cdc.gov/flu/weekly/overview.htm) (<http://www.cdc.gov/flu/weekly/overview.htm>) page.

Last Reviewed: June 28, 2024, 11:00 AM

Source: Centers for Disease Control and Prevention (<https://www.cdc.gov/>), National Center for Immunization and Respiratory Diseases (NCIRD) (<https://www.cdc.gov/ncird/index.html>)

Was this page helpful?

Yes

Partly

No