Age-adjusted rate (per 100,000 standard

Jurisdiction	Deaths	per 100,000 standar population)	Comparison to the U.S. rate
United States (overall)	107,941	32.6	-
Alabama	1,492	31.5	Not significantly different than the U.S. rate
Alaska	254	34.3	Not significantly different than the U.S. rate
Arizona	2,664	37.2	Rate significantly higher than U.S. rate
Arkansas	617	21.7	Rate significantly lower than the U.S. rate
California	10,952	26.9	Rate significantly lower than the U.S. rate
Colorado	1,811	29.8	Rate significantly lower than the U.S. rate
Connecticut	1,482	40.3	Rate significantly higher than U.S. rate
Delaware	549	55.3	Rate significantly higher than U.S. rate
District of Columbia	451	64.3	Rate significantly higher than U.S. rate
Florida	7,551	35.2	Rate significantly higher than U.S. rate
Georgia	2,687	24.9	Rate significantly lower than the U.S. rate
Hawaii	284	18.6	Rate significantly lower than the U.S. rate
Idaho	385	20.7	Rate significantly lower than the U.S. rate
Illinois	3,849	30.0	Rate significantly lower than the U.S. rate
Indiana	2,682	41.0	Rate significantly higher than U.S. rate
Iowa	469	15.3	Rate significantly lower than the U.S. rate
Kansas	754	26.5	Rate significantly lower than the U.S. rate
Kentucky	2,271	53.2	Rate significantly higher than U.S. rate
Louisiana	2,376	54.5	Rate significantly higher than U.S. rate
Maine	707	54.3	Rate significantly higher than U.S. rate
Maryland	2,573	40.3	Rate significantly higher than U.S. rate
Massachusetts	2,642	37.4	Rate significantly higher than U.S. rate
Michigan	2,997	30.7	Rate significantly lower than the U.S. rate
Minnesota	1,384	24.8	Rate significantly lower than the U.S. rate
Mississippi	758	27.6	Rate significantly lower than the U.S. rate
Missouri	2,192	36.9	Rate significantly higher than U.S. rate
Montana	208	19.4	Rate significantly lower than the U.S. rate
Nebraska	225	11.8	Rate significantly lower than the U.S. rate
Nevada	1,003	30.3	Rate significantly lower than the U.S. rate
New Hampshire	486	36.0	Rate significantly higher than U.S. rate
New Jersey	2,985	31.6	Not significantly different than the U.S. rate
New Mexico	1,024	50.3	Rate significantly higher than U.S. rate
New York	6,358	31.4	Rate significantly lower than the U.S. rate
North Carolina	4,310	41.8	Rate significantly higher than U.S. rate
North Dakota	148	19.8	Rate significantly lower than the U.S. rate
Ohio	5,144	45.6	Rate significantly higher than U.S. rate
Oklahoma	1,196	30.7	Rate significantly lower than the U.S. rate
Oregon	1,363	31.1	Not significantly different than the U.S. rate
Pennsylvania	5,169	40.9	Rate significantly higher than U.S. rate

Rhode Island	424	38.1	Rate significantly higher than U.S. rate
South Carolina	2,279	44.7	Rate significantly higher than U.S. rate
South Dakota	95	11.3	Rate significantly lower than the U.S. rate
Tennessee	3,825	56.0	Rate significantly higher than U.S. rate
Texas	5,489	18.2	Rate significantly lower than the U.S. rate
Utah	627	19.8	Rate significantly lower than the U.S. rate
Vermont	276	45.9	Rate significantly higher than U.S. rate
Virginia	2,496	28.8	Rate significantly lower than the U.S. rate
Washington	2,725	33.7	Not significantly different than the U.S. rate
West Virginia	1,335	80.9	Rate significantly higher than U.S. rate
Wisconsin	1,792	31.8	Not significantly different than the U.S. rate
Wyoming	126	21.9	Rate significantly lower than the U.S. rate

NOTES: Drug overdose deaths were identified using International Classification of Diseases, Tenth Revision underlying cause-of-death codes X40–X44, X60–X64, X85, and Y10–Y14. Age-adjusted drug overdose death rates were calculated using the direct method and the 2000 U.S. standard population.

SOURCE: National Center for Health Statistics, National Vital Statistics System, Mortality Data.