



HHS Public Access

Author manuscript

Aerosol Air Qual Res. Author manuscript; available in PMC 2024 August 02.

Published in final edited form as:

Aerosol Air Qual Res. 2022 November ; 22(11): . doi:10.4209/aaqr.220047.

Understanding Air Quality Changes after Implementation of Mitigation Measures during a Pandemic: A Scoping Review of Literature in the United States

Sara McElroy^{1,2,*}, Ambarish Vaidyanathan²

¹Oak Ridge Institute for Science and Education, Oak Ridge, TN USA and Climate and Health Program, National Center for Environmental Health, Centers for Disease Control and Prevention, Atlanta, GA 30341, USA

²Climate and Health Program, DEHSP, NCEH, CDC, National Center for Environmental Health, Centers for Disease Control and Prevention, Atlanta, GA 30341, USA

Abstract

Traffic-related emissions continue to be a significant source of air pollution in the United States (US) and around the globe. Evidence has shown that previous policies implemented to restrict-traffic flows have affected air pollution levels. Thus, mitigation strategies associated with the COVID-19 pandemic that modified population-level mobility patterns provide a unique opportunity to study air pollution change across the US. For instance, to slow the spread of the pandemic, state and local governments started implementing various mitigation actions, including stay-at-home directives, social distancing measures, school closures, and travel restrictions. This scoping review aimed to summarize the existing evidence about how air quality changed through mitigation practices throughout the pandemic in the US. We found 66 articles that fit our inclusion criteria. Generally, the consolidated results revealed that nitrogen dioxide (NO₂) and carbon monoxide (CO) decreased across the country. Studies observed mixed directions and magnitudes of change for fine and coarse particulate matter (PM_{2.5}, PM₁₀), ozone (O₃), and sulfur dioxide (SO₂). Few articles tried to explain this notable heterogeneity in air quality changes by associating contextual factors, such as mobility, traffic flow, and demographic factors. However, all studies agreed that the change in air pollution was nonuniform across the US and even varied within a city.

Keywords

COVID-19; Air pollution; Mitigation measures; Mobility; Public health interventions

This is an open access article distributed under the terms of the [Creative Commons Attribution License \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are cited.

*Corresponding Author: Qwi4@cdc.gov.

SUPPLEMENTARY MATERIAL

Supplementary material for this article can be found in the online version at <https://doi.org/10.4209/aaqr.220047>

1 INTRODUCTION

Traffic-related emissions are an important source of ambient air pollution to which the majority of the population is exposed to in the United States (US) and other parts of the world. On-road combustion sources are a major contributor of various air pollutants, including criteria pollutants such as nitrogen dioxide (NO₂), carbon monoxide (CO), fine particulate matter (PM_{2.5}), and ozone (O₃) (Bradley *et al.*, 1999, Russell *et al.*, 2012). In the past, there has been evidence showing that strategies implemented to restrict traffic-related mobility have led to changes in air quality levels (Peel *et al.*, 2010). For example, policies that were implemented to reduce traffic congestion during the Atlanta and Beijing Olympic Games, were found to be associated with reduced air pollutant concentrations (Friedman *et al.*, 2001; Li *et al.*, 2010). These strategies included 24-hour public transportation, alternative work hours and telecommuting for local businesses, banning of vehicles that failed emission standards, restricting private vehicles to drive every other day, and closures of areas to private vehicles (Balagas, 1996; Porter, 1997; The People's Government of Beijing Municipality, 2009). The changes in population-level mobility during the COVID-19 pandemic has provided a similar opportunity to study how mitigation strategies have impacted air quality in the United States.

The World Health Organization (WHO) issued a public health emergency on January 30, 2020, when coronavirus, SARS-CoV-2, was starting to spread across the world (WHO, 2020; Bashir *et al.*, 2020; Wu *et al.*, 2020). The WHO declared the disease a global pandemic on March 11, 2020 (Cucinotta and Vanelli, 2020), soon followed by Centers for Disease Control and Prevention (CDC) emphasizing the evidence of community spread in the United States. Many major cities across the United States took actions to mitigate community spread and began implementing stay-at-home orders, lockdown measures, social distancing measures, travel restrictions, school closures, and promoting telecommuting and remote work options (Cohen and Kupferschmidt, 2020; Pepe *et al.*, 2020; Schlosser *et al.*, 2020).

These mitigation efforts and countermeasures for preventing the spread of the pandemic have brought about behavioral changes in our society. For instance, remote work and telecommuting options combined with stay-at-home directives have brought about shifts in mobility patterns. These personal behavioral modifications and community-level mitigation practices provide an opportunity to assess the relationship between interventions and their collective impact on air quality levels.

However, community-level mitigation policies vary significantly across states or even within counties in the US (Raifman *et al.*, 2020). Researchers have assessed changes in air pollution, which reflect the impact of mitigation efforts on emission changes associated with various air pollution sources, and changes in standard daily activity patterns within cities by analyzing mobility, traffic, and economic information (Campbell *et al.*, 2021; Heintzelman *et al.*, 2021; Winchester *et al.*, 2021). For instance, researchers have investigated the impact of stay-at-home orders on air quality levels under the assumption that changes in transportation patterns and mobility would lead to lower levels of traffic-related air pollutants. Of note, air pollutants such as NO₂, O₃, CO, coarse particulate matter (PM₁₀) and PM_{2.5} are

determinants of economic and human development (Venter *et al.*, 2021). Primary sources of these air pollutants include transportation and other non-road combustion sources (Russell *et al.*, 2012; Bradley *et al.*, 1999; Manousakas *et al.*, 2017).

We conducted this scoping review to summarize published scientific articles examining how air quality changed due to community-level pandemic mitigation measures within the US. This scoping review aims to summarize findings from published research exploring change in air pollution levels occurring in the period with COVID-19 mitigation practices (e.g., shelter-in-place directives). Specifically, we report on exposure data used, methodologies, the results of these assessments, contextual factors, mediators, and other studies outcomes. Our review summarizes the available evidence to provide an overview of the topic for researchers interested in linking measurable relationships between policies that reduce emissions and pollutant concentration.

Specifically, the review question is: how did ambient air pollution change from previous years to the period where COVID-19 restrictions were enacted in the US? Furthermore, how did this observed change in air pollution vary by geography and time? In addition, have studies explored contextual factors that can explain the relationship between air quality changes and COVID-19 mitigation measures?

2 METHODS

2.1 Study Design

We developed our scoping review using the methodology framework proposed by the Joanna Briggs Institute, built on the previous methodology proposed by Arksey and O'Malley (Arksey and O'Malley, 2005; Peters *et al.*, 2015). This review was conducted utilizing the five steps recommended by Arksey and O'Malley (2005): 1) identifying a valid research question and defining search strategies, 2) identifying relevant studies, 3) study selection, 4) extracting and charting data, and 5) collating, summarizing, discussing, and reporting results.

2.2 Search Strategy

The leading search strategy aims to identify published scientific articles, or articles in preprint, in English. An initial limited search of MEDLINE (PubMed) and Google Scholar was executed to detect articles pertaining to this topic. These articles provided insights for text words that were included in titles, abstracts, and keywords. The information found from the preliminary search informed the development of a search strategy, including identified keywords and index terms that were tailored for each database. Since COVID-19 is a very recent development, we decided to constrict our scoping review to articles that have been published or are in preprint after the implementation of COVID-19 mitigation strategies (March 1, 2020, through October 1, 2021). The reference list of included studies was also assessed to find additional studies. The literature for this review was identified by searching the following online databases: PubMed, Web of Science, Embase, Global Health, Environmental Science Collection, Scopus, and Google Scholar. Our inclusion criteria included scientific articles published in English within the designated date range

that discussed changes in air quality within the United States. We considered scientific commentaries, editorials, descriptive studies, comparison studies, exposure studies, and epidemiological study designs, including before and after studies, ecological studies, cross-sectional studies, cohort studies, case-control studies, spatiotemporal studies, and cluster analyses. Conference abstracts are not included in this study. The scope of this review does not include nonscientific publications, news articles, unpublished and gray literature, and systematic literature reviews. The complete list of search terms is located in Appendix A.

2.3 Identification and Selection of Relevant Studies

The seven online databases were searched using the predetermined keywords. An example of these key words search is “air quality” OR “air pollution” OR “ambient exposure” AND “COVID-19*” OR “SARS Coronavirus 2 Infection” OR “SARS*” AND “US” OR “United States” If a question arose about the inclusion or exclusion of literature, an independent second researcher was brought in to make the final decision. Duplicate articles were eliminated. Fig. 1 depicts the process of our inclusion and exclusion of articles by a flow diagram per the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) methodology (Moher *et al.*, 2009).

2.4 Data Extraction

Data extracted from articles included in the scoping review were entered into an excel spreadsheet by a reviewer. Specific details about the populations, concept, context, and study methodologies of significance to the scoping review research question and aims were abstracted. More specifically, the author, journal’s name, study population, study location, study design, statistical analyses, source of exposure data, outcome, study period, covariates, main findings, and reviewer comments were included in the data abstracted.

2.5 Methodology Quality and Appraisal

Because our research aims were to summarize available evidence and identify gaps in the literature, we did not appraise the methodology quality or bias of the included articles, consistent with guidance on scoping review conduct (Peters *et al.*, 2015).

2.6 Presentation of Results and Summarizing the Findings

Following our proposed research objectives, articles fell into three main categories. The first category entailed publications that only examined changes in air pollution between pre-and-during-COVID-19. The pre-COVID-19 period in the U.S. is defined as any time before March 1st, 2020, and the during-COVID-19 period is defined as any time after March 31st, 2020, to December 31st, 2021. The second group of articles examined changes across space and time by examining changes in air pollution across several geographic locations. The third group included contextual factors to explain the differences in air quality between pre-and current-COVID-19 periods. All results and statements regarding changes in air quality throughout the pandemic restrictions were abstracted from published information.

3 RESULTS

We identified 538 published articles; 309 articles were excluded as duplicates. After title and abstract screening were conducted for the remaining 229 articles, 154 were excluded because they did not fit our inclusion criteria. If the title indicated the study was performed outside the United States, involved examining COVID-19 mortality/case rates, studied indoor air pollution, or examined the biology of the virus, the article was excluded. A full-text review was implemented for the remaining 75 articles, 12 articles were excluded due to subject matter, or the article was a conference abstract. After examining citations from these articles, an additional three studies were found, resulting in a total of 66 scientific articles included in this scoping review (Fig. 1). The data abstraction table for the included articles in this scoping review can be found in Appendix B.

3.1 Literature Characteristics

Among the 66 included articles in this study, publication dates ranged from June 2020 to preprint articles that will be published in November of 2021. A little less than half the studies were published in 2020 (46%) compared to 2021 (54%). Included articles were conducted in a single city ($n = 10$, 15%), the continental US or multiple cities within the US ($n = 14$, 17%), one state in the US ($n = 7$, 9%), a US city and other international cities ($n = 5$, 8%), several countries including the US ($n = 9$, 15%) and globally ($n = 21$, 34%). The included articles were published in 29 different peer-review journals, with the most articles published in *Science of the Total Environment* ($n = 12$, 18%). Study periods ranged from 6 weeks to the entire year of 2020 (Table 1).

3.2 Air pollution Data

The included articles studied air pollutants or proxy measures of air quality including NO₂ and/or other nitrogen oxides (NO_x), CO, PM_{2.5}, PM₁₀, O₃, aerosol optical depth (AOD, a proxy for particulates), black carbon (BC), and SO₂. Forty-three studies included more than one air pollutants (65%). Twenty-three studies examined one air pollutant, which was either NO₂, PM_{2.5}, or CO₂. The distribution of the number of studies that included each air quality measure can be seen in Fig. 2. Over half of the studies measured changes in air pollution concentrations, 19% measured changes in both concentrations and emissions, and 25% measured changes in vertical column density. There were 34 studies that retrieved air pollutant measurements from ground monitors (51%), 13 studies combined information from ground monitors and satellite data (20%), 14 studies relied on satellite data alone (21%), and five studies utilized mobile air pollution monitors (8%) to measure air pollutants. Articles which included ground monitor measurements retrieved those data from the US Environmental Protection Agency (EPA), AirNow (a partnership with the EPA), National Oceanic Atmospheric Administration (NOAA), National Park Service, NASA, CDC, and tribal, state, and local air quality agencies (<https://www.airnow.gov/>), Aerosol Robotic Network (AERONET), OpenAQ API, local county air quality measures, and the World Air Quality Index (WAQI) project. Remotely sensed data were obtained from multiple sources, which include the Tropospheric Monitoring Instrument (TROPOMI) onboard Sentinel-5 Precursor, the Ozone Monitoring Instrument (OMI) on Aura satellite, Moderate Resolution Imaging Spectroradiometer (MODIS) instrument on NASA's Terra and Aqua satellites, the

NOAA Suomi-NPP, and the Measure of Pollution in the Troposphere (MOPITT) on NASA's Terra satellite (Fig. 2).

3.3 Study Design

Many of the included articles' study design capitalized on the opportunity that arose from COVID-19 restrictions, which created a natural experiment, allowing for comparison of before and after periods (Bar *et al.*, 2021; Campbell *et al.*, 2021; Mendoza *et al.*, 2021). Percent change methodology was utilized the most ($n = 24$) to quantify the change in air quality between a historical period and the COVID-19 period (See Fig. 4). Additional methodologies include unsupervised machine learning algorithms (Venter *et al.*, 2020; Keller *et al.*, 2021; Zhang *et al.*, 2021), anomalies calculation (Zhang *et al.*, 2020), time-lagged linear regression with ANCOVA and F-tests (Zangari *et al.*, 2020), difference-in-difference quasi-experimental method (Brodeur *et al.*, 2021), probability density functions (Mishra *et al.*, 2021), two-sample t-tests (Berman and Ebisu, 2020; Jia *et al.*, 2020; Chadwick *et al.*, 2021; Elshorbany *et al.*, 2021), non-parametric Wilcoxon rank-sum test (Hudda *et al.*, 2020), regression discontinuity design (Hudda *et al.*, 2020), robust differences with median and interquartile range (IQR) (Bekbulat *et al.*, 2021) and general linear models (Connerton *et al.*, 2020). Four articles were descriptive and did not perform any statistical analyses (Balasubramaniam *et al.*, 2020; Hoang *et al.*, 2021; Irfan *et al.*, 2021; Mousazadeh *et al.*, 2021). Ten studies were time-series analyses utilizing multivariate autoregressive models (Xiang *et al.*, 2020), seasonal autoregressive integrated moving average with exogenous factors (SARIMAX) (Wong *et al.*, 2021), Stochastic Time-Inverted Lagrangian Transport model (Turner *et al.*, 2020), general additive model (GAM) (Liu *et al.*, 2021a), moving averages (Liu *et al.*, 2021b, 2021c), autoregressive moving averages models (ARIMA) (Winchester *et al.*, 2021), principal component analysis (Savtchenko and Khayat, 2021), and mixed effect models (Ghosal and Saha, 2021). Lastly, one study performed a spatial cluster analysis using Getis-Ord (G_i^*) spatial statistic (Straka *et al.*, 2021).

3.4 Principal Findings

The change in ambient air quality from pre-COVID mitigation strategies was found to have increased, decreased, or stay the same depending on air pollutant and geographical location. There was a consensus that NO_2 and CO decreased within the COVID-19 lockdown period, but the decreased level varied across geographical regions. Reductions in NO_2 varied widely across studies from 18% to 63%. Also, many studies agreed that O_3 increased in urban areas and decreased in rural areas throughout the policies implemented to stop the spread of COVID-19. Increases in O_3 percent change ranged from 17% to 86%, and the decrease in O_3 ranged from 1% to 10%. There were mixed results when examining $\text{PM}_{2.5}$. Some studies found decreases in $\text{PM}_{2.5}$, increases in $\text{PM}_{2.5}$, and other studies found no variation from historical levels. Finally, two studies found no change in any air pollution criterion levels in 2020 than historical data. The effect sizes and results from each study can be seen in Appendix: Table S1.

City-specific results fit the general pattern observed from the overall findings. NO_2 and CO level reductions were reported in all cities, with the most significant NO_2 and CO emissions decrease in Los Angeles and New York City, respectively. $\text{PM}_{2.5}$, O_3 , and SO_2

levels varied across cities and within cities, with both observed increases and decreases of the air pollutant. The reduction of $PM_{2.5}$ levels was the largest in Los Angeles, and the largest increase was recorded in New York City. O_3 increased in New York City but decreased in Memphis and Los Angeles. Directions and magnitudes of air pollutant changes from city-specific studies can be seen in Fig. 3. Methodologies used in each of these studies can be seen in Fig. 4.

The sources of emission of NO_2 consist of 90% from on-road and non-road combustion and biomass burning, and the remaining 10% primarily comes from chemical productions at power plants (Richter *et al.*, 2004; Otmani *et al.*, 2020). Studies that found a reduction in NO_2 during the COVID-19 restrictions tied this reduction to both sources. The stay-at-home orders significantly reduced motor vehicle traffic, which resulted in a significant reduction of vehicle exhaust emissions (Archer *et al.*, 2020; Chen *et al.*, 2020; Connerton *et al.*, 2020; Wang and Li, 2021). Furthermore, some factories were closed during the early stages of the pandemic, which curbed the production of NO_2 (Wang and Li, 2021). CO reductions were consistent across studies, and these reductions were also linked to vehicle emissions, as vehicle traffic is the primary source of both NO_2 and CO (Tanzer-Gruener *et al.*, 2020). Moreover, NO_2 and CO are primary pollutants; thus, changes in activity can be directly connected with emission levels.

Several hypotheses were proposed to explain why some studies did not observe a decrease in $PM_{2.5}$ during the lockdown period. One such proposed reason was that the stay-at-home orders mainly affected individuals' driving patterns. However, commercial vehicles (fueled primarily by diesel) and electricity demand (often met by burning of fossil fuels) remained unchanged. As a result, a reduction in $PM_{2.5}$ might not be prominent (Archer *et al.*, 2020). Secondly, $PM_{2.5}$ is essentially a pollutant created from complex and nonlinear atmospheric chemistry. A key component in this relationship is the volatile organic compound to NO_x ratio (VOC: NO_x). When this ratio increases, it increases the production of secondary organic aerosols, a primary component in $PM_{2.5}$; thus, leading to an increase in $PM_{2.5}$ (Connerton *et al.*, 2020; Bekbulat *et al.*, 2021). Lastly, PM can also be attributed to residential fuel combustion, waste management, and fugitive dust, which may not have been affected by COVID-19 restrictions (Chen *et al.*, 2020).

Increases in O_3 concentrations have been attributed to the nonlinear photochemical reaction, which involves NO_x , VOCs, and sunlight (Jin *et al.*, 2017). Effects of the precursor concentration production rate can be categorized as either NO_x limited, or VOC limited. NO_x -limited is when the VOC: NO_x ratio is large, and the VOC-limited environment happens when the VOC: NO_x ratio is smaller. In areas with a VOC-limited regime, a decrease in NO_x concentrations will cause an increase in O_3 production (Chen *et al.*, 2020; Fu *et al.*, 2020). Most of the studies that observed increases in O_3 cited this photochemical reaction as an explanation.

3.5 Contextual Factors

Twenty articles tried to explain the change of air pollution levels through different contextual factors. Eight studies incorporated mobility data into the analyses. Sources of these mobility data were from Google, Apple, and a dataset developed by Descartes Labs that provides an

aggregated mobility measure based on anonymized mobile device locations. Nine articles examined how traffic flow was related to changes in air quality. Traffic flow information was obtained via TRACFLOW (Xiang *et al.*, 2020), TomTom traffic congestion data (Liu *et al.*, 2020; Winchester *et al.*, 2021), Caltrans Performance Measurement System (Liu *et al.*, 2021a; Naeger and Murphy, 2020; Parker *et al.*, 2020), Transit app (Pan *et al.*, 2020), Streetlight VMT (Jia *et al.*, 2020), and the Federal Highway Administration (Elshorbany *et al.*, 2021). Parameters used in these analyses were total vehicle volume (TOV), road occupancy (%), public transit demand, vehicle miles traveled (VMT), and daily traffic counts. Two studies performed a health assessment of how many prevented premature deaths resulted from reducing air pollution during the COVID-19 restrictions. Finally, two studies examined the disparities of NO₂ pollution for disproportionately impacted populations across the US (Winchester *et al.*, 2021). Their findings revealed the most significant reductions occurred in marginalized areas, but the effect of lockdowns on racial, ethnic, and socioeconomic NO₂ disparities was mixed and, for many cities, nonsignificant (Kerr *et al.*, 2021).

4 DISCUSSION

Generally, we saw a decrease in NO₂ and CO and mixed directions of other air pollutants in response to COVID-19 mitigation strategies, and the effects of pandemic mitigation practices on air pollutants were heterogeneous across the US. This heterogeneity was observed across both large and small spatial scales. Changes in air pollutants were found to differ by and within regions, counties, and cities. This observed heterogeneity highlights the complex interactions between air pollutants, meteorological conditions, pandemic mitigation efforts, geographical location, and other factors that could be driving this array of effects. Studies that examined only metropolitan areas found nonuniform effects of COVID-19 mitigation policies on the same air pollutant. Only a few studies tried to elucidate these nonuniform effects by examining individual mobility and traffic flow patterns.

We identified crucial gaps in the literature through our scoping review, which examined how air quality changed through the span of this scoping review. One such gap that was elucidated related to the dates of publication and the length of study periods. Nearly all studies included in this scoping review encompassed study periods that only covered a portion of 2020. Most of the published articles in 2020 examined changes in air quality only until May of 2020. Furthermore, many studies published in 2021 did not assess the change in air pollution for 2020 in its entirety. These truncations of study periods resulted in missed information about the renewed COVID-19 mitigation strategies implemented because of the significant surge of COVID-19 cases at the end of 2020 in the US, and the emergence and additional practices employed to reduce the spread of the subsequent novel coronavirus variants. Truncated study periods do ignore potential changes to air quality during other times when mitigation strategies were in effect to address the surges of COVID-19 cases, however, the impact of these strategies on air quality changes can be different. For instance, understanding changes in mobility patterns during the initial wave of cases and comparing that to subsequent periods when mitigation strategies were in effect can help quantify impact on relevant air pollution sources.

While mitigation measures have influenced transportation-related emissions, there are other emission sources that could have contributed to an increase in air pollutant concentrations. Most notably, many of these articles only explored air quality changes during the initial stages of the pandemic and did not include time periods when record-breaking wildfires occurred, during the latter part of 2020. We acknowledge that there was an urgency to disseminate findings from these studies, which could have led to the short study periods. However, these short study periods would not allow for ascertainment of contributions from individual sources in areas where air pollution is from multiple sources. Wildfires are a significant source of air pollutant emissions, including PM_{2.5} (Ford *et al.*, 2018; Gupta *et al.*, 2018; Aguilera *et al.*, 2021). These increased emissions of PM_{2.5} could have influenced the findings of studies that examined air pollutant changes throughout 2020, especially in the Western US. The possible effect of excess PM_{2.5} from wildfires was supported by the findings of two articles included in this scoping review that examined air pollution changes until September and December of 2020, which found PM_{2.5} and AOD levels were higher than expected (Bekbulat *et al.*, 2021; Elshorbany *et al.*, 2021).

The majority of articles included in this scoping review relied upon ground monitors or remote sensing data. We believe the researchers had a good argument to utilize remote sensing data because it gives the opportunity to retrieve near real-time aerosol data (if data from ground monitors are not available), even though calibration and cloud cover can be an issue (Berman and Ebisu, 2020). Although we did not quantitatively assess the quality of the articles included in the scoping review, we found that over half the articles utilized basic methodologies that included descriptive studies, correlation studies, and comparison studies calculating percent change. Ideally, a measurement-based study could address some of the limitations identified in these papers. For example, performing a time-series analysis using autoregressive moving averages model adjusted for meteorological and mobility confounders and accounting for seasonality of the pollutants and meteorological variables could be a valid approach. Additionally, articles that examined change in air quality as a function of mobility should have chosen an appropriate data source that addresses the research question at hand, as these data vary by individual and aggregated scale and across various spatial and temporal scales (Hu *et al.*, 2021).

A concerning body of evidence suggests that climate hazards such as extreme heat, wildfires, drought, floods, and hurricanes, which are increasing in frequency and intensity in many regions under climate change, intersect with the COVID-19 pandemic and public health response (Phillips *et al.*, 2020). Thus, it might be equally important to understand how exposure to these additional climate hazards was affected by pandemic mitigation strategies in the US, as there is growing evidence to suggest that cascading health effects resulting from exposure to compound hazards (e.g., air pollution and extreme heat) is a significant public health concern. Furthermore, evidence from this scoping review, which summarized the relationships between changes in air pollution and mitigation practices, can shed light on how climate adaptation and mitigation efforts aimed at reducing greenhouse gas emissions might influence air quality levels at local and regional levels scales.

There are several limitations to this scoping review. Articles published after the retrieval date were not able to be included in our study. Also, we did not rank the suitability of

the methodology or any inherent biases due to the study design implemented for these articles because our main goal was to summarize the change in air quality linked to COVID-19 mitigation strategies. Future research could aim to weigh the methodologies used across these studies differently. Furthermore, due to the numerous studies examining air quality changes within the same geographical area, there were instances where the magnitude of change for air pollutants was reported using different concentration units that were not comparable (e.g., vertical column density [moles m⁻²] versus routinely reported concentration units [ppb]). In such instances, we excluded studies using those non-traditional reporting units from Fig. 3.

5 CONCLUSION

Overall, we identified 66 scientific articles that assessed the change in air quality related to COVID-19 mitigation efforts. Evidence from these studies showed nonuniform changes in air pollution at the regional, state, county, and local scales across the US. CO and NO₂ generally decreased across all geographic locations, O₃ was found to increase in urban areas and decrease in rural areas, and PM_{2.5} concentrations changes were the most variable across the US. A portion of the studies included in this review attempted to explain the observed heterogeneity in air quality change by examining how various contextual factors, such as mobility, traffic patterns, and socioeconomic status, could be associated and found that traffic-related emissions and mobility decreased the most in early 2020. This scoping review has summarized the available evidence and can be utilized to elucidate further the relationship between pandemic mitigation practices and changes in air quality. Lastly, from a climate change perspective, understanding the relationship between air pollution and contextual factors, such as measures of baseline health equity, behavioral modification due to compounding hazards, changes in mobility patterns as measured before and after a public health intervention, can help inform the effectiveness of policies aimed at reducing greenhouse gas emissions.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENT

Dr. Sara McElroy's work was supported in part by an appointment to the research participation program at the Centers for Disease Control and Prevention, administered by Oak Ridge Institute of Science and Education through an Interagency agreement between the U.S. Department of Energy and the Centers for Disease Control and Prevention

REFERENCES

- Acharya P, Barik G, Gayen BK, Bar S, Maiti A, Sarkar A, Ghosh S, De SK, Sreekesh S (2021). Revisiting the levels of Aerosol Optical Depth in south-southeast Asia, Europe and USA amid the COVID-19 pandemic using satellite observations. *Environ. Res* 193, 110514. 10.1016/j.envres.2020.110514
- Aguilera R, Corringham T, Gershunov A, Benmarhnia T (2021). Wildfire smoke impacts respiratory health more than fine particles from other sources: Observational evidence from Southern California. *Nat. Commun* 12, 1493. 10.1038/s41467-021-21708-0 [PubMed: 33674571]

- Amouei Torkmahalleh M, Akhmetvaliyeva Z, Omran AD, Faezeh Darvish Omran F, Kazemitabar M, Naseri Mahtab, Naseri Motahareh, Sharifi H, Malekipirbazari M, Kwasi Adotey E, Gorjinezhad S, Eghtesadi N, Sabanov S, Alastuey A, de Fátima Andrade M, Buonanno G, Carbone S, Cárdenas-Fuentes DE, Cassee FR, Dai Q, et al. (2021). Global air quality and COVID-19 pandemic: Do we breathe cleaner air? *Aerosol Air Qual. Res* 21, 200567. 10.4209/aaqr.200567
- Archer CL, Cervone G, Golbazi M, Al Fabel N, Hultquist C (2020). Changes in air quality and human mobility in the USA during the COVID-19 pandemic. *Bull. Atmos. Sci. Technol* 1, 491–514. 10.1007/s42865-020-00019-0 [PubMed: 38624442]
- Arksey H, O'Malley L (2005). Scoping studies: Towards a methodological framework. *Int. J. Soc. Res. Methodol* 8, 19–32. 10.1080/1364557032000119616
- Balagas R (1996). Improved Air Quality in Atlanta During the 1996 Olympic Summer Games. Atlanta: Georgia Environmental Protection Division. Georgia Department of Natural Resources.
- Balasubramaniam D, Kanmanipappa C, Shankarlal B, Saravanan M (2020). Assessing the impact of lockdown in US, Italy and France—What are the changes in air quality? *Energy Sources Part A* 1–11. 10.1080/15567036.2020.1837300
- Bar S, Parida BR, Mandal SP, Pandey AC, Kumar N, Mishra B (2021). Impacts of partial to complete COVID-19 lockdown on NO₂ and PM_{2.5} levels in major urban cities of Europe and USA. *Cities* 117, 103308. 10.1016/j.cities.2021.103308
- Bashir MF, Ma BJ, Bilal, Komal B, Bashir MA, Farooq TH, Iqbal N, Bashir M (2020). Correlation between environmental pollution indicators and COVID-19 pandemic: A brief study in Californian context. *Environ. Res* 187, 109652. 10.1016/j.envres.2020.109652
- Bauwens M, Compennolle S, Stavrakou T, Müller JF, Gent J van, Eskes H, Levelt PF, van der A. R, Veefkind JP, Vlietinck J, Yu H, Zehner C (2020). Impact of coronavirus outbreak on NO₂ pollution assessed using TROPOMI and OMI observations. *Geophys. Res. Lett* 47, e2020GL087978. 10.1029/2020GL087978
- Bekbulat B, Apte JS, Millet DB, Robinson AL, Wells KC, Presto AA, Marshall JD (2021). Changes in criteria air pollution levels in the US before, during, and after Covid-19 stay-at-home orders: Evidence from regulatory monitors. *Sci. Total Environ* 769, 144693. 10.1016/j.scitotenv.2020.144693
- Berman JD, Ebisu K (2020). Changes in US air pollution during the COVID-19 pandemic. *Sci. Total Environ* 739, 139864. 10.1016/j.scitotenv.2020.139864
- Bradley KS, Stedman DH, Bishop GA (1999). A global inventory of carbon monoxide emissions from motor vehicles. *Chemosphere Global Change Sci.* 1, 65–72. 10.1016/S1465-9972(99)00017-3
- Bray CD, Nahas A, Battye WH, Aneja VP (2021). Impact of lockdown during the COVID-19 outbreak on multi-scale air quality. *Atmos. Environ* 254, 118386. 10.1016/j.atmosenv.2021.118386
- Brodeur A, Cook N, Wright T (2021). On the effects of COVID-19 safer-at-home policies on social distancing, car crashes and pollution. *J. Environ. Econ. Manage* 106, 102427. 10.1016/j.jeem.2021.102427
- Campbell PC, Tong D, Tang Y, Baker B, Lee P, Saylor R, Stein A, Ma S, Lamsal L, Qu Z (2021). Impacts of the COVID-19 economic slowdown on ozone pollution in the U.S. *Atmos. Environ* 264, 118713. 10.1016/j.atmosenv.2021.118713
- Chadwick E, Le K, Pei Z, Sayahi T, Rapp C, Butterfield AE, Kelly KE (2021). Technical note: Understanding the effect of COVID-19 on particle pollution using a low-cost sensor network. *J. Aerosol Sci* 155, 105766. 10.1016/j.jaerosci.2021.105766
- Chauhan A, Singh RP (2020). Decline in PM_{2.5} concentrations over major cities around the world associated with COVID-19. *Environ. Res* 187, 109634. 10.1016/j.envres.2020.109634
- Chen LWA, Chien LC, Li Y, Lin G (2020). Nonuniform impacts of COVID-19 lockdown on air quality over the United States. *Sci. Total Environ* 745, 141105. 10.1016/j.scitotenv.2020.141105
- Chossière GP, Xu H, Dixit Y, Isaacs S, Eastham SD, Allroggen F, Speth RL, Barrett SRH (2021). Air pollution impacts of COVID-19–related containment measures. *Sci. Adv* 7, eabe1178. 10.1126/sciadv.abe1178
- Cohen J, Kupferschmidt K (2020). Strategies shift as coronavirus pandemic looms. *Science* 367, 962–963. 10.1126/science.367.6481.962 [PubMed: 32108093]

- Connerton P, Vicente de Assunção J, Maura de Miranda R, Dorothée Slovic A, José Pérez-Martínez P, Ribeiro H (2020). Air quality during COVID-19 in four megacities: Lessons and challenges for public health. *IJERPH* 17, 5067. 10.3390/ijerph17145067 [PubMed: 32674410]
- Cucinotta D, Vanelli M (2020). WHO declares COVID-19 a pandemic. *Acta Biomed.* 91, 157–160. 10.23750/abm.v91i1.9397 [PubMed: 32191675]
- Dang HAH, Trinh TA (2021). Does the COVID-19 lockdown improve global air quality? New cross-national evidence on its unintended consequences. *J. Environ. Econ. Manage* 105, 102401. 10.1016/j.jeem.2020.102401
- El-Sayed MM, Elshorbany YF, Koehler K (2021). On the impact of the COVID-19 pandemic on air quality in Florida. *Environ. Pollut* 285, 117451. 10.1016/j.envpol.2021.117451
- Elshorbany YF, Kapper HC, Ziemke JR, Parr SA (2021). The status of air quality in the United States during the COVID-19 pandemic: A remote sensing perspective. *Remote Sens.* 13, 369. 10.3390/rs13030369
- Ford B, Val Martin M, Zelasky SE, Fischer EV, Anenberg SC, Heald CL, Pierce JR (2018). Future fire impacts on smoke concentrations, visibility, and health in the contiguous United States. *GeoHealth* 2, 229–247. 10.1029/2018GH000144 [PubMed: 32159016]
- Friedman MS, Powell KE, Hutwagner L, Graham LM, Teague WG (2001). Impact of changes in transportation and commuting behaviors during the 1996 Summer Olympic Games in Atlanta on air quality and childhood asthma. *JAMA* 285, 897–905. 10.1001/jama.285.7.897 [PubMed: 11180733]
- Fu F, Purvis-Roberts KL, Williams B (2020). Impact of the COVID-19 pandemic lockdown on air pollution in 20 major cities around the world. *Atmosphere* 11, 1189. 10.3390/atmos11111189
- Ghosal R, Saha E (2021). Impact of the COVID-19 induced lockdown measures on PM_{2.5} concentration in USA. *Atmos. Environ* 254, 118388. 10.1016/j.atmosenv.2021.118388
- Goldberg DL, Anenberg SC, Griffin D, McLinden CA, Lu Z, Streets DG (2020). Disentangling the impact of the COVID-19 lockdowns on urban NO₂ from natural variability. *Geophys. Res. Lett* 47, e2020GL089269. 10.1029/2020GL089269
- Gupta P, Doraiswamy P, Levy R, Pikelnaya O, Maibach J, Feenstra B, Polidori A, Kiros F, Mills KC (2018). Impact of California fires on local and regional air quality: The role of a low-cost sensor network and satellite observations. *GeoHealth* 2, 172–181. 10.1029/2018GH000136 [PubMed: 31157310]
- Habibi H, Awal R, Fares A, Ghahremannejad M (2020). COVID-19 and the improvement of the global air quality: The bright side of a pandemic. *Atmosphere* 11, 1279. 10.3390/atmos11121279
- Hammer MS, van Donkelaar A, Martin RV, McDuffie EE, Lyapustin A, Sayer AM, Hsu NC, Levy RC, Garay MJ, Kalashnikova OV, Kahn RA (2021). Effects of COVID-19 lockdowns on fine particulate matter concentrations. *Sci. Adv* 7, eabg7670. 10.1126/sciadv.abg7670
- Heintzelman A, Filippelli G, Lulla V (2021). Substantial decreases in U.S. cities' ground-based NO₂ concentrations during COVID-19 from reduced transportation. *Sustainability* 13, 9030. 10.3390/su13169030
- Hoang AT, Huynh TT, Nguyen XP, Nguyen TKT, Le TH (2021). An analysis and review on the global NO₂ emission during lockdowns in COVID-19 period. *Energy Sources Part A* 1–21. 10.1080/15567036.2021.1902431
- Hu T, Wang S, She B, Zhang M, Huang X, Cui Y, Khuri J, Hu Y, Fu X, Wang X, Wang P, Zhu X, Bao S, Guan W, Li Z (2021). Human mobility data in the COVID-19 pandemic: Characteristics, applications, and challenges. *Int. J. Digital Earth* 14, 1126–1147. 10.1080/17538947.2021.1952324
- Hudda N, Simon MC, Patton AP, Durant JL (2020). Reductions in traffic-related black carbon and ultrafine particle number concentrations in an urban neighborhood during the COVID-19 pandemic. *Sci. Total Environ* 742, 140931. 10.1016/j.scitotenv.2020.140931
- Irfan M, Ahmad M, Fareed Z, Iqbal N, Sharif A, Wu H (2022). On the indirect environmental outcomes of COVID-19: short-term revival with futuristic long-term implications. *Int. J. Environ. Health Res* 32, 1271–1281. 10.1080/09603123.2021.1874888 [PubMed: 33448868]

- Jia C, Fu X, Bartelli D, Smith L (2020). Insignificant impact of the “stay-at-home” order on ambient air quality in the Memphis metropolitan Area, U.S.A. *Atmosphere* 11, 630. 10.3390/atmos11060630
- Jiang Z, Shi H, Zhao B, Gu Y, Zhu Y, Miyazaki K, Lu X, Zhang Y, Bowman KW, Sekiya T, Liou KN (2021). Modeling the impact of COVID-19 on air quality in southern California: Implications for future control policies. *Atmos. Chem. Phys.* 21, 8693–8708. 10.5194/acp-21-8693-2021
- Jin X, Fiore AM, Murray LT, Valin LC, Lamsal LN, Duncan B, Folkert Boersma K, De Smedt I, Abad GG, Chance K, Tonnesen GS (2017). Evaluating a space-based indicator of surface ozone-NO_x-VOC sensitivity over midlatitude source regions and application to decadal trends: Space-based indicator of O₃ sensitivity. *J. Geophys. Res.* 122, 10439–10461. 10.1002/2017JD026720
- Keller CA, Evans MJ, Knowland KE, Hasenkopf CA, Modekurty S, Lucchesi RA, Oda T, Franca BB, Mandarino FC, Díaz Suárez MV, Ryan RG, Fakes LH, Pawson S (2021). Global impact of COVID-19 restrictions on the surface concentrations of nitrogen dioxide and ozone. *Atmos. Chem. Phys.* 21, 3555–3592. 10.5194/acp-21-3555-2021
- Kerr GH, Goldberg DL, Anenberg SC (2021). COVID-19 pandemic reveals persistent disparities in nitrogen dioxide pollution. *PNAS* 118, e2022409118. 10.1073/pnas.2022409118
- Kondragunta S, Wei Z, McDonald BC, Goldberg DL, Tong DQ (2021). COVID-19 induced fingerprints of a new normal urban air quality in the United States. *J. Geophys. Res.* 126, e2021JD034797. 10.1029/2021JD034797
- Li Y, Wang W, Kan H, Xu X, Chen B (2010). Air quality and outpatient visits for asthma in adults during the 2008 Summer Olympic Games in Beijing. *Sci. Total Environ* 408, 1226–1227. 10.1016/j.scitotenv.2009.11.035 [PubMed: 19959207]
- Liu Z, Ciaia P, Deng Z, Lei R, Davis SJ, Feng S, Zheng B, Cui D, Dou X, Zhu B, Guo Rui, Ke P, Sun T, Lu C, He P, Wang Yuan, Yue X, Wang Yilong, Lei Y, Zhou H, et al. (2020). Near-real-time monitoring of global CO₂ emissions reveals the effects of the COVID-19 pandemic. *Nat. Commun* 11, 5172. 10.1038/s41467-020-18922-7 [PubMed: 33057164]
- Liu J, Lipsitt J, Jerrett M, Zhu Y (2021a). Decreases in near-road NO and NO₂ concentrations during the COVID-19 pandemic in California. *Environ. Sci. Technol. Lett* 8, 161–167. 10.1021/acs.estlett.0c00815
- Liu Q, Harris JT, Chiu LS, Sun D, Houser PR, Yu M, Duffy DQ, Little MM, Yang C (2021b). Spatiotemporal impacts of COVID-19 on air pollution in California, USA. *Sci. Total Environ* 750, 141592. 10.1016/j.scitotenv.2020.141592
- Liu Q, Malarvizhi AS, Liu W, Xu H, Harris JT, Yang J, Duffy DQ, Little MM, Sha D, Lan H, Yang C (2021c). Spatiotemporal changes in global nitrogen dioxide emission due to COVID-19 mitigation policies. *Sci. Total Environ* 776, 146027. 10.1016/j.scitotenv.2021.146027
- Manousakas M, Papaefthymiou H, Diapouli E, Migliori A, Karydas AG, Bogdanovic-Radovic I, Eleftheriadis K (2017). Assessment of PM_{2.5} sources and their corresponding level of uncertainty in a coastal urban area using EPA PMF 5.0 enhanced diagnostics. *Sci. Total Environ* 574, 155–164. 10.1016/j.scitotenv.2016.09.047 [PubMed: 27631196]
- Mendoza DL, Benney TM, Bares R, Crosman ET (2021). Intra-city variability of fine particulate matter during COVID-19 lockdown: A case study from Park City, Utah. *Environ. Res* 201, 111471. 10.1016/j.envres.2021.111471
- Miech JA, Herckes P, Fraser MP (2021). Effect of COVID-19 travel restrictions on Phoenix air quality after accounting for boundary layer variations. *Atmos. Environ* 10, 100105. 10.1016/j.aeoa.2021.100105
- Mishra G, Ghosh K, Dwivedi AK, Kumar M, Kumar S, Chintalapati S, Tripathi SN (2021). An application of probability density function for the analysis of PM_{2.5} concentration during the COVID-19 lockdown period. *Sci. Total Environ* 782, 146681. 10.1016/j.scitotenv.2021.146681
- Miyazaki K, Bowman K, Sekiya T, Takigawa M, Neu JL, Sudo K, Osterman G, Eskes H (2021). Global tropospheric ozone responses to reduced NO_x emissions linked to the COVID-19 worldwide lockdowns. *Sci. Adv* 7, eabf7460. 10.1126/sciadv.abf7460
- Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med.* 6, e1000097. 10.1371/journal.pmed.1000097

- Mousazadeh M, Paital B, Naghdali Z, Mortezaia Z, Hashemi M, Karamati Niaragh E, Aghababaei M, Ghorbankhani M, Lichtfouse E, Sillanpää M, Hashim KS, Emamjomeh MM (2021). Positive environmental effects of the coronavirus 2020 episode: A review. *Environ. Dev. Sustain* 23, 12738–12760. 10.1007/s10668-021-01240-3 [PubMed: 33558801]
- Muhammad S, Long X, Salman M (2020). COVID-19 pandemic and environmental pollution: A blessing in disguise? *Sci. Total Environ* 728, 138820. 10.1016/j.scitotenv.2020.138820
- Naeger AR, Murphy K (2020). Impact of COVID-19 containment measures on air pollution in California. *Aerosol Air Qual. Res* 20, 2025–2034. 10.4209/aaqr.2020.05.0227
- Otmani A, Benchrif A, Tahri M, Bounakhla M, Chakir EM, El Bouch M, Krombi M (2020). Impact of Covid-19 lockdown on PM₁₀, SO₂ and NO₂ concentrations in Salé City (Morocco). *Sci. Total Environ* 735, 139541. 10.1016/j.scitotenv.2020.139541
- Pan S, Jung J, Li Z, Hou X, Roy A, Choi Y, Gao HO (2020). Air quality implications of COVID-19 in California. *Sustainability* 12, 7067. 10.3390/su12177067
- Parker HA, Hasheminassab S, Crouse JD, Roehl CM, Wennberg PO (2020). Impacts of Traffic Reductions Associated With COVID-19 on Southern California Air Quality. *Geophys. Res. Lett* 47. 10.1029/2020GL090164
- Peel JL, Klein M, Dana Flanders W, Mulholland JA, Tolbert PE (2010). Impact of improved air quality during the 1996 Summer Olympic Games in Atlanta on multiple cardiovascular and respiratory outcomes. Health Effects Institute, Boston, Mass., USA. <https://www.healtheffects.org/system/files/Peel148.pdf>
- Pepe E, Bajardi P, Gauvin L, Privitera F, Lake B, Cattuto C, Tizzoni M (2020). COVID-19 outbreak response, a dataset to assess mobility changes in Italy following national lockdown. *Sci Data* 7, 230. 10.1038/s41597-020-00575-2 [PubMed: 32641758]
- Peters MDJ, Godfrey CM, Khalil H, McInerney P, Parker D, Soares CB (2015). Guidance for conducting systematic scoping reviews. *Int. J. Evid. Based Healthc* 13, 141–146. 10.1097/XEB.0000000000000050 [PubMed: 26134548]
- Phillips CA, Caldas A, Cleetus R, Dahl KA, Declat-Barreto J, Licker R, Merner LD, Ortiz-Partida JP, Phelan AL, Spanger-Siegfried E, Talati S, Trisos CH, Carlson CJ (2020). Compound climate risks in the COVID-19 pandemic. *Nat. Clim. Chang* 10, 586–588. 10.1038/s41558-020-0804-2
- Porter C (1997). Changes in Air Quality and Transportation Associated With the 1996 Atlanta Summer Olympics. National Cooperative Highway Research Program. M. C. S. I. Cambridge.
- Raifman J, Nocka K, Jones D, Bor J, Lipson S, Jay J, Cole M, Krawczyk N, Benfer EA, Chan P, Galea S (2022). COVID-19 US State Policy Database. Ann Arbor, MI: Inter-university Consortium for Political and Social Research. 10.3886/E119446V143
- Richter A, Eyring V, Burrows JP, Bovensmann H, Lauer A, Sierk B, Crutzen PJ (2004). Satellite measurements of NO₂ from international shipping emissions. *Geophys. Res. Lett* 31, L23110. 10.1029/2004GL020822
- Russell AR, Valin LC, Cohen RC (2012). Trends in OMI NO₂ observations over the United States: Effects of emission control technology and the economic recession. *Atmos. Chem. Phys* 12, 12197–12209. 10.5194/acp-12-12197-2012
- Sahraei MA, Ku kapan E, Çodur MY (2021). Public transit usage and air quality index during the COVID-19 lockdown. *J. Environ. Manage* 286, 112166. 10.1016/j.jenvman.2021.112166
- Sannigrahi S, Kumar P, Molter A, Zhang Q, Basu B, Basu AS, Pilla F (2021). Examining the status of improved air quality in world cities due to COVID-19 led temporary reduction in anthropogenic emissions. *Environ. Res* 196, 110927. 10.1016/j.envres.2021.110927
- Savtchenko AK, Khayat MG (2021). NO₂ anomalies - Economy attribution and rapid climate response. *Atmos. Environ* 254, 118351. 10.1016/j.atmosenv.2021.118351
- Schlosser F, Maier BF, Jack O, Hinrichs D, Zachariae A, Brockmann D (2020). COVID-19 lockdown induces disease-mitigating structural changes in mobility networks. *PNAS* 117, 32883–32890. 10.1073/pnas.2012326117 [PubMed: 33273120]
- Shakoar A, Chen X, Farooq TH, Shahzad U, Ashraf F, Rehman A, Yan W (2020). Fluctuations in environmental pollutants and air quality during the lockdown in the USA and China: Two sides of COVID-19 pandemic. *Air Qual. Atmos. Health* 13, 1335–1342. 10.1007/s11869-020-00888-6 [PubMed: 32837622]

- Shehzad K, Bilgili F, Koçak E, Xiaoxing L, Ahmad M (2021). COVID-19 outbreak, lockdown, and air quality: fresh insights from New York City. *Environ. Sci. Pollut. Res* 28, 41149–41161. 10.1007/s11356-021-13556-8
- Sokhi RS, Singh V, Querol X, Finardi S, Targino AC, Andrade M. de F., Pavlovic R, Garland RM, Massagué J, Kong S, Baklanov A, Ren L, Tarasova O, Carmichael G, Peuch VH, Anand V, Arbilla G, Badali K, Beig G, Belalcazar LC, et al. (2021). A global observational analysis to understand changes in air quality during exceptionally low anthropogenic emission conditions. *Environ. Int* 157, 106818. 10.1016/j.envint.2021.106818
- Straka W, Kondragunta S, Wei Z, Zhang H, Miller SD, Watts A (2020). Examining the economic and environmental impacts of COVID-19 using earth observation data. *Remote Sens.* 13, 5. 10.3390/rs13010005
- Tanzer-Gruener R, Li J, Eilenberg SR, Robinson AL, Presto AA (2020). Impacts of modifiable factors on ambient air pollution: A case study of COVID-19 shutdowns. *Environ. Sci. Technol. Lett* 7, 554–559. 10.1021/acs.estlett.0c00365 [PubMed: 37566291]
- The People's Government of Beijing Municipality (2009). Public notice on temporary traffic control measures during the 2008 Beijing Olympic and Paralympic Games. The People's Government of Beijing Municipality, China.
- Turner AJ, Kim J, Fitzmaurice H, Newman C, Worthington K, Chan K, Wooldridge PJ, Köehler P, Frankenberg C, Cohen RC (2020). Observed impacts of COVID-19 on urban CO₂ emissions. *Geophys. Res. Lett* 47, e2020GL090037. 10.1029/2020GL090037
- Venter ZS, Aunan K, Chowdhury S, Lelieveld J (2020). COVID-19 lockdowns cause global air pollution declines. *PNAS* 117, 18984–18990. 10.1073/pnas.2006853117 [PubMed: 32723816]
- Venter ZS, Aunan K, Chowdhury S, Lelieveld J (2021). Air pollution declines during COVID-19 lockdowns mitigate the global health burden. *Environ. Res* 192, 110403. 10.1016/j.envres.2020.110403
- Wang Q, Li S (2021). Nonlinear impact of COVID-19 on pollutions—Evidence from Wuhan, New York, Milan, Madrid, Bandra, London, Tokyo and Mexico City. *Sustainable Cities Soc.* 65, 102629. 10.1016/j.scs.2020.102629
- Winchester AK, Peterson RA, Carter E, Sammel MD (2021). Impact of COVID-19 social distancing policies on traffic congestion, mobility, and NO₂ pollution. *Sustainability* 13, 7275. 10.3390/su13137275
- Wong MS, Zhu R, Yin Tung Kwok C, Kwan MP, Santi P, Ho Liu C, Qin K, Lee KH, Heo J, Li H, Ratti C (2021). Association between NO₂ concentrations and spatial configuration: A study of the impacts of COVID-19 lockdowns in 54 US cities. *Environ. Res. Lett* 16, 054064. 10.1088/1748-9326/abf396
- World Health Organization (WHO) (2020). Coronavirus Disease 2019 (COVID-19) Situation Report.
- Wu JT, Leung K, Leung GM (2020). Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: A modelling study. *Lancet* 395, 689–697. 10.1016/S0140-6736(20)30260-9 [PubMed: 32014114]
- Xiang J, Austin E, Gould T, Larson T, Shirai J, Liu Y, Marshall J, Seto E (2020). Impacts of the COVID-19 responses on traffic-related air pollution in a Northwestern US city. *Sci. Total Environ* 747, 141325. 10.1016/j.scitotenv.2020.141325
- Yusup Y, Ramli NK, Kayode JS, Yin CS, Hisham S, Mohamad Isa H, Ahmad MI (2020). Atmospheric carbon dioxide and electricity production due to lockdown. *Sustainability* 12, 9397. 10.3390/su12229397
- Zambrano-Monserrate MA, Ruano MA, Sanchez-Alcalde L (2020). Indirect effects of COVID-19 on the environment. *Sci. Total Environ* 728, 138813. 10.1016/j.scitotenv.2020.138813
- Zangari S, Hill DT, Charette AT, Mirowsky JE (2020). Air quality changes in New York City during the COVID-19 pandemic. *Sci. Total Environ* 742, 140496. 10.1016/j.scitotenv.2020.140496
- Zhang H, Lin Y, Wei S, Loo BPY, Lai PC, Lam YF, Wan L, Li Y (2021). Global association between satellite-derived nitrogen dioxide (NO₂) and lockdown policies under the COVID-19 pandemic. *Sci. Total Environ* 761, 144148. 10.1016/j.scitotenv.2020.144148

Zhang Z, Arshad A, Zhang C, Hussain S, Li W (2020). Unprecedented temporary reduction in global air pollution associated with COVID-19 forced confinement: A continental and city scale analysis. *Remote Sens.* 12, 2420. 10.3390/rs12152420

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Fig. 1. PRISMA Flow chart of scoping review inclusion and exclusion criteria.




Fig. 2. A thematic map of number of studies included in this scoping review, by air pollutants. Studies could examine more than one pollutant. The numbers reflect the total number of studies that examined a particular pollutant.




Fig. 3. Location-specific summary of magnitude and direction of change in air pollutants.




Fig. 4. Methodologies utilized in this scoping review, by location and air pollutant.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1.

List of articles included in this scoping review.

Author	Journal	Published date	Study Period	Study Location	Pollutants
Amouei Torkmahalleh <i>et al.</i> (2021)	Aerosol and Air Quality Research	1/28/2021	1/1/2020/4/30/2020	Global	NO ₂ , PM _{2.5} , O ₃
Naeger and Murphy (2020)	Aerosol and Air Quality Research	8/10/2020	3/9/2020/4/10/2020	California	NO ₂ , PM _{2.5}
Shakoor <i>et al.</i> (2020)	Air Quality, Atmosphere & Health	8/9/2020	1/18/2020/3/25/2020	Global	CO, NO ₂ , PM _{2.5} , PM ₁₀
Habibi <i>et al.</i> (2020)	Atmosphere	11/27/2020	1/1/2020/4/30/2020	Global	CO, NO ₂ , PM _{2.5} , O ₃
Fu <i>et al.</i> (2020)	Atmosphere	11/3/2020	1/23/2020/5/10/2020	Global	CO, NO ₂ , PM _{2.5} , PM ₁₀ , O ₃ , SO ₂
Jia <i>et al.</i> (2020)	Atmosphere	6/14/2020	3/25/2020/4/23/2020	Memphis, Tennessee	NO ₂ , PM _{2.5} , O ₃
Keller <i>et al.</i> (2021)	Atmospheric Chemistry and Physics	1/21/2021	1/1/2020/7/1/2020	Global	NO ₂ , O ₃
Jiang <i>et al.</i> (2021)	Atmospheric Chemistry and Physics	6/9/2021	2/18/2020/4/23/2020	Southern California	CO, NH ₃ , NO _x , PM _{2.5} , PM ₁₀ , SO _x
Miech <i>et al.</i> (2021)	Atmospheric Environment	4/1/2021	1/6/2020/4/8/2020	Phoenix, Arizona	CO, NO ₂ , PM ₁₀
Savtchenko and Khayat (2021)	Atmospheric Environment	6/1/2021	10/1/2004/4/30/2020	Global	NO ₂
Bray <i>et al.</i> (2021)	Atmospheric Environment	6/21/2021	1/1/2020/4/30/2020	Global	CO, NO ₂ , PM _{2.5} , PM ₁₀ , O ₃ , SO ₂
Campbell <i>et al.</i> (2021)	Atmospheric Environment	11/1/2021	3/1/2020/9/30/2020	Continental US	O ₃
Ghosal and Saha (2021)	Atmospheric Environment	6/1/2021	1/1/2020/6/29/2020	Continental US	PM _{2.5}
Archer <i>et al.</i> (2020)	Bulletin of Atmospheric Science and Technology	10/26/2020	3/31/2020/4/30/2020	Continental US	NO ₂ , PM _{2.5}
Bar <i>et al.</i> (2021)	Cities	10/1/2021	3/1/2020/5/30/2020	Europe and US	NO ₂ , PM _{2.5}
Balasubramaniam <i>et al.</i> (2020)	Energy Sources, Part A: Recovery, Utilization, and Environmental Effects	10/1/2020	3/1/2020/5/30–2020	US, Italy, and France	NO ₂ , PM _{2.5} , O ₃
Hoang <i>et al.</i> (2021)	Energy Sources, Part A: Recovery, Utilization, and Environmental Effects	3/1/2021	NA	Global	NO ₂
Mousazadeh <i>et al.</i> (2021)	Environment, Development, and Sustainability	2/4/2021	NA	Global	CO, NO, NO ₂ , PM _{2.5} , PM ₁₀ , O ₃ , SO ₂
Sokhi <i>et al.</i> (2021)	Environment International	12/1/2021	1/1/2020/9/30–2020	Global	CO, NO, NO ₂ , PM _{2.5} , PM ₁₀ , O ₃ , SO ₂
El-Sayed <i>et al.</i> (2021)	Environmental Pollution	9/15/2021	3/15/2020/4/15/2020	Florida	CO, NO ₂ , PM _{2.5} , O ₃ , SO ₂
Acharya <i>et al.</i> (2021)	Environmental Research	2/1/2021	3/22/2020/4/30/2020	Southeast Asia, Europe, and the US	AOD, NO ₂ , SO ₂

Author	Journal	Published date	Study Period	Study Location	Pollutants
Sannigrahi <i>et al.</i> (2021)	Environmental Research	3/4/2021	2/1/2020-5/11/2020	Global	CO, NO ₂ , PM _{2.5} , PM ₁₀
Venter <i>et al.</i> (2021)	Environmental Research	1/1/2021	1/1/2020–5/15/2020	Global	NO ₂ , PM _{2.5} , O ₃
Mendoza <i>et al.</i> (2021)	Environmental Research	10/1/2021	2/3/2020–7/23/2020	Park City, Utah	PM _{2.5}
Chauhan and Singh (2020)	Environmental Research	8/1/2020	12/1/2019–3/31/2020	Global	PM _{2.5}
Wong <i>et al.</i> (2021)	Environmental Research Letters	5/11/2021	2/15/2020–5/6/2020	Continental US	NO ₂
Shehzad <i>et al.</i> (2021)	Environmental Science and Pollution Research	3/29/2021	1/1/2020–5/3/2020	New York City, New York	CO, NO ₂ , PM _{2.5} , O ₃ , SO ₂
Liu <i>et al.</i> (2021a)	Environmental Science and Technology Letters	12/4/2020	1/1/2020–4/30/2020	California	NO, NO ₂
Tanzer-Gruener <i>et al.</i> (2020)	Environmental Science and Technology Letters	6/1/2020	3/1/2020–3/31/2020	Pittsburg, PA	CO, NO ₂ , PM _{2.5}
Bauwens <i>et al.</i> (2020)	Geophysical Research Letters	6/16/2020	1/1/2020–4/30/2020	Global	NO ₂
Turner <i>et al.</i> (2020)	Geophysical Research Letters	10/30/2020	2/2/2020–5/2/2020	San Francisco, CA	CO ₂
Parker <i>et al.</i> (2020)	Geophysical Research Letters	11/1/2020	3/19/2020–6/30/2020	South Coast Air Basin (SoCAB), California	NO ₂ , NO _x , PM _{2.5} , O ₃
Goldberg <i>et al.</i> (2020)	Geophysical Research Letters	8/17/2020	1/1/2020–4/30/2020	Continental US	NO ₂
Irfan <i>et al.</i> (2021)	International Journal of Environmental Research	1/1/2021	NA	China, USA, Italy, and Spain	NO ₂
Connerton <i>et al.</i> (2020)	International Journal of Environmental Research and Public Health	7/14/2020	3/1/2020–3/31/2020	São Paulo, Brazil; Paris, France; Los Angeles, California; and New York	CO, NO ₂ , PM _{2.5} , O ₃
Chadwick <i>et al.</i> (2021)	Journal of Aerosol Science	6/1/2021	2/11/2020–5/11/2020	Salt Lake City, Utah	PM _{2.5}
Sahraei <i>et al.</i> (2021)	Journal of Environmental Management	5/1/2021	1/15/2020–5/31/2020	Global	CO, NO ₂ , PM _{2.5} , PM ₁₀ , O ₃ , SO ₂
Dang <i>et al.</i> (2021)	Journal of Environmental Economics and Management	1/1/2021	1/1/2020–3/31/2020	Global	NO ₂ , PM _{2.5}
Brodeur <i>et al.</i> (2021)	Journal of Environmental Economics and Management	3/1/2020	3/19/2020–5/31/2020	Continental US	PM _{2.5}
Kondragunta <i>et al.</i> (2021)	Journal of Geophysical Research: Atmosphere	8/17/2021	2/11/2020–5/11/2020	Continental US	NO _x
Liu <i>et al.</i> (2020)	Nature Communications	10/14/2020	1/1/2020–6/30/2020	Global	CO ₂
Kerr <i>et al.</i> (2021)	Proceedings of the National Academy of Sciences	8/19/2021	3/13/2020–6/13/2020	Continental US	NO ₂
Venter <i>et al.</i> (2020)	Proceedings of the National Academy of Sciences	7/28/2020	1/1/2020–5/31/2020	Global	AOD, NO ₂ , O ₃
Straka <i>et al.</i> (2021)	Remote Sensing	12/22/2020	2/1/2020–4/30/2020	Los Angeles, California; Chicago, Illinois; Washington DC	NO ₂
Zhang <i>et al.</i> (2020)	Remote Sensing	7/1/2020	1/1/2020–5/15/2020	Global	AOD, CO, NO ₂ , PM _{2.5} , O ₃ , SO ₂

Author	Journal	Published date	Study Period	Study Location	Pollutants
Elshorbany <i>et al.</i> (2021)	Remote Sensing	1/1/2021	1/1/2020–12/31/2020	New York, Illinois, Florida, Texas, and California	AOD, CO, NO ₂ , O ₃
Miyazaki <i>et al.</i> (2021)	Science Advances	6/9/2021	2/1/2020–7/31/2020	Global	NO _x , O ₃
Hammer <i>et al.</i> (2021)	Science Advances	6/23/2021	1/1/2020–4/30/2020	Global	PM _{2.5}
Chossiere <i>et al.</i> (2021)	Science Advances	5/21/2021	1/1/2020–7/6/2020	Global	NO ₂ , PM _{2.5} , O ₃
Liu <i>et al.</i> (2021c)	Science of the Total Environment	7/1/2021	1/1/2020–6/30/2020	China, USA, UK, Brazil, South Africa, and India	NO ₂
Zhang <i>et al.</i> (2021)	Science of the Total Environment	3/20/2021	1/1/2020–4/30/2020	Australia, Canada, China, and United States	NO ₂
Liu <i>et al.</i> (2021b)	Science of the Total Environment	1/1/2021	1/26/2020–6/14/2020	California	CO, NO ₂ , PM _{2.5} , PM ₁₀ , O ₃ , SO ₂
Muhammad <i>et al.</i> (2020)	Science of the Total Environment	8/1/2020	NA	Global	NO ₂
Mishra <i>et al.</i> (2021)	Science of the Total Environment	8/15/2021	3/24/2020–6/15/2020	India, China, France, Brazil and United States	PM _{2.5}
Zangari <i>et al.</i> (2020)	Science of the Total Environment	11/1/2020	1/1/2020–5/30/2020	New York	NO ₂ , PM _{2.5}
Xiang <i>et al.</i> (2020)	Science of the Total Environment	12/1/2020	2/17/2020–3/31/2020	Seattle, Washington	BC, CO, NO, NO ₂ , NO _x , PM _{2.5}
Chen <i>et al.</i> (2020)	Science of the Total Environment	7/21/2020	1/1/2020–4/30/2020	Continental US	CO, NO ₂ , PM _{2.5} , PM ₁₀ , O ₃
Zambrano-Monserrate <i>et al.</i> (2020)	Science of the Total Environment	8/1/2020	NA	China, USA, Italy, and Spain	NO ₂
Berman and Ebisu (2020)	Science of the Total Environment	10/1/2020	1/8/2020–4/21/2020	Continental US	NO ₂ , PM _{2.5}
Hudda <i>et al.</i> (2020)	Science of the Total Environment	11/10/2020	3/27/2020–5/14/2020	Somerville, Massachusetts	BC, PNC
Bekbulat <i>et al.</i> (2021)	Science of the Total Environment	5/15/2020	1/1/2020–10/1/2020	Continental US	CO, NO ₂ , PM _{2.5} , PM ₁₀ , O ₃
Heintzelman <i>et al.</i> (2021)	Sustainability	8/12/2021	3/1/2020–4/30/2020	Continental US	NO ₂
Winchester <i>et al.</i> (2021)	Sustainability	6/29/2021	1/1/2020–8/31/2020	Continental US	NO ₂
Pan <i>et al.</i> (2020)	Sustainability	8/1/2020	3/20/2020–5/30/2020	California	PM _{2.5} , O ₃
Yusup <i>et al.</i> (2020)	Sustainability	11/1/2020	3/1/2020–5/31/2020	China, the United States, Europe, and India	CO ₂
Wang and Li (2021)	Sustainable Cities and Society	2/1/2021	1/23/2020–10/30/2020	Wuhan, New York, Milan, Madrid, Bandra, London, Tokyo and Mexico City	NO ₂ , PM _{2.5} , O ₃ , SO ₂