

Supplemental Table 1: Number, weighted number and weighted percentage of maternal ART-use, infant feeding practices and infant nevirapine use amongst mothers with consecutive follow-up visits at 6 weeks, 3 months, 6 months, 9 months, 12 months, 15 months and 18 months

	6-weeks Unweighted number/ Weighted number (%)	3 months Unweighted number/ Weighted number (%)	6 months Unweighted number/ Weighted number (%)	9 months Unweighted number/ Weighted number (%)	12 months Unweighted number/ Weighted number (%)	15 months Unweighted number/ Weighted number (%)	18-months Unweighted number/ Weighted number (%)
Maternal ART-use	951/1637 (52.4)	892/1637 (55.7)	715/1257 (56.9)	774/1266 (59.0)	765/1236 (58.9)	765/1230 (59.0)	785/1246 (59.5)
HIV exposed infant on nevirapine prophylaxis	2001/2127 (93.2)	1176/1770 (65.0%)	739/1687 (41.9)	455/1661 (24.8)	267/1647 (15.0)	139/1671 (7.1)	62/1787 (3.1)
Any breastfeeding	1056/1603 (64.1)	876/1603 (40.2)	538/1236 (29.6)	364/1182 (19.8)	260/1050 (13.7)	150/1141 (9.2)	93/1195 (5.2)