MORBIDITY AND MORTALITY WEEKLY REPORT

INDEX ISSUE

INDEX - Volume 27, 1978

Each issue (number) of Volume 27 of the Morbidity and Mortality Weekly Report contains the following pages, inclusive. In the index, numbers in bold face are errata.

Issue	Pages	Issue	Pages	Issue	Pages	Issue	Pages
. 1	1-8	14	115-122	28	235-246	41	391-402
2	9-16	15	123-130	29	247-258	42	403-414
3	17-24	16	131-138	30	259-270	43	415-426
4	25-32	17	139-146	31	271-282	44	427-438
5	33-40	18	147-154	32	283-294	45	439-450
6	41-48	19	155-166	33	295-306	46	451-462
7	49-56	20	167-174	34	307-318	47	463-474
8	57-64	21	175-182	35	319-330	48	475-486
9	65-72	22	183-190	36	331-342	49	487-498
10	73-80	23	191-198	37	343-354	50	499-510
10*	81-90	24	199-206	38	355-366	51	511-522
11	91-98	25	207-214	39	367-378	52	523-538
12	99-106	26	215-222	40	379-390	53	539-550*
13	107-114	27	223-234				

^{*}Supplement. Supplements to the MMWR are:

Chemoprophylaxis of Malaria, 10 March 1978. Issued as supplement to No. 10

Health Information for International Travel 1978. Issued September 1978

Reported Morbidity and Mortality in the United States, 1973. Annual Supplement Summary, to be published in mid-1979 as No. 54

Index. 15 January 1979. Issued as No. 53

A

Abortion: associated with consumption of pennyroyal oil 511; complications of 71, 175; deaths associated with 71, 175; patterns among teenagers 460; surveillance in 1976, 175

Abscesses: associated with DTP vaccine 331

ACIP recommendation: on cholera vaccine 173, influenza vaccine 205, 285, 351, 447, measles prevention 427, meningococcal polysaccharide vaccines 327, plague vaccine 255, pneumococcal polysaccharide vaccine 25, rubella vaccine 451, smallpox vaccine 156, typhoid vaccine 231, typhus vaccine 189, vaccines for international travel 449, yellow fever vaccine 268

Adhesive bandages: Elastoplast, associated with Rhizopus infections 33

Amies transport medium: associated with falsepositive Gram stains 23

Antibiotic resistance: of Mycobacterium tuberculosis 355, Neisseria gonorrhoeae 10, 121, Plasmodium falciparum 84, 221, 463, pneumococcus 1

Antihemophilic factor: recovery of non-A, non-B hepatitis virus from 199

Arboviral activity: in humans 375, in United States 375, Caribbean 375, Central America 375. See also California encephalitis, Dengue fever, Eastern equine encephalitis, Japanese B encephalitis, St. Louis encephalitis, Western equine encephalitis.

Arsine poisoning: associated with maintenance workers at a chemical plant 165; in Georgia 165

Ascaris lumbricoides: prevalence of, in U.S. stool survey 167

Asphyxia: fatal, among fishermen 309

B

Bacteremia: caused by Klebsiella pneumoniae 307, Pseudomonas aeruginosa 307, 308; in a dialysis unit 307 Bandages: associated with *Rhizopus* infections 33, 80, 190, 243

Beans: associated with Clostridium perfringens food poisoning 164

Birth control: used by teenagers 460. See also Family planning.

Birth defects: caused by Rh hemolytic disease 487; congenital rubella syndrome 495, 522

Births: among teenagers in 1976, 460

Blindness: incidence of, worldwide 279; prevention of 279; remediable 279

Botfly infestation: in participants of an archaeologic expedition in Guatemala 405

Botulism: associated with country club restaurant 138, 145, fish 356, honey 17, olives 501, potato salad 138, 145, 483, 498, soil 17, three-bean salad 138, 145, vacuum cleaner dust 17; caused by type A toxin 17, 138, 174, 250, 466, 501, type B toxin 17, 250; in infants 17, 100, 249; outbreak 138, 145, 174, 466, 483, 501; in California 483, 501, Colorado 466, 483, 498, New Mexico 138, 145, 174, Puerto Rico 356, Texas 138, 174, United States 17, 105, 138, 145, 174, 466, 501, England 100. See also Infant botulism.

Burn injuries: epidemiologic features of 124; in New York 124

Burritos: associated with Clostridium perfringens food poisoning 164

C

California encephalitis: fatal 279, 375; in Arkansas 375, Illinois 375, Minnesota 279, 375, New York 375, Ohio 375, Tennessee 375, Wisconsin 279, 375

Campylobacter fetus infection: caused by C. fetus sub. jejuni 207, 226; milk-borne 226; waterborne 207; in Colorado 226, Vermont 207

Cardiopulmonary resuscitation (CPR) training: manikins' risk of transmitting hepatitis and other disease 132, 154, 247

Chemical poisoning: associated with traffic accident 50; caused by 1,3-dichloropropene 50; long-term effects of 50; in California 50

Cholera: caused by serogroup O1, 341, 351, 366, 367, 388, 402, 413, 442, 520; endemic 351, 365, 367, 388, 402, 413, 442; treatment for

413; vaccine 173, 413; in Louisiana 341, 351, 365, 367, 388 (map), 402, 413, 442, 520, United States 181, 341, 351, 365, 367, 402, 413, 442, 520, Africa 181, Asia 181, Canada 181, Europe 181, Oceania 181, worldwide 181

Cholera vaccine: ACIP recommendation concerning 173, 449

Ciguatera fish poisoning: associated with red snapper 39; in British Virgin Islands 39

Clostridium perfringens food poisoning: in California 164

Coccidioidomycosis: associated with windstorm 55, 106; in California 55, 106

Congenital rubella syndrome: in United States from 1977-1978, 495, 522

Conjunctivitis: caused by silver nitrate prophylaxis 107, 258

Contraception: See Family planning.

Crab: associated with cholera outbreak 341, 367, 388, 402, 413; recommendations for boiling before eating 402. See also Cholera.

Cruise ship: gastrointestinal illness aboard 65

Culture-collection devices: associated with falsepositive Gram stains 23

D

Decontamination: of CPR training manikins 132, 154, 247

Dengue fever: caused by type 1, 74, 183, 225, 304, type 2, 74, 183, 224, 304, type 3, 74, 183, 304; hemorrhagic 476; in Puerto Rico 183, 224, 304, United States 74, 183, 304, 376, 476, Caribbean 74, 183, 304, 376, Central America 376. See also Dengue hemorrhagic fever.

Dengue hemorrhagic fever: in Ohio resident of Indian descent 476; WHO's clinical classification of 478; in Southeast Asia 477

Diarrheal illness: associated with herbal teas 248.

See also Gastrointestinal illness, Gastroenteritis.

Drug resistance: See Antibiotic resistance.

DTP vaccine: reactions associated with 331, 471

Eastern equine encephalitis: in Florida 375, United States 214, 375, Dominican Republic 213

Echo 9 infection: associated with aseptic meningitis 393, 507, pharyngitis in children 393; in Maryland 501, New York 392

Elastoplast bandages: associated with Rhizopus infections 33, 80, 190, 243

Encephalitis: See California encephalitis, Eastern equine encephalitis, Japanese B encephalitis, St. Louis encephalitis, Western equine encephalitis.

Entamoeba histolytica: prevalence of, in U.S. stool survey 167

Enterobius vermicularis: prevalence of, in U.S. stool survey 167

F

Falls: injuries due to, studies of 192

False-positive tests: associated with Amies transport medium 23

Family planning: and teenage childbearing 131, 460; excess births in 7 southeastern states as a guide for 73

Fertility: excess 73, 131; of teenagers 460

Floods: and vaccination for typhoid fever 500

Food poisoning: associated with bean-filled burritos 164, marijuana-contaminated cakes 404; caused by Clostridium perfringens 164. See also Botulism, Infant botulism.

G

Gases: toxic, produced by farm's liquid manure system 47

Gastroenteritis: associated with a sewage leak 183, summer camps 403; caused by Campy-lobacter fetus sub. jejuni 207, 226; of possible viral etiology 183, 404, 450; waterborne 403, 450; in Arkansas 183, Missouri 183, Pennsylvania 403, 450

- Gastrointestinal illness: aboard a naval vessel 237, the S/S Statendam 65; associated with burritos 164, herbal teas 248, hydroquinone poisoning 237, seafood salad 65, water 65; caused by Vibrio parahaemolyticus 65. See also Gastroenteritis.
- Genetic disease: congenital rubella 495, 522; Rh hemolytic 487
- Giardia lamblia: most frequently identified organism in U.S. stool survey 167
- Giardiasis: waterborne 155; in Colorado 155, Michigan 155, New York 155. See also Giardia lamblia.
- Gonococcal ophthalmia neonatorum: and silver nitrate prophylaxis 107, 258
- Gonorrhea: gonococcal ophthalmia neonatorum as complication of 107, 258; in 1977, 98, 166; in 1978, 448; penicillinase-producing 10, 121; in Philippines 10, United States 10, 98, 166, 448, England 10, worldwide 10

H

- Heart valve prostheses: contaminated porcine 92
- Hemodialysis: associated with outbreak of bacteremia and pyrogenic reactions 307
- Hepatitis: non-A, non-B, recovery of virus-like particles of 199; photograph of 199; risk of acquiring in CPR training classes 132, 154, 247
- Herbal teas: associated with diarrhea 248. See also Pennyroyal oil.
- Honey: associated with infant botulism 18, 249
- Hydrogen sulfide poisoning: associated with farm's liquid manure system 47; in Wisconsin 47
- Hydroquinone: poisoning 237

- Immunization laws: and nationwide Immunization Initiative 7; enforcement of 1, 245, 282, 303; in Michigan 1
- Indiana Memorial Union: associated with Legionnaires' disease 191, 216, 283, 317

- Infant botulism: associated with honey 18, 249; isolation of toxin from serum from patient with 411; in Arizona 411, California 17, 18, 249, United States 17, 100, 249, 411, England 100
- Infantile periarteritis nodosa: and Kawasaki disease 9
- Influenza: A/FM/1/47(H1N1) 8, 16; A/FW/1/50. 16; A/Brazil/11/78, 521; A/Texas/1/77, 8, 16, 32, 40, 48, 56, 64, 80, 114, 224, 285; A/USSR/77, 16, 24, 32, 40, 48, 55, 64, 72, 80, 91, 114, 224, 285, 437, 462, 520; A/Victoria/3/75-like 24, 32, 40, 48, 56, 80, 114, 285; B/Hong Kong/5/72, 8, 16, 285; caused by type A 8, 16, 24, 32, 40, 48, 56, 64, 72, 80, 91, 114, 224, 285, 437, 462, 474, 484, 507, 521, 537, type B 8, 24, 32, 40, 114, 285, 437, 462, type C 507; mortality due to pneumonia-influenza, charts of 32, 48, 64, 80, 91, 114, 473; in Arizona 8, 16, 72, 521, 537, California 55, 64, 438, 474, 507, 521, 537, Colorado 8, 16, 48, 64, 537, Connecticut 24, 32, 72, Delaware 24, 64, District of Columbia 24, 40, 55, 64, Florida 24, 32, 522, 537, Georgia 24, 72, 537, Illinois 8, 16, 24, 32, 56, 64, 537, Indiana 24, 32, Louisiana 64, Maine 32, Maryland 16, 24, 32, 64, Massachusetts 24, Michigan 16, 24, 48, 64, Missouri 8, 16, 64, New Hampshire 32, New Jersey 8, 16, 24, 32, 56, 64, New Mexico 16, 537, New York 8, 24, 32, 56, 64, 507, North Carolina 24, 32, North Dakota 64, Ohio 24, 64, Oregon 91, 537, Pennsylvania 8, 24, 32, 64, Philippines 24, 438, Puerto Rico 40, 484, Rhode Island 24, 32, South Carolina 64, Tennessee 8, 56, 64, Texas 64, 114, 462, 521, 537, Utah 521, 537, Vermont 32, 72, Virginia 24, 72, Washington 522, Wisconsin 8, 16, 32, 64, Wyoming 40, 48, 64, United States 8, 16, 24, 32, 40, 48, 55, 64, 72, 80, 91, 114, 285, 462, 474, 507, 521, 537, Argentina 224, Australia 437, 438, 462, Brazil 224, Bulgaria 48, Canada 56, 80, 438, Canal Zone 91, Chile 224, Czechoslovakia 48, Ecuador 224, England 32, 40, 48, 522, Finland 8, France 537, Germany 91, Hong Kong 8, 16, 24, 437, 462, Hungary 48, 91, 538, Italy 91, Jamaica 522, Japan 48, 91, Malaysia 462, Netherlands 91, New Zealand 437, 462, Norway 40, People's Republic of China 24, Scotland 522, Singapore 522, South America 224, Spain 537, Sweden 40. Taiwan 8, 24, 32, 64, 438, Thailand 522, United Kingdom 24, 32, 40, 48, U.S.S.R. 8, 16, 538, worldwide 40, 48, 64, 91, 521, 537. See also Influenza vaccine, Russian flu.
- Influenza vaccine: ACIP statement on 205, 285, 351; American Academy of Pediatrics' recommendation on, for high-risk children 447; shortage of 447

Injuries: due to falls 192

J

Japanese B encephalitis: in India 449, 464

K

Kawasaki disease: and infantile periarteritis nodosa 9; in United States 9, Japan 9, worldwide 9

ŀ

Lead poisoning: adult 191; associated with ore smelter 57, stripping older homes 191; causing encephalopathy 8; investigation of effects of, on Texas community 57; surveillance of childhood, 4th quarter FY 1977, 66, 71, 1st quarter 1978, 137, 2nd quarter FY 1978, 339; in Massachusetts 191, New Jersey 8, Texas 57

Legionnaires' disease: associated with air conditioning 283, 368, 415, 439, country club 293, 415, Indiana Memorial Union 191, 216, 283, 317, N.Y. garment district 352, 365, Spanish resorts 205, 523, Veterans of Foreign Wars Convention 387, water 283, 317, 368, 439; bacterium, environmental isolation of 283, 368, 415, 439, identification of Togus strain of 293; international symposium on 176; outbreaks of, in Atlanta 293, 415, 439, Bloomington 191, 216, 283, 317, 435, Los Angeles 394, 439, Memphis 353, 365, 387, 439, New York City 352, 365, 439; seasonal distribution of 441; sporadic cases of 439; in Georgia 293, 415, Indiana 191, 216, 283, 317, Missouri 387, New Jersey 387, New York 352, 365, 387, Tennessee 353, 365, 368, Washington 387, United States 439, Australia 523, England 200, Scotland 200, 205, Wales 200, worldwide 523

Leukemia: childhood, cluster of 129; in New York 129

Liquid manure system: farm worker's death associated with toxic gases from 47 Liquid protein diet: deaths associated with 223; FDA regulations concerning 223

Lymphocyte dysfunction: associated with polybrominated biphenyl exposure 207

M

Malaria: acquired in Kenya 214, 258, 463, Tanzania 463; caused by *Plasmodium falciparum* 81, 214, 221, 332, 463, *P. malariae* 81, 273, 333, *P. ovale* 81, 333, *P. vivax* 81, 332; chemoprophylaxis 81 (supplement), 221, 274; chloroquine-resistant strains of 84, 221, 463; fatal 214, 258, 332; risk by country 86, 221; surveillance summary for 1976, 41, for 1977, 332; in California 214, Georgia 463, New York 463, United States 41, 214, 332, 463, Denmark 463, Grenada 273, worldwide 86, 221

Mammoplasty: augmentation, associated with postoperative mycobacterial infections 513

Manure: associated with hydrogen sulfide poisoning 47

Marijuana: foodborne outbreak of 404

Measles: ACIP recommendation concerning prevention of 427; associated with ball pythons 321, school outbreaks 41, 303, 321, 489, 510, 519; fatal 424; goal to eliminate from United States by October 1, 1982, 391; in children with religious exemptions from vaccination 401; in Indiana 369, Maryland 401, Michigan 41, North Carolina 244, Oregon 519, Texas 489, 510, Utah 344, United States in 1972-1976, 39, in 1976, 39, in 1977, 39, 235, 303, in 1978, 235, 282, 303, 424. See also Immunization laws.

Meningitis: aseptic 393, 507; associated with echovirus 9, 393, 508; in Maryland 508, New York 392

Meningococcal disease: associated with alcoholics 121; caused by serogroup A 121, serogroup B 121, C 121, Y 121, 147, W135, 121, 154; hospital-acquired 147; in cancer patients 147; vaccine recommendations for 327; in Alaska 121, 154, Connecticut 147

Meningococcemia: nosocomial 358; in Wisconsin 358

Meningoencephalitis: primary amebic 343

Mexican food: associated with Clostridium perfringens food poisoning 164 MMWR

545

Milk: associated with bovine tuberculosis 108

MLNS: See Kawasaki disease.

Mucocutaneous lymph node syndrome: See Kawasaki disease.

Mucormycosis: nosocomial outbreak of, associated with Elastoplast wound dressings 33, 80

Mumps: surveillance summary for 1976, 379; in United States 379

Mycobacterial infection: associated with augmentation mammoplasty 513, porcine heart valves 92; caused by Mycobacterium chelonei 92, M. fortuitum complex 513

Mycoplasmal pneumonia: in singing group 271, boy campers 271; in Wisconsin 271

Myiasis: cutaneous 405

THE YEAR OF BUILDING THE PROPERTY OF

N

National Center for Health Statistics: joint study with CDC on efficacy of nosocomial infection control (SENIC) 139

Neisseria gonorrhoeae: See Gonorrhea.

stockers with a few control of the

Nematodes: prevalence of, in U.S. stool survey 167

Nosocomial infections: associated with Elastoplast bandages 33, 80, 190, 243, porcine heart valve prostheses 92; caused by adenovirus type 4, 49, Group Y Neisseria meningitidis 147, Mycobacterium chelonei 92, respiratory syncytial virus 260; exposure of pregnant women to, by medical personnel 123; in an intensive care nursery 260, cancer patients 147; SENIC phase I study of 139; U.S. hospitals' attempts to control 139. See also Cardiopulmonary resuscitation (CPR) training.

0

Olives: associated with botulism outbreak 501

Umin 11 ft (Amery \$14) bussed by Journage Character \$14, in Q v Year \$15.05 whereast

PAM: See Primary amebic meningoencephalitis.

Parainfluenza infection: in extended-care facilities 475; initially diagnosed and reported as Russian flu 198

Paralytic shellfish poisoning: in Washington 416

Parasites: intestinal, surveillance of in United States 167

Parvovirus-like agents: associated with waterborne gastroenteritis 403

PBB: See Polybrominated biphenyls.

PCB: See Polychlorinated biphenyls.

Penicillin: resistance of gonococci 10, 121, pneumococci 1

Pennyroyal oil: used unsuccessfully as an abortifacient 511

Pesticide poisoning: acute and possible long-term effects of 50

Pharyngoconjunctival fever: caused by adenovirus type 4, 49; in New York 49

Photograph: of virus-like particles associated with non-A, non-B hepatitis 199

Plague: bubonic 259; in Arizona 259, California 259, New Mexico 259

Plague vaccine: ACIP recommendations concerning 255, 449

Pneumococcus: See Streptococcus pneumoniae.

Pneumonia: influenza deaths in 121 U.S. cities, charts of 32, 48, 64, 80, 91, 114, 473

Poisoning: acute arsine 165; associated with bean-filled burritos 164, farm's liquid manure system 47, hydrogen sulfide 47, polybrominated biphenyls in dairy feed 115, 207, polychlorinated biphenyls in sludge 99, red snapper 39; childhood lead 66, 137, 484; hydroquinone 237; of fisherman associated with hydrogen sulfide, ammonia, carbon dioxide, and sodium bisulfite in ship's hold 310. See also Food poisoning, Lead poisoning, Pesticide poisoning.

Poliomyelitis: caused by type 1 poliovirus 222, 273, 294, 536; in Dutch visitor to Canada and United States 273, 294; in Canada 273, 294, 381, Netherlands 222, 273, 294, 381, 536

Poliovirus vaccine: type used, in Netherlands 222, 536

Polybrominated biphenyls: accidentally introduced into dairy feed 115, 207; prospective Michigan cohort study of humans exposed to 115; study of lymphocyte dysfunction in Michigan persons exposed to 207 Polychlorinated biphenyls: contamination of sludge with 99

Primary amebic meningoencephalitis: caused by Acanthamoeba species 343, Naegleria fowleri 343; non-fatal 343; in California 343, Florida 343. New York 343

Prostheses: contaminated heart valve 92; silicone mammary, associated with mycobacterial infections 513

Psittacosis: in a pet store owner 417; traced to pet birds 417; in Connecticut 417

Pythons: associated with Q fever 321

Q

Q fever: associated with ball pythons 321, ticks 321; in New York 321

Quarantine measures: See Vaccination certificate requirements.

R

Rabbit paws: associated with tularemia 105

Rabies: animal 224, 399, 426, 447, 465, 499; canine 224; fatal 267, 306; human 267, 306, 465, 499; in bats 500, raccoons 465, 500, skunks 399, 426, 465, 500, squirrels 447; sources of exposure for, in 20 human cases 267, 306; vaccine-induced in dogs 224; in Alabama 447, 465, Florida 465, Georgia 465, South Carolina 465, Texas 267, 306, United States 499

Rabies vaccine: available at CDC 333, 413; human diploid cell strain 333, 413

Raccoons: increasing reports of rabies in 465

Recycling: of farm waste associated with worker's death 47

Relapsing fever: in a camper 526; in California 526

Respiratory illness: in Wisconsin 271

Respiratory syncytial virus infection: in an intensive care nursery 260; in California 260

Reye syndrome: and influenza B 15; in United States 15

RH hemolytic disease: in United States from 1968-1977, 487

Rhizopus infection: associated with Elastoplast bandages 33, 80, 190, 243; caused by R. oryzae 33, 80, 190, 243; R. rhizopodiformis 243; in Minnesota 33, 80, 190, 243, United States 33, 80, 190, 243

Rift Valley fever: in Egypt 113

Roast beef: associated with salmonellosis 315; USDA recommendations for cooking 315

Rocky Mountain spotted fever: in Arkansas 113, Georgia 113, Maryland 113, North Carolina 113, Oklahoma 113, South Carolina 113, Tennessee 113, Virginia 113, United States in 1977, 113

Roundworms: prevalence of, in U.S. stool survey 167

RSV: See Respiratory syncytial virus infection.

Rubella: ACIP recommendation concerning vaccine for 451; and congenital rubella syndrome 495, 522; exposure of pregnant women to 123; outbreak among medical personnel 123; in California 123. United States 495

Rubeola: See Measles.

Russian flu: hemagglutination and neuraminidase cross-reactions of 16; mistakenly reported from Oregon 91, 198; similar to A/FM/1/47, 8, 16, A/FW/1/50, 16; in United States 40, 48, 55, 64, 72, 80, 91, 114, 198, 462, Australia 462, Asia 64, Bulgaria 48, Czechoslovakia 48, Europe 64, 91, Germany 91, Hungary 48, 91, Italy 91, Japan 48, 91, Malaysia 462, Netherlands 91, People's Republic of China 24, 64, Taiwan 24, 32, 64, United Kingdom 24, 32, 40, 48, U.S.S.R. 8, worldwide 48, 64, 91. See also Influenza.

S

Salad: associated with botulism 138, 174, Vibrio parahaemolyticus outbreak on a Caribbean cruise 65

Salmonellosis: associated with precooked roast beef 315, turkey 514; caused by Salmonella muenster 514; in New York 315, Washington 514

Seafood: salad associated with Vibrio parahaemo. lyticus outbreak on a Caribbean cruise 65. See also Crab, Paralytic shellfish poisoning, Shrimp.

- SENIC Project: and results of Preliminary Screening Questionnaire 139
- Sewage: leak from lagoon associated with gastroenteritis 183
- Shigellosis: caused by Shigella flexneri 79, S. sonnei 79; decline in cases of, caused by S. flexneri 79; surveillance summary for 1976, 79: in United States 79
- Shrimp: associated with Vibrio parahaemolyticus foodborne outbreak 345; "dip" associated with fatal asphyxia among fishermen 307
- Silver nitrate: as prophylaxis for gonococcal ophthalmia neonatorum 107, 258
- Skunks: and CDC recommendation against keeping when captured in the wild 400, 426; associated with rabies 399, 426, 465
- Sludge: contaminated with polychlorinated biphenyls 99
- Smallpox: airborne 319, 341; laboratory-associated 319, 341, 346, 415; United States ends requirement for vaccination for 295; WHO \$1,000 award for case of 415; in Bangladesh 8, England 319, 341, 346, 364, Ethiopia 8, 153, Kenya 8, 153, Somalia 8, 153, 415, worldwide 8, 153, 319, 341, 346, 415. See also Smallpox vaccine, Vaccination certificate requirements.
- Smallpox vaccine: ACIP recommendations for use of 156, 449
- S/S Statendam: gastrointestinal illness aboard 65
- St. Louis encephalitis: in United States 375
- Streptococcus pneumoniae: penicillin-resistant 1; multiply-resistant 1; vaccine for 25; in South Africa 1
- Surveillance summary: on abortion in 1976, 175, intestinal parasites in 1976, 167, malaria in 1976, 41, in 1977, 332, mumps in 1976, 379, rabies in 1977, 499, shigellosis in 1976, 79
- **Syphilis:** in United States in 1977, 31, 63, in 1978, 105, 130, 190, 296, 411, 423, 536

in this registrical province. based on workly rel.

The recording week concludes at close of business of

- Telone II: See Pesticide poisoning.
- Tick-borne encephalitis: acquired in Hungary 164; caused by *Ixodes ricinus* 164; in Ohio 164, eastern Europe 164

- Tick paralysis: in Colorado 297, New Mexico 297
- Trichuris trichiura: prevalence of, in U.S. stool survey 167
- Trust Territory of the Pacific Islands: begins reporting morbidity and mortality data to CDC 370
- Tuberculosis: and criteria for reporting/counting cases of 317; associated with a college 145, foreign students 145, immigrants 145, nursing home 523; caused by *Mycobacterium bovis* 108, 145; foreign-acquired 108, 145; with resistance to isoniazid, para-aminosalicylic acid, and streptomycin 355; in California 145, Maryland 108, Mississippi 355, North Dakota 523, United States 63, 316
- Tularemia: associated with rabbits 105; pneumonic 105, 352, 366; on Martha's Vineyard 352, 366, in Massachusetts 352, 366, Washington 105
- Typhoid fever: ACIP recommendation concerning vaccine for 231, 449; vaccination requirements during natural disasters 500
- Typhus fever: endemic 215; laboratory-acquired 215; louse-borne (epidemic), ACIP recommendation concerning vaccine for 189, 449; in Maryland 215

V

Vaccination certificate requirements: and changes in names of countries in Afars and the Issas, French Territory of the (to Djibouti, Republic of) 7, Cambodia (to Democratic Kampuchea) 56, Central African Republic (to Central African Empire) 56, Laos (to Lao People's Democratic Republic) 154. Libyan Arab Republic (to Libyan Arab Jamahiriya) 154; changes in cholera vaccination requirements in Afghanistan 56, Brunei 56, China, People's Republic of 153, Guinea-Bissau 154, Italy 154, Kuwait 154, Mozambique 154, Oman 376, Seychelles 376, Saudi Arabia 377, United States 367; changes in smallpox vaccination requirements in Afghanistan 56, Antiqua 56, Argentina 376, Australia 364, Bahrain 56, Bulgaria 56, 153, Chile 56, China, Republic of (Taiwan) 7, Colombia 54, Cook Islands 58, Czechoslovakia 58, France 113, French Guiana 106, Gabon 106, Gambia 106, German Democratic Republic (East) 106, Germany, Federal Republic of (West) 106, 154, Gibraltar 106, Guadeloupe 106, Guatemala 106, Guernsey,

Alderney, and Sark 106, Guinea-Bissau 106, Guyana 7, 106, Honduras 154, Hong Kong 364, Israel 154, Italy 154, Jamaica 154, Japan 364, Jordan 154, Liberia 221, Macao 122. Malaysia 122, Malta 364, Martinique 122, Mauritius 122, 364, Mexico 122, Monaco 122, Morocco 364, Niue 122, Oman 376, Peru 122, Poland 130, Reunion 154, Romania 130, Saint Helena 221, 377, Saudi Arabia 377, Singapore 364, Trinidad and Tobago 353, Turkey 364, United States 295; changes in vellow fever vaccination requirements in Afghanistan 56, Canal Zone 56, China, People's Republic of 153, Cuba 58, India 154, Indonesia 154, Macao 122, Malaysia 122, Montserrat 154, Niue 122, Pakistan 122, Samoa (American) 154, Saudi Arabia 377

Vaccine: for cholera 173, 449, DTP 331, 471, influenza 205, 285, 351, louse-borne (epidemic) typhus fever 189, 449, measles 427, meningococcal disease 327, plague 255, 449, poliomyelitis 222, rabies 224, 333, 413, rubella 451, smallpox 156, 295, 449, Streptococcus pneumoniae 25, typhoid fever 231, 449, yellow fever 58, 268, 449; induced rabies 224

Varicella-Zoster Immune Globulin: availability of 508

Vibrio cholerae infection: in Louisiana 341, 351, 365, 367, 388, 389, 402, 412, 520

Vibrio parahaemolyticus infection: associated with shrimp 345; foodborne 345; in Louisiana

January 15, 1979



Western equine encephalitis: in United States 376

Windstorm: associated with coccidioidomycosis 55, 106

Wound infections: associated with augmentation mammoplasty 513, porcine heart valve implantation 92; caused by mycobacteria 92, 513

Y

Yellow fever: jungle 509; in The Gambia 520, Trinidad 509; on Colombia-Venezuela border 255

Yellow fever vaccine: ACIP recommendation concerning 268, 449. See also Vaccination certificate requirements.

The Morbidity and Mortality Weekly Report, circulation 84,000, is published by the Center for Disease Control, Atlanta, Georgia. The data in this report are provisional, based on weekly telegraphs to CDC by state health departments. The reporting week concludes at close of business of Friday; compiled data on a national basis are officially released to the public on the succeeding Friday.

The editor welcomes accounts of interesting cases, outbreaks, environmental hazards, or other public health problems of current interest to health officials. Send reports to: Center for Disease Control, Attn: Editor, Morbidity and Mortality Weekly Report, Atlanta, Georgia 30333.

Send mailing list additions, deletions, and address changes to: Center for Disease Control, Attn: Distribution Services, GSO, 1-SB-36, Atlanta, Georgia 30333. When requesting changes be sure to give your former address, including zip code and mailing list code number, or send an old address label.

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE PUBLIC HEALTH SERVICE / CENTER FOR DISEASE CONTROL ATLANTA, GEORGIA 30333 OFFICIAL BUSINESS

Director, Center for Disease Control William H. Foege, M.D. Director, Bureau of Epidemiology Philip S. Brachman, M.D. Editor Michael B. Gregg, M.D. Managing Editor Anne D. Mather, M.A. Postage and Fees Paid U.S. Department of HEW HEW 396

AIR MAIL



T.5

Principles of Control for