

**Supplemental Digital Content 3: Table of Corrected Mortality Estimates at 1 Year of Follow-up among 306,335 Adults Starting ART in Mozambique during 2004–2013\***

Year of ART initiation	No. Lost by 1 year	No. Dead by 1 year	Denominator (No. Starting ART in Calendar Year)	Mortality at 1 Year of ART Follow-up				LTFU at 1 year		Retention at 1 year
				Uncorrected	(95% Confidence Interval)**	Corrected	(95% Confidence Interval)†	Uncorrected	Corrected	
2004	453	156	1,784	9.0%	(7.7-10.4%)	16.8%	(10.5-24.2%)	25.9%	18.2%	65.0%
2005	967	281	4,694	6.1%	(5.4-6.8%)	13.9%	(8.6-19.8%)	20.9%	13.1%	73.0%
2006	2,915	860	12,634	6.9%	(6.5-7.4%)	15.0%	(9.2-21.7%)	23.5%	15.0%	70.0%
2007	6,305	1,529	25,616	6.1%	(5.8-6.4%)	14.5%	(8.5-21.7%)	25.0%	16.5%	69.0%
2008	6,840	1,499	26,704	5.7%	(5.5-6.0%)	14.4%	(8.2-21.8%)	26.1%	17.6%	68.0%
2009	6,503	1,664	31,129	5.5%	(5.2-5.7%)	13.5%	(8.1-19.5%)	21.4%	13.5%	73.0%
2010	7,483	1,704	37,449	4.7%	(4.4-4.9%)	12.6%	(7.4-18.4%)	20.5%	12.4%	75.0%
2011	9,892	1,567	41,991	3.8%	(3.6-4.0%)	12.7%	(6.8-19.5%)	24.0%	15.3%	72.0%
2012	17,325	1,648	58,892	2.9%	(2.8-3.0%)	12.7%	(5.9-21.4%)	31.0%	21.3%	66.0%
2013	15,159	892	65,442	2.0%	(1.8-2.1%)	11.2%	(5.4-17.9%)	42.6%	33.3%	55.5%
Overall	73,842	11,800	306,335	4.3%	(4.2-4.4%)	13.1%	(7.1-20.1%)	26.8%	18.0%	68.9%

Abbreviations: ART, antiretroviral therapy; LTFU, lost to follow-up.

\*Corrected using the published web-based nomogram available at: <http://www.iedea-sa.org/index.php?id=2785>.

\*\*Confidence intervals were estimated using the stcomp package in STATA and do not account for clustering.

†Confidence intervals were estimated through the web-based nomogram at: <http://www.iedea-sa.org/index.php?id=2785>.