

Appendix Table. In vitro susceptibilities of 92 isolates of group G *Streptococcus*, April 1998–August 2004, Taiwan

Antimicrobial agent	Range	MIC ($\mu\text{g/mL}$) [*]		Susceptible isolates, %
		MIC ₉₀	MIC ₅₀	
Penicillin	0.03–0.12	0.06	0.03	100
Cefepime	0.03–0.5	0.06	0.03	100
Imipenem	0.03	0.03	0.03	100
Ertapenem	0.03–0.5	0.06	0.03	100
Azithromycin	0.03→128	128	0.12	67.4
Clarithromycin	0.03→128	4	0.06	73.9
Telithromycin	0.03–32	0.5	0.03	96.7†
Clindamycin	0.06→128	64	0.25	87.0
Ciprofloxacin	0.06–1	0.5	1	100
Levofloxacin	0.06–1	0.5	1	100
Moxifloxacin	0.06–0.5	0.25	0.12	100
Vancomycin	0.12–1	0.5	0.25	100
Quinupristin-dalfopristin	0.5–4	4	2	33.7
Linezolid	1–2	2	2	100
Tigecycline	0.03–1	0.5	0.12	100‡

^{*}MIC₅₀, 50% inhibition; MIC₉₀, 90% inhibition.†Defined as MICs <0.5 $\mu\text{g/mL}$.‡Defined as MICs <2 $\mu\text{g/mL}$.