

Italy - Medical Students 2010 (3rd Year Students Only) Global Health Professions Student Survey (GHPSS)

FACT SHEET

The Italy medical students GHPSS includes data on prevalence of cigarette smoking and use of other tobacco products, knowledge and attitudes about tobacco use, exposure to second-hand smoke, desire for smoking cessation, and training received during medical school regarding patient counseling on smoking cessation techniques.

This information could be very useful in monitoring and developing Italy's comprehensive tobacco control program. The Italy medical students GHPSS was a sample of third year medical students attending 9 medical schools in Italy conducted in 2010. A total of 730 3rd year students participated in the Italy medical students GHPSS. The overall response rate of all medical students surveyed was 41.3%.

Prevalence

64.9% ever smoked cigarettes (Males = 69.3%, Females = 62.0%)
20.0% currently smoke cigarettes (Males = 22.2%, Females = 18.6%)
18.8% ever used any form of tobacco other than cigarettes (Males = 26.8%, Females = 13.5%)
3.6% currently use any form of tobacco other than cigarettes (Males = 5.6%, Females = 2.3%)
31.6% of ever smokers smoked on school premises/property during the past year
10.8% of ever smokers smoked in school buildings during the past year

Factors Influencing Tobacco Use

28.4% were exposed to second-hand smoke at home, during the past week.
75.5% were exposed to second-hand smoke in public places during the past week
89.1% reported that their school had a ban on smoking in school buildings and clinics
79.1% reported that their school enforced the ban on smoking in school buildings and clinics
97.5% thought smoking should be banned in all enclosed public places
87.9% thought tobacco sales to adolescents should be banned
82.4% thought there should be a complete ban on advertising tobacco products
60.8% of current smokers wanted to stop smoking
31.5% of current smokers ever received help/advice to stop smoking cigarettes
90.4% were taught about the dangers of smoking during medical school training

Role Model and Cessation Training

65.0% thought health professionals serve as role models for their patients
94.4% thought health professionals have a role in giving advice about smoking cessation to patients
10.6% received formal training in smoking cessation approaches during m school
87.7% thought health professionals should get specific training on cessation techniques

Highlights

64.9% currently smoke cigarettes and 3.6% currently use other tobacco products

Over six in 10 of the smokers indicated that they want to quit

Over three-quarters were exposed to smoke in public places

Almost all support a ban on smoking in all enclosed public places

One in 10 received training to provide patients with cessation approaches

87.7% think health professionals should get specific training on cessation techniques to use with patients