

Morocco - Dental Students 2010 (3rd Year Students Only) Global Health Professions Student Survey (GHPSS)

FACT SHEET

The Morocco Dental students GHPSS includes data on prevalence of cigarette smoking and use of other tobacco products, knowledge and attitudes about tobacco use, exposure to second-hand smoke, desire for smoking cessation, and training received during Dental school regarding patient counseling on smoking cessation techniques.

This information could be very useful in monitoring and developing Morocco's comprehensive tobacco control program. The Morocco Dental students GHPSS was a census of 3rd year Dental students attending two Dental schools in Morocco conducted in 2010. A total of 119 3rd year students participated in the Morocco Dental students GHPSS. The overall response rate of all Dental students surveyed was 87.1%.

Prevalence

- 27.9% ever smoked cigarettes (Males = 66.0%, Females = 13.5%)
- 4.4% currently smoke cigarettes (Males = 14.9%, Females = 1.1%)
- 25.6% ever used any form of tobacco other than cigarettes (Males = 51.6%, Females = 16.4%)
- 3.7% currently use any form of tobacco other than cigarettes (Males = 7.5%, Females = 2.5%)
- 0.9% currently smoke shisha (Males = 3.7%, Females = 0.0%)
- 18.6% of ever smokers smoked on school premises/property during the past year
- 9.9% of ever smokers smoked in school buildings during the past year

Factors Influencing Tobacco Use

- 17.6% were exposed to second-hand smoke at home, during the past week
- 64.7% were exposed to second-hand smoke in public places during the past week
- 22.0% reported that their school had a ban on smoking in school buildings and clinics
- 58.2% reported that their school enforced the ban on smoking in school buildings and clinics
- 97.5% thought smoking should be banned in all enclosed public places
- 92.5% thought tobacco sales to adolescents should be banned
- 85.9% thought there should be a complete ban on advertising tobacco products
- 42.3% were taught about the dangers of smoking during dental school training

Role Model and Cessation Training

- 84.8% thought health professionals serve as role models for their patients
- 92.9% thought health professionals have a role in giving advice about smoking cessation to patients
- 4.4% received formal training in smoking cessation approaches during dental school
- 96.7% thought health professionals should get specific training on cessation techniques

Highlights

4.4% currently smoke cigarettes and 3.7% currently use other tobacco products

Almost two-thirds were exposed to smoke in public places

Almost all support a ban on smoking in all enclosed public places

More than nine in 10 thought tobacco sales to adolescents should be banned

4.4% received training to provide patients with cessation approaches

Almost all think health professionals should get specific training on cessation techniques to use with patients