

Czech Republic - Pharmacy Students 2011 (3rd Year Students Only) Global Health Professions Student Survey (GHPSS)

FACT SHEET

The Czech Republic pharmacy students GHPSS includes data on prevalence of cigarette smoking and use of other tobacco products, knowledge and attitudes about tobacco use, exposure to second-hand smoke, desire for smoking cessation, and training received during pharmacy school regarding patient counseling on smoking cessation techniques.

This information could be very useful in monitoring and developing Czech Republic's comprehensive tobacco control program. The Czech Republic pharmacy students GHPSS was a census of third year pharmacy students attending two pharmacy schools in Czech Republic conducted in 2011. A total of 331 3rd year students participated in the Czech Republic pharmacy students GHPSS. The overall response rate of all pharmacy students surveyed was 96.0%.

Prevalence

82.7% ever smoked cigarettes (Males = 84.9%, Females = 82.3%)
22.4% currently smoke cigarettes (Males = 20.8%, Females = 22.7%)
53.5% ever used any form of tobacco other than cigarettes (Males = 64.2%, Females = 51.4%)
8.2% currently use any form of tobacco other than cigarettes (Males = 13.2%, Females = 7.2%)
17.5% of ever smokers smoked on school premises/property during the past year
0.0% of ever smokers smoked in school buildings during the past year

Factors Influencing Tobacco Use

28.7% were exposed to second-hand smoke at home, during the past week
60.3% were exposed to second-hand smoke in enclosed public places during the past week
85.0% were exposed to second-hand smoke in outdoor public places during the past week
89.4% were exposed to second-hand smoke in public places during the past week
85.5% reported that their school had a ban on smoking in school buildings and clinics
77.2% reported the ban on smoking is enforced among those who reported their school had a ban on smoking in school building and clinics
79.4% thought smoking should be banned in all enclosed public places
97.0% thought tobacco sales to adolescents should be banned
69.5% thought there should be a complete ban on advertising tobacco products
66.7% of current smokers wanted to stop smoking
14.7% of current smokers ever received help/advice to stop smoking cigarettes
91.8% were taught about the dangers of smoking during pharmacy school training

Role Model and Cessation Training

55.8% thought health professionals serve as role models for their patients
82.4% thought health professionals have a role in giving advice about smoking cessation to patients
1.5% received formal training in smoking cessation approaches during pharmacy school
71.3% thought health professionals should get specific training on cessation techniques

Highlights

22.4% currently smoke cigarettes and 8.2% currently use other tobacco products

Two-thirds of the smokers indicated that they want to quit

Three in five were exposed to smoke in enclosed public places

Almost eight in 10 support a ban on smoking in all enclosed public places

Over seven in 10 think health professionals should get specific training on cessation techniques to use with patients