

# Thailand - Dental Students 2011 (3<sup>rd</sup> Year Students Only) Global Health Professions Student Survey (GHPSS)

## FACT SHEET . . . . .

The Thailand Dental students GHPSS includes data on prevalence of cigarette smoking and use of other tobacco products, knowledge and attitudes about tobacco use, exposure to second-hand smoke, desire for smoking cessation, and training received during Dental school regarding patient counseling on smoking cessation techniques.

This information could be very useful in monitoring and developing Thailand's comprehensive tobacco control program. The Thailand Dental students GHPSS was a census of 3<sup>rd</sup> year Dental students attending 8 Dental schools in Thailand conducted in 2011. A total of 577 students participated out of which 566 belonged to 3<sup>rd</sup> year. The overall response rate of all Dental students surveyed was 84.6%.

### Prevalence of Tobacco Use

- 13.9% ever smoked cigarettes (Males = 27.8%, Females = 7.8%)
- 2.5% currently smoke cigarettes (Males = 6.8%, Females = 0.5%)
- 16.8% ever used any form of tobacco other than cigarettes (Males = 25.9%, Females = 12.5%)
- 2.3% currently use any form of tobacco other than cigarettes (Males = 2.8%, Females = 2.1%)
- 5.9% of ever smokers smoked on school premises/property during the past year.
- 0.0% of ever smokers smoked in school buildings during the past year.

### Factors Influencing Tobacco Use

- 32.4% were exposed to second-hand smoke at home, during the past week
- 49.8% were exposed to second-hand smoke in enclosed public places during the past week
- 61.5% were exposed to second-hand smoke in outdoor public places during the past week
- 68.0% reported that their school had a ban on smoking inside and outside the whole perimeter of their school buildings and clinics
- 92.3% reported that their school enforced the ban on smoking inside and outside the whole perimeter of their school buildings and clinics
- 98.9% thought smoking should be banned in all enclosed public places
- 94.9% thought tobacco sales to adolescents should be banned
- 89.5% thought there should be a complete ban on advertising tobacco products
- 15.3% of current smokers ever received help/advice to stop smoking cigarettes<sup>†</sup>
- 89.8% were taught about the dangers of smoking during dental school training

### Role Model and Cessation Training

- 98.3% thought health professionals serve as role models for their patients
- 96.0% thought health professionals have a role in giving advice about smoking cessation to patients
- 32.7% received formal training in smoking cessation approaches during dental school
- 91.9% thought health professionals should get specific training on cessation techniques

(<sup>†</sup> The percentage is based on observations between 10 and 34)

### Highlights

**2.5% currently smoke cigarettes and 2.3% currently use any form of tobacco other than cigarettes**

**Half were exposed to smoke in public places**

**Almost all support a ban on smoking in all enclosed public places**

**One-third received training to provide patients with cessation approaches**

**Over nine in 10 think health professionals should get specific training on cessation techniques to use with patients**

