

Thailand - Nursing Students 2011 (3rd Year Students Only) Global Health Professions Student Survey (GHPSS)

FACT SHEET

The Thailand Nursing students GHPSS includes data on prevalence of cigarette smoking and use of other tobacco products, knowledge and attitudes about tobacco use, exposure to second-hand smoke, desire for smoking cessation, and training received during Nursing school regarding patient counseling on smoking cessation techniques.

This information could be very useful in monitoring and developing Thailand's comprehensive tobacco control program. The Thailand Nursing students GHPSS was a sample of 3rd year Nursing students attending 14 Nursing schools in Thailand conducted in 2011. A total of 1,917 students participated out of which 1,845 belonged to 3rd year. The overall response rate of all Nursing students surveyed was 91.2%.

Prevalence of Tobacco Use

- 9.8% ever smoked cigarettes (Males = 33.2%, Females = 8.6%)
- 0.5% currently smoke cigarettes (Males = 6.6%, Females = 0.2%)
- 14.1% ever used any form of tobacco other than cigarettes (Males = 33.2%, Females = 12.9%)
- 1.9% currently use any form of tobacco other than cigarettes (Males = 6.1%, Females = 1.7%)
- 3.6% of ever smokers smoked on school premises/property during the past year.
- 3.3% of ever smokers smoked in school buildings during the past year

Factors Influencing Tobacco Use

- 22.7% were exposed to second-hand smoke at home, during the past week
- 36.2% were exposed to second-hand smoke in enclosed public places during the past week
- 45.1% were exposed to second-hand smoke in outdoor public places during the past week
- 79.0% reported that their school had a ban on smoking inside and outside the whole perimeter of their school buildings and clinics
- 98.4% reported that their school enforced the ban on smoking inside and outside the whole perimeter of their school buildings and clinics
- 99.4% thought smoking should be banned in all enclosed public places
- 96.7% thought tobacco sales to adolescents should be banned
- 93.2% thought there should be a complete ban on advertising tobacco products
- 92.0% were taught about the dangers of smoking during nursing school training

Role Model and Cessation Training

- 99.8% thought health professionals serve as role models for their patients
- 97.9% thought health professionals have a role in giving advice about smoking cessation to patients
- 29.7% received formal training in smoking cessation approaches during nursing school
- 97.9% thought health professionals should get specific training on cessation techniques

Highlights

0.5% currently smoke cigarettes and 1.9% currently use any form of tobacco other than cigarettes

36.2% were exposed to smoke in public places

Almost all support a ban on smoking in all enclosed public places

Three in 10 received training to provide patients with cessation approaches

Almost all think health professionals should get specific training on cessation techniques to use with patients

