

Slovenia (Ages 13-15)

Global Youth Tobacco Survey (GYTS)

FACT SHEET



The Slovenia GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Slovenia could include in a comprehensive tobacco control program.

The Slovenia GYTS was a school-based survey of students in primary years 8 and 9, and secondary year 1 conducted in

2007. A two-stage cluster sample design was used to produce representative data for all of Slovenia. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 95.0%, the class response rate was 97.3%, and the student response rate was 87.5%, and the overall response rate was 80.8%. A total of 2,516 students aged 13-15 participated in the Slovenia GYTS.

Prevalence

- 57.3% of students had ever smoked cigarettes (Boys = 56.1%, Girls = 56.7%)
- 21.8% currently use any tobacco product (Boys = 16.9%, Girls = 24.2%)
- 20.3% currently smoke cigarettes (Boys = 15.2%, Girls = 23.0%)
 - 6.9% currently daily cigarette smokers (Boys = 4.8%, Girls = 8.0%)
 - 6.4% currently smoke cigars (Boys = 6.4%, Girls = 5.5%)
- 23.9% ever smokers initiated smoking before age ten (Boys = 27.8%, Girls = 19.8%)
- 20.6% of never smokers are likely to initiate smoking next year (Boys = 17.2%, Girls = 24.1%)

Access and Availability - Current Smokers

- 9.9% usually smoke at home
- 66.3% buy cigarettes in a store
- 71.2% who bought cigarettes in a store were NOT refused purchase because of their age

Exposure to Secondhand Smoke (SHS)

- 75.9% think smoking should be banned from public places
- 63.3% think smoke from others is harmful to them
- 45.1% have one or more parents who smoke
- 16.3% have most or all friends who smoke

Cessation - Current Smokers

- 39.7% want to stop smoking
- 67.0% tried to stop smoking during the past year
- 44.2% have ever received help to stop smoking
- 11.6% always have or feel like having a cigarette first thing in the morning

Media and Advertising

- 73.6% saw anti-smoking media messages vs. 43.4% saw pro media messages on TV
- 51.8% saw anti messages vs. 46.0% saw pro messages on billboards
- 53.4% saw anti smoking ads vs. 51.8% saw pro-cigarette ads in newspapers or magazines
- 13.3% have an object with a cigarette brand logo
- 7.0% were offered free cigarettes by a tobacco company representative

School

- 61.6% had been taught in class, during the past year, about the dangers of smoking
- 49.1% had discussed in class, during the past year, reasons why people their age smoke
- 51.3% had been taught in class, during the past year, the effects of tobacco use

Highlights

- 21.8% of students currently use any form of tobacco; 20.3% currently smoke cigarettes; 6.9% are current daily smokers; 6.4% currently smoke cigars; 1 in 5 never smokers likely to initiate smoking next year.
- SHS exposure is high – over 2 in 5 have at least a parent who smokes; 16.3% have most or all their friends smoke
- Close to two-thirds of the students think smoke from others is harmful to them.
- Three-quarters of the students think smoking in public places should be banned.
- More than 1 in 7 students have an object with a cigarette brand logo on it.
- Almost three-quarters of students saw anti-smoking messages on TV vs. more than 4 in 10 saw pro messages; over half saw anti messages on billboards vs. close to 5 in 10 saw pro messages.
- Six in 10 students were taught in school about dangers of smoking.