

Sri Lanka (Ages 13-15)

Global Youth Tobacco Survey (GYTS)

FACT SHEET



The Sri Lanka GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Sri Lanka could include in a comprehensive tobacco control program.

The Sri Lanka GYTS was a school-based survey of students in grades 8 through 10 conducted in 2007. A two-stage cluster

sample design was used to produce representative data for all of Sri Lanka. At the first stage all schools were selected to participate in the survey. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 100%, the class response rate was 100%, the student response rate was 85%, and the overall response rate was 85%. A total of 1,428 students aged 13-15 participated in the Sri Lanka GYTS.

Prevalence

- 5.1% of students had ever smoked cigarettes (Boy = 6.9%, Girl = 3.4%)
- 9.1% currently use any tobacco product (Boy = 12.4%, Girl = 5.8%)
- 1.2% currently smoke cigarettes (Boy = 1.6%, Girl = 0.9%)
- 8.6% currently use other tobacco products (Boy = 11.6%, Girl = 5.6%)
- 3.7% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

- 36.0% think boys and 16.9% think girls who smoke have more friends
- 29.5% think boys and 26.1% think girls who smoke look more attractive

Exposure to Secondhand Smoke (SHS)

- 35.4% live in homes where others smoke in their presence
- 65.9% are around others who smoke in places outside their home
- 87.9% think smoking should be banned from public places
- 77.8% think smoke from others is harmful to them
- 29.9% have one or more parents who smoke
- 3.3% have most or all friends who smoke

Media and Advertising

- 84.5% saw anti-smoking media messages, in the past 30 days
- 67.4% saw pro-cigarette ads on billboards, in the past 30 days
- 68.4% saw pro-cigarette ads in newspapers or magazines, in the past 30 days
- 5.7% have an object with a cigarette brand logo
- 3.0% were offered free cigarettes by a tobacco company representative

School

- 72.8% had been taught in class, during the past year, about the dangers of smoking
- 35.7% had discussed in class, during the past year, reasons why people their age smoke
- 56.0% had been taught in class, during the past year, the effects of tobacco use

Highlights

- 9.1% of students currently use any form of tobacco; 1.2% of students currently smoke cigarettes and 8.6% currently use some other form of tobacco.
- SHS exposure is high – over one-third of the students live in homes where others smoke and almost two-thirds of the students are exposed to smoke around others outside of the home; 3 in 10 students have at least one parent who smokes and 3.3% of students have friends who smoke.
- More than three-quarters of the students think smoke from others is harmful to them.
- Nearly 9 in 10 students think smoking in public places should be banned.
- 5.7% of students have an object with a cigarette brand logo.
- Over 8 in 10 students saw anti-smoking media messages in the past 30 days; more than two-thirds of the students saw pro-cigarette ads in the past 30 days.