



HHS Public Access

Author manuscript

Am J Obstet Gynecol. Author manuscript; available in PMC 2025 January 03.

Published in final edited form as:

Am J Obstet Gynecol. 2021 August ; 225(2): 183.e1–183.e16. doi:10.1016/j.ajog.2021.02.028.

Urban-Rural Differences in Pregnancy-Related Deaths, United States, 2011–2016

Peter T. MERKT, MPH,
Michael R. KRAMER, PhD,
David A. GOODMAN, PhD,
Mary D. BRANTLEY, MPH,
Chloe M. BARRERA, MPH,
Lindsay ECKHAUS, MPH,
Emily E. PETERSEN, MD

National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention (Ms. Brantley, Ms. Eckhaus, Mr. Merkt, and Drs. Goodman and Petersen); and the Department of Epidemiology, Rollins School of Public Health, Emory University (Ms. Barrera and Dr. Kramer), Atlanta, Georgia.

Abstract

Background: The U.S. pregnancy-related mortality ratio has not improved over the past decade and includes striking disparities by race and ethnicity and by state. Understanding differences in pregnancy-related mortality across and within urban and rural areas can guide the development of interventions for preventing future pregnancy-related deaths.

Objective: We sought to compare pregnancy-related mortality across and within urban and rural counties by race and ethnicity and age.

Study Design: We conducted a descriptive analysis of 3747 pregnancy-related deaths during 2011–2016 (the most recent available data) with available zone improvement plan code or county data in the Pregnancy Mortality Surveillance System, among Hispanic and non-Hispanic White, Black, American Indian or Alaska Native, and Asian or Pacific Islander women aged 15 to 44 years. We aggregated data by US county and grouped counties per the National Center for Health Statistics Urban-Rural Classification Scheme for Counties. We used R statistical software, *epitools*, to calculate the pregnancy-related mortality ratio (number of pregnancy-related deaths per 100,000 live births) for each urban-rural grouping, obtain 95% confidence intervals, and perform exact tests of ratio comparisons using the Poisson distribution.

Results: Of the total 3747 pregnancy-related deaths analyzed, 52% occurred in large metro counties, and 7% occurred in noncore (rural) counties. Large metro counties had the lowest pregnancy-related mortality ratio (14.8; 95% confidence interval, 14.2–15.5) whereas noncore counties had the highest (24.1; 95% confidence interval, 21.4–27.1), including race and ethnicity

Corresponding author: Toby Merkt, yzq1@cdc.gov, 4770 Buford Hwy NE, Atlanta, GA 30341.

CDC Disclaimer: The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

and age groups. Pregnancy-related mortality ratio age disparities increased with rurality. Women aged 25 to 34 years and aged 35 to 44 years living in noncore counties had pregnancy-related mortality ratios 1.5 and 3 times higher, respectively, than women of the same age groups in large metro counties. Within each urban-rural category, pregnancy-related mortality ratios were higher among non-Hispanic Black women than non-Hispanic White women. Non-Hispanic American Indian or Alaska Native pregnancy-related mortality ratios in small metro, micropolitan, and noncore counties were 2 to 3 times that of non-Hispanic White women in the same areas.

Conclusion: Although more than half of pregnancy-related deaths occurred in large metro counties, the pregnancy-related mortality ratio rose with increasing rurality. Disparities existed in urban-rural categories, including by age group and race and ethnicity. Geographic location is an important context for initiatives to prevent future deaths and eliminate disparities. Further research is needed to better understand reasons for the observed urban-rural differences and to guide a multifactorial response to reduce pregnancy-related deaths.

Keywords

maternal mortality; maternal death; pregnancy-related death; pregnancy mortality surveillance system; policy; age disparities; racial and ethnic disparities; noncore; metropolitan

Introduction

Pregnancy-related complications claim the lives of approximately 700 women annually in the United States; approximately two-thirds of these deaths are preventable.^{1,2} Over the past decade, the pregnancy-related mortality ratio (PRMR, the number of pregnancy-related deaths per 100,000 live births) has not improved.³ There are well-documented disparities in pregnancy-related deaths: non-Hispanic Black women and American Indian or Alaska Native women have a PRMR 2 to 3 times higher than non-Hispanic White women, and the PRMR varies among states, from fewer than 10 to more than 30 deaths per 100,000 births.^{4,5}

Rural areas of the United States have unique health challenges. All-cause mortality in rural areas was 18.9% higher than that of urban areas in 2016.⁶ As measured by the hospital discharge data from deliveries, the likelihood of severe maternal morbidity and mortality in similar groups is 9% higher in rural areas than that of urban areas.⁷ Access to hospital-based and community-based obstetrical care in rural areas declined during 1995–2015, which may have affected pregnancy outcomes.^{8,9,10}

There are large racial and ethnic variations in population distributions in urban and rural areas. Non-Hispanic White individuals make up most rural (79%) and suburban (68%) populations. The proportion of the population who are non-Hispanic Black is twice as large in urban counties (17%) than that of rural counties (8%).¹¹ The proportion of the population who are Hispanic in urban counties (27%) is twice as large as that of suburban counties (14%) and 3 times as large as that of rural counties (8%).¹¹ Nearly half of the American Indian or Alaska Native population lives in rural counties.¹²

Despite racial and ethnic disparities in the PRMR, geographic differences in population distributions, and documented health challenges, little is known nationally about potential

urban-rural differences in pregnancy-related deaths. Previous studies have focused on subnational geographic areas¹³ or were based on death record data^{14,15} with documented challenges,^{16,17} limiting interpretation. Our study used the national Pregnancy Mortality Surveillance System (PMSS) data to examine pregnancy-related deaths in urban and rural counties in the United States. Examining geographic variations in pregnancy-related deaths can inform interventions to prevent future pregnancy-related deaths.

Materials and Methods

Numerator

Here, we analyzed 6 years of PMSS data, from 2011 to 2016. The PMSS methodology to identify pregnancy-related deaths has been published previously.¹⁸ Briefly, clinically-trained epidemiologists reviewed the death certificates of women who died within a year of pregnancy and matched birth certificates or fetal death reports, and other available information (eg, autopsy reports). Deaths were determined to be pregnancy-related if the death occurred during or within 1 year of pregnancy and was caused by a pregnancy complication, a chain of events initiated by pregnancy, or an aggravation of an unrelated condition from the physiological effects of pregnancy. The PMSS receives deidentified records from 52 reporting areas (50 states, New York City, and Washington, DC).

As vital records information is recorded in the PMSS data, reviewers abstracted the geographic location data, including the state, county, or zone improvement plan (ZIP) code of residence, when available, from the death certificate and/or other sources (eg, a matching birth or fetal death certificate). County of residence (not recorded in the PMSS until 2015) is unavailable for records from 2011 to 2014. To address this gap, we developed a geocoding process. Because the residence ZIP code, regardless of source, was the most complete geographic location variable available across pregnancy-related PMSS records during 2011–2016 (93% available), it was first used to assign a 5-digit Federal Information Processing System (FIPS) code¹⁹ to each death by using Centrus Desktop.²⁰ The 5-digit FIPS code is a combination of numeric identifiers for state and county.²¹ Next, the state associated with the resulting FIPS code was compared with the state of residence recorded in the PMSS to ensure the accuracy of the geocoding. If the state identified from the FIPS geocode did not match the state in the PMSS, or if the PMSS record did not have a ZIP code to geocode, then the state and county of residence, if available in the PMSS, were coded to a 5-digit FIPS. PMSS records were excluded from analysis if the geocoded FIPS code did not match and the county was missing in PMSS or if the ZIP code and county were unavailable in the PMSS.

We matched FIPS codes from the geocoded deaths to the 2013 National Center for Health Statistics (NCHS) Urban-Rural Classification Scheme for Counties. This Scheme assigns all U.S. counties to 1 of 6 categories,²² which are defined as:

- Large central metro: metropolitan statistical areas (MSA) with a population of at least 1 million that contain the entire population of the largest principal city within the MSA, or that have their entire population contained in the largest principal city of the MSA, or that contain at least 250,000 inhabitants of any principal city of the MSA.

- Large fringe metro: MSA with a population of at least 1 million that do not fit the definition of large central metropolitan.
- Medium metro: MSA with a population of 250,000–999,999.
- Small metro: MSA with a population of less than 250,000.
- Micropolitan: Urban cluster with a population of 10,000–49,999.
- Noncore: Nonmetropolitan counties that did not qualify as micropolitan (most rural areas).

Maternal race or ethnicity was classified as Hispanic (any race) and non-Hispanic White, Black, American Indian or Alaska Native, and Asian or Pacific Islander. Maternal age was considered in 5-year age groups (15–19, 20–24, 25–29, 30–34, 35–39, 40–44 years).

Denominators

This analysis developed a methodology to estimate county level counts of births to women aged 15 to 44 years by year and race and ethnicity, which can be used to examine urban-rural differences in pregnancy-related deaths. To estimate PRMRs in this analysis, we used 2 methods to obtain birth denominators depending on the stratification variables (county, year, race and ethnicity, or maternal age). Data came from the National Vital Statistics System (NVSS) via the Centers for Disease Control and Prevention, Wide-ranging Online Data for Epidemiologic Research (CDC WONDER).²³

The first method to obtain denominators was for county-level estimates of live births by race and ethnicity and year to account for data suppression (NVSS subnational data with fewer than 10 births were suppressed). We obtained state fertility rates from the NVSS stratified by year, state, and race and ethnicity and multiplied by estimated county population counts of women aged 15 to 44 years from US Census Bureau population data stratified by year, county, and race and ethnicity (Supplemental Online Content; Supplemental Table 1).

We validated our estimate with true births data aggregated by urban-rural category obtained from the NVSS and found close agreement (Supplemental Online Content). As with the numerator data, estimated births were matched to the 2013 NCHS urban-rural classification scheme. However, we cannot use this method to estimate births by county, race and ethnicity, year, and maternal age because nearly 7% of fertility rates stratified by state, race and ethnicity, year, and age were suppressed via CDC WONDER.

A second method was used to obtain the births denominator for urban-rural birth counts stratified by maternal age. For this denominator, we used birth counts from the NVSS aggregated to 2013 NCHS urban-rural classification scheme and maternal age in 5-year age groups (15–19, 20–24, 25–29, 30–34, 35–39, and 40–44 years).

Statistical analysis

PRMRs were calculated for each urban-rural category and stratified by year, race and ethnicity, or maternal age. PRMRs based on fewer than 8 deaths were suppressed as they may be unreliable. Furthermore, 8 deaths represented a relative standard error of 35%. We

used R version 4.0.2 (R Foundation for Statistical Computing, Vienna, Austria) and the *epitools* package²⁴ to calculate PRMRs, obtain 95% confidence intervals (CIs), and perform exact tests of ratio comparisons using the Poisson distribution. We created race and ethnicity and age group PRMR disparity ratios by dividing the PRMR in 1 group by the PRMR in a reference group. Disparity ratios used non-Hispanic White women and women ages 30 to 34 years as the reference groups, respectively, as these were the groups with the highest number of pregnancy-related deaths. This study did not involve human subjects as defined in 45 CFR 46.102(e) and therefore was not reviewed by an institutional review board.

Results

There were 4076 pregnancy-related deaths recorded in the PMSS during 2011–2016. Furthermore, 5-digit FIPS codes were successfully assigned for 3788 records (92.9%) from ZIP code of residence and for 109 records (0.3%) from county of residence. There were 179 records (4%) excluded from this analysis because of missing ZIP code and county information. Of the 3897 records successfully assigned a 5-digit FIPS, 5 deaths were unable to be matched to the NCHS urban-rural classification scheme because of unresolvable data discrepancies, and 12 deaths had an unknown race and ethnicity; these were excluded from analysis. Finally, deaths of women older than 44 years (n=126) and 7 deaths of women younger than 15 years or where age was unknown were excluded because the estimated birth denominator is for women aged 15 to 44 years. The remaining 3747 pregnancy-related deaths were used in this analysis (Table 1 and Supplemental Table 2), representing 92% of all pregnancy-related deaths in the United States that occurred during 2011–2016.

After assigning deaths and births into the 6 NCHS Urban-Rural Classifications, we observed no difference between large central and large fringe metropolitan counties, and we combined these into a single large metro area. More than half of pregnancy-related deaths among Black (61%), Asian or Pacific Islander (71%), and Hispanic (64%) women, and 39% of deaths among White women, occurred in large metro counties. In contrast, approximately 52% of pregnancy-related deaths among American Indian or Alaska Native women occurred in micropolitan or noncore counties. Urban-rural estimated births (Table 2) closely aligned with true birth counts used for validation, by state, NCHS urban-rural category, and year for Hispanic women and non-Hispanic women of all race groups (Supplemental Figures 1 and 2).

The PRMR increased as rurality increased (Figure 1 and Supplemental Table 3, and Supplemental Table 4). The PRMR was higher in micropolitan (19.8; 95% CI, 17.9–22.0) and noncore (23.8; 95% CI, 21.1–26.7) counties than in large metro counties (14.6; 95% CI, 13.9–15.2).

The pattern of increasing PRMR with increasing rurality was observed among most race and ethnicity and age groups; however, age and race- and ethnicity-specific counts were small, resulting in wide overlapping CIs (Table 3). The PRMR among women aged 15 to 24 years was relatively similar by urban-rural category. However, for women in older age groups (ages, 25–29, 30–34, 35–39, and 40–44 years), the disparity between noncore and large metro counties increased with age (Figure 2). Women aged 25 to 29 and 30 to 34 years

living in noncore counties had a PRMR 1.5 times higher than women of the same age groups in large metro counties. Comparing noncore counties with large metro counties, women aged 35 to 39 years and 40 to 44 years had 3.1 and 3.3 times higher PRMR, respectively.

Racial and ethnic and age disparities persisted within each county group (Table 4). Black women, regardless of NCHS Urban-Rural Classification, had a PRMR 3 to 4 times higher than White women in the same area. The PRMR of American Indian or Alaska Native women living in small metro, micropolitan, and noncore counties was 2 to 3 times than that of White women living in these same geographies. The PRMR of non-Hispanic White women was 2 times the PRMR of Hispanic women in noncore counties. In micropolitan and noncore counties, women aged 35 to 39 years had about 3 times PRMR, and women aged 40 to 44 years had 7 to 8 times the PRMR than women aged 30 to 34 years (Table 5).

Comment

Principal Findings

The PRMR increased with increasing rurality. Although large metro counties represented 52% of pregnancy-related deaths, they had the lowest PRMR. Noncore and micropolitan counties represented 17% of pregnancy-related deaths but had the highest PRMR. This difference was observed among all non-Hispanic race groupings and in women aged 30 to 44 years. We found that women aged 25 to 29, 30 to 34, 35 to 39, and 40 to 44 years living in noncore counties had higher PRMR than women of the same age groups in large metro counties, and the urban-rural disparity increased among older age groups, signaling that the “exposure” of living in rural areas has a different effect on PRMR for older women compared with younger women.

Results

These findings add to the limited data about urban-rural differences in pregnancy-related deaths in the United States. Our results were consistent with general findings that mortality is higher in rural areas. However, the PRMR in noncore areas was 1.6 times the PRMR in large metropolitan areas, while all-cause mortality in rural areas was 1.2 times the all-cause mortality in urban areas.⁶ This suggests that there are specific factors present in urban and rural areas that impact pregnancy-related mortality differently than all-cause mortality. An analysis of pregnancy-related deaths in Georgia from 2010 to 2012 found no significant difference in PRMR by rural, urban, or nonrural classifications, or between Black and White women in rural areas.¹³ A national-level study using NVSS data during 1990–1992 and 2005–2009 similarly found no difference in maternal mortality between metropolitan and non-metropolitan areas.¹⁴ We found that the urban-rural PRMR disparity increased with age. Women aged 25 to 29 and 30 to 34 years in noncore counties had a PRMR 1.5 times higher than women in the same age groups in large metro counties; women aged 35 to 39 and 40 to 44 years in noncore counties had a PRMR more than 3 times higher than women of the same age groups in large metro counties. The PRMRs for Black women were significantly higher than that of White women within each urban-rural classification level, which mirror documented national- and state-level disparities.⁴ We found that PRMR was lower for Hispanic women than in non-Hispanic White women in noncore counties. Our analysis

suggested that the PRMR disparity between American Indian or Alaska Native women and White women observed nationally is driven by disparities in small metro, micropolitan, and noncore counties.

Clinical Implications

Differences among geographic location groups might reflect the intersection of age, chronic health conditions, and access to care, including risk-appropriate care. Furthermore, our analysis suggested that increasing rurality affects the PRMR of older women differently compared with the PRMR of younger women. Both the prevalence of delivery hospitalizations of women with substance use disorder and depression,^{25,26} and the prevalence of smoking,²⁷ heart disease, hypertension,²⁸ and obesity,²⁹ were highest among women in rural areas. Infant mortality was highest in rural counties.³⁰ Compared with urban areas, decreased access to obstetrical care among women in rural areas,^{8,10,31} including fewer maternal-fetal medicine subspecialists,³² further driving distances to access perinatal care,¹⁰ and fewer health care providers per capita in rural areas³³ may contribute to disparities among populations by geographic location. Efforts that may help address geographic location disparities include risk-appropriate care,³⁴ transportation reimbursement policies,³⁵ statewide participation in perinatal quality collaboratives,³⁶ and telemedicine or telehealth.^{37,38} Differences among groups by geographic location may reflect the intersection of interpersonal racism or implicit bias, such as health care provider preferential bias toward White patients,³⁹ and structural racism manifested in residential or economic segregation, which are linked to poorer health outcomes.⁴⁰ Indeed, indicators of structural racism such as college graduation, employment, and homeownership inequalities have been shown to differ in their health impacts by urban-rural category.⁴¹ In addition, in rural areas, poverty is highest among Black and American Indian or Alaska Native⁴² populations and is associated with lower odds of identifying a steady source of healthcare;⁴³ furthermore, infant mortality rate is higher among Black and American Indian or Alaska Native populations.⁴⁴ In urban areas, American Indian or Alaska Native individuals have higher odds of cost-related barriers to care and of not having a usual place to go when sick than White individuals.⁴⁵ Across the nation, Black and American Indian or Alaska Native women with at least some college education have higher PRMRs than White women with less than a high school diploma,⁴ which indicates that access to care only partly explains observed urban-rural disparities.⁵ Observed differences between non-Hispanic White women and Hispanic women in noncore counties were consistent with findings sometimes referred to as the “Hispanic paradox.”⁴⁶ However, research also points to differences in cardiovascular mortality rates comparing non-Hispanic White persons individuals with individuals in 3 Hispanic subgroups,⁴⁷ evidence of heterogeneity in outcomes based on acculturation,^{48,49} and severe maternal morbidity being higher among Hispanic women than non-Hispanic White women.^{50,51} Infant mortality data suggested similar associations between mortality and race within Hispanic populations and non-Hispanic populations,⁵² which could be examined in future analyses of pregnancy-related deaths.

Research Implications

Ultimately, pregnancy-related mortality is a multidimensional issue. Exploring differences in causes of death, timing of death relative to pregnancy, and associations between pregnancy-

related deaths and ecologic county-level characteristics, such as access to care, residential segregation, or poverty, may yield additional information about disparities among urban-rural categories.

Strengths and Limitations

This analysis has several limitations. By grouping women by age and race and ethnicity, we masked potential intra-group heterogeneity. Although there are documented urban-rural differences in fertility rates,¹⁹ we assumed that race- and ethnicity-specific fertility rates obtained from the NVSS were the same across both urban and rural counties within a state for a given year. Validation checks (Supplemental Online Content) showed that the estimated births closely agreed with data from the NVSS. Vital records, such as those used during PMSS review, can be flawed by misclassifications, including pregnancy checkbox errors on the death certificate^{16,17} and race and ethnicity coding inaccuracies. PMSS reviewers use additional information whenever possible to improve the accuracy of pregnancy-related classification.¹⁸ Misclassification of American Indian or Alaska Native mortality data was well documented,^{53, 54, 55} particularly among urban areas, which tend to have a higher rate of misclassification than rural areas.⁵⁶ To reduce racial and ethnic misclassifications, PMSS reviewers obtain self-reported data for race and Hispanic origin from matched birth certificates when available. The PRMR does not include injury deaths (eg, caused by drug overdose, suicide, homicide) or cancer-related deaths,⁵⁷ which may skew the results by urban-rural category. Residence ZIP codes in PMSS were abstracted first from the death certificate, but if information were not available, then they were abstracted from other linked sources (eg, the birth or fetal death certificate), which may reduce accuracy if the woman moved since the birth or fetal death certificate was issued. Based on a 10% (n=924) random sample of all PMSS records, 2011–2015, we estimated that 75% of PMSS records contained a ZIP code abstracted from the death certificate.

Conclusions

Noncore and micropolitan counties represented 17% of pregnancy-related deaths and had the highest pregnancy-related mortality ratios. Disparities in pregnancy-related mortality ratios in urban-rural categories indicate that geographic location is an important context for efforts to prevent pregnancy-related deaths. Urban-rural disparities may stem from differences in population health status, access to or quality of care, causes and timing of death, and other socioenvironmental characteristics and exposure to bias and structural racism. A better understanding of these factors' interaction is needed as one piece of a multifactorial response to guide and improve effective policy and program interventions to prevent pregnancy-related deaths.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgement

We thank Sabrina Madni, MPH,¹ for assistance replicating the analysis and ensuring the accuracy of all tables.

Disclosure Statement:

The authors report no conflict of interest or funding source.

References

1. Centers for Disease Control and Prevention. Pregnancy-related deaths. 2019; <https://www.cdc.gov/reproductivehealth/maternalinfanthealth/pregnancy-relatedmortality.htm>. Accessed August 8, 2019.
2. Davis NL, Smoots A, Goodman D. Pregnancy-related deaths: data from 14 U.S. Maternal Mortality Review Committees, 2008–2017. 2019. https://www.cdc.gov/reproductivehealth/maternal-mortality/erase-mm/MMR-Data-Brief_2019-h.pdf. Accessed February 6, 2020.
3. Centers for Disease Control and Prevention. Pregnancy mortality surveillance system. 2019; <https://www.cdc.gov/reproductivehealth/maternalinfanthealth/pregnancy-mortality-surveillance-system.htm>. Accessed August 8, 2019.
4. Petersen EE, Davis NL, Goodman D, et al. Racial/ethnic disparities in pregnancy-related deaths — United States, 2007–2016. *MMWR Morb Mortal Wkly Rep*. 2019;68(35):762–765. [PubMed: 31487273]
5. Kramer MR, Strahan AE, Preslar J, et al. Changing the conversation: applying a health equity framework to maternal mortality reviews. *Am J Obstet Gynecol*. 2019;221(6):609.e601–609.e609.
6. Cosby AG, McDoom-Echebiri MM, James W, Khandekar H, Brown W, Hanna HL. Growth and persistence of place-based mortality in the United States: the rural mortality penalty. *Am J Public Health*. 2019;109(1):155–162. [PubMed: 30496008]
7. Kozhimannil KB, Interrante JD, Henning-Smith C, Admon LK. Rural-urban differences in severe maternal morbidity and mortality in the U.S., 2007–15. *Health Aff*. 2019;38(12):2077–2085.
8. Cohen D, Coco A. Declining trends in the provision of prenatal care visits by family physicians. *The Annals of Family Medicine*. 2009;7(2):128–133. [PubMed: 19273867]
9. Xierali IM, Nivet MA, Rayburn WF. Relocation of obstetrician–gynecologists in the United States, 2005–2015. *Obstet Gynecol*. 2017;129(3):543–550. [PubMed: 28178048]
10. Hung P, Casey MM, Kozhimannil KB, Karaca-Mandic P, Moscovice IS. Rural-urban differences in access to hospital obstetric and neonatal care: how far is the closest one? *J Perinatol*. 2018;38(6):645–652. [PubMed: 29453436]
11. Parker K, Horowitz J, Brown A, Fry R, Cohn D, Igielnik R. What unites and divides urban, suburban and rural communities. 2018:24. <https://www.pewsocialtrends.org/wp-content/uploads/sites/3/2018/05/Pew-Research-Center-Community-Type-Full-Report-FINAL.pdf>. Accessed March 22, 2020.
12. Probst J, Anmal F. Findings brief: social determinants of health among rural American Indian and Alaska Native populations. 2019. https://www.sc.edu/study/colleges_schools/public_health/research/research_centers/sc_rural_health_research_center/documents/socialdeterminantsofhealthamongruralamericanindianandalaskanativepopulations.pdf. Accessed January 5, 2020.
13. Platner M, Loucks TL, Lindsay MK, Ellis JE. Pregnancy-associated deaths in rural, nonrural, and metropolitan areas of Georgia. *Obstet Gynecol*. 2016;128(1):113–120. [PubMed: 27275792]
14. Singh GK, Siahpush M. Widening rural-urban disparities in all-cause mortality and mortality from major causes of death in the USA, 1969–2009. *J Urban Health*. 2014;91(2):272–292. [PubMed: 24366854]
15. Hansen A, Moloney M. Pregnancy-related mortality and severe maternal morbidity in rural Appalachia: established risks and the need to know more. *J Rural Health*. 2020;36(1):3–8. [PubMed: 31246338]
16. Baeva S, Saxton DL, Ruggiero K, et al. Identifying Maternal Deaths in Texas Using an Enhanced Method, 2012. *Obstet Gynecol*. 2018;131(5):762–769. [PubMed: 29630012]

¹National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention, Atlanta, Georgia

17. Catalano A, Davis NL, Petersen EE, et al. Pregnant? Validity of the pregnancy checkbox on death certificates in four states, and characteristics associated with pregnancy checkbox errors. *Am J Obstet Gynecol.* 2020;222(3):269 e261–269 e268.
18. Creanga AA, Syverson C, Seed K, Callaghan WM. Pregnancy-related mortality in the United States, 2011–2013. *Obstet Gynecol.* 2017;130(2):366–373. [PubMed: 28697109]
19. Daniels K, Martinez G, Nugent C. Urban and rural variation in fertility-related behavior among U.S. women, 2011–2015. *NCHS Data Brief.* 2018;297:1–8. <https://www.cdc.gov/nchs/data/databriefs/db297.pdf>. Accessed May 21, 2020.
20. Pitney Bowes Inc. Centrus Desktop. Version 6.0. Troy, NY2018. <https://www.pitneybowes.com/content/dam/support/software/product-documentation/public/geostan/v2016-04/en-us/centrus-desktop-v6-04-00-n-quickstart-guide.pdf>. Accessed September 17, 2019
21. United States Census. Understanding geographic identifiers (GEOIDs). 2020. <https://www.census.gov/programs-surveys/geography/guidance/geo-identifiers.html>. Accessed March 6, 2020.
22. Ingram D, Franco S. 2013 NCHS urban–rural classification scheme for counties. National Center for Health Statistics. 2014; *Vital Health Stat* 2(166). https://www.cdc.gov/nchs/data/series/sr_02/sr02_166.pdf. Accessed February 6, 2020.
23. Centers for Disease Control and Prevention. About natality, 2007–2019. <http://wonder.cdc.gov/natality-current.html>. Accessed March 6, 2020.
24. Aragon TJ. epitools: Epidemiology Tools. R package version 0.5-10.1.2020. <https://cran.r-project.org/package=epitools>. Accessed March 16, 2021
25. Admon LK, Winkelman TNA, Moniz MH, Davis MM, Heisler M, Dalton VK. Disparities in chronic conditions among women hospitalized for delivery in the United States, 2005–2014. *Obstet Gynecol.* 2017;130(6):1319–1326. [PubMed: 29112666]
26. Soni A, Fingar K, L R. Obstetric delivery inpatient stays involving substance use disorders and related clinical outcomes, 2016. <https://www.hcup-us.ahrq.gov/reports/statbriefs/sb254-Delivery-Hospitalizations-Substance-Use-Clinical-Outcomes-2016.jsp>. Accessed October 2019.
27. Nighbor TD, Doogan NJ, Roberts ME, et al. Smoking prevalence and trends among a U.S. national sample of women of reproductive age in rural versus urban settings. *PLoS One.* 2018;13(11):e0207818. [PubMed: 30485376]
28. Shaw KM, Theis KA, Self-Brown S, Roblin DW, Barker L. Chronic disease disparities by county economic status and metropolitan classification, Behavioral Risk Factor Surveillance System, 2013. *Prev Chronic Dis.* 2016;13:E119. [PubMed: 27584875]
29. Lundeen EA, Park S, Pan L, O’Toole T, Matthews K, Blanck HM. Obesity prevalence among adults living in metropolitan and nonmetropolitan counties — United States, 2016. *MMWR Morb Mortal Wkly Rep.* 2018;67(23):653–658. [PubMed: 29902166]
30. Ely D, Hoyert D. Differences between rural and urban areas in mortality rates for the leading causes of infant death: United States, 2013–2015. *NCHS Data Brief.* 2018;300:1–8. <https://www.cdc.gov/nchs/data/databriefs/db300.pdf> Accessed March 25, 2020.
31. Kozhimannil KB, Hung P, Henning-Smith C, Casey MM, Prasad S. Association between loss of hospital-based obstetric services and birth outcomes in rural counties in the United States. *J Am Med Assoc.* 2018;319(12):1239–1247.
32. Rayburn WF, Klagholz JC, Elwell EC, Strunk AL. Maternal-fetal medicine workforce in the United States. *Am J Perinatol.* 2012;29(9):741–746. [PubMed: 22773289]
33. Agency for Healthcare Research and Quality. National healthcare quality and disparities report chartbook on rural health care. <https://www.ahrq.gov/sites/default/files/wysiwyg/research/findings/nhqdr/chartbooks/qdr-ruralhealthchartbook-update.pdf>. Accessed October 2017. AHRQ Pub. No. 17(18)-0001–2-EF.
34. Catalano A, Bennett A, Busacker A, et al. Implementing CDC’s Level of Care Assessment Tool (LOCATe): A National Collaboration to Improve Maternal and Child Health. *J Womens Health (Larchmt).* 2017;26(12):1265–1269. [PubMed: 29240547]

35. Okoroh EM, Kroelinger CD, Lasswell SM, Goodman DA, Williams AM, Barfield WD. United States and territory policies supporting maternal and neonatal transfer: review of transport and reimbursement. *J Perinatol.* 2016;36(1):30–34. [PubMed: 26334399]
36. Gupta M, Donovan EF, Henderson Z. State-based perinatal quality collaboratives: Pursuing improvements in perinatal health outcomes for all mothers and newborns. *Semin Perinatol.* 2017;41(3):195–203. [PubMed: 28646987]
37. Mann S, McKay K, Brown H. The maternal health compact. *N Engl J Med.* 2017;376(14):1304–1305. [PubMed: 28379803]
38. Marcin JP, Shaikh U, Steinhorn RH. Addressing health disparities in rural communities using telehealth. *Pediatr Res.* 2016;79(1–2):169–176. [PubMed: 26466080]
39. Hall WJ, Chapman MV, Lee KM, et al. Implicit racial/ethnic bias among health care professionals and its influence on health care outcomes: a systematic review. *Am J Public Health.* 2015;105(12):e60–76.
40. Williams DR, Collins C. Racial residential segregation: a fundamental cause of racial disparities in health. *Public Health Rep.* 2001(116):404–416. [PubMed: 12042604]
41. Bell CN, Owens-Young JL. Self-rated health and structural racism indicated by county-level racial inequalities in socioeconomic status: the role of urban-rural classification. *J Urban Health.* 2020;97(1):52–61. [PubMed: 31898201]
42. U.S. Department of Agriculture. Economic Research Service. Rural poverty & well-being. 2018. <https://www.ers.usda.gov/topics/rural-economy-population/rural-poverty-well-being/#demographics>. Accessed May 1, 2020.
43. Caldwell JT, Ford CL, Wallace SP, Wang MC, Takahashi LM. Racial and ethnic residential segregation and access to health care in rural areas. *Health Place.* 2017;43:104–112. [PubMed: 28012312]
44. Singh GK, Daus GP, Allender M, et al. Social determinants of health in the United States: addressing major health inequality trends for the nation, 1935–2016. *Int J MCH AIDS.* 2017;6(2):139–164. [PubMed: 29367890]
45. Rutman S, Phillips L, Sparck A. Health care access and use by urban American Indians and Alaska Natives: findings from the National Health Interview Survey (2006–09). *J Health Care Poor Underserved.* 2016;27(3):1521–1536. [PubMed: 27524782]
46. Chen Y, Freedman ND, Rodriguez EJ, et al. Trends in Premature Deaths Among Adults in the United States and Latin America. *JAMA Network Open.* 2020;3(2).
47. Rodriguez F, Hastings KG, Boothroyd DB, et al. Disaggregation of Cause-Specific Cardiovascular Disease Mortality Among Hispanic Subgroups. *JAMA Cardiology.* 2017;2(3).
48. Flores ME, Simonsen SE, Manuck TA, Dyer JM, Turok DK. The “Latina epidemiologic paradox”: contrasting patterns of adverse birth outcomes in U.S.-born and foreign-born Latinas. *Womens Health Issues.* 2012;22(5):e501–507. [PubMed: 22944904]
49. Hoggatt KJ, Flores M, Solorio R, Wilhelm M, Ritz B. The “Latina epidemiologic paradox” revisited: the role of birthplace and acculturation in predicting infant low birth weight for Latinas in Los Angeles, CA. *J Immigr Minor Health.* 2012;14(5):875–884. [PubMed: 22160842]
50. Howell EA, Egorova NN, Janevic T, et al. Race and Ethnicity, Medical Insurance, and Within-Hospital Severe Maternal Morbidity Disparities. *Obstet Gynecol.* 2020;135(2):285–293. [PubMed: 31923076]
51. Mujahid MS, Kan P, Leonard SA, et al. Birth hospital and racial and ethnic differences in severe maternal morbidity in the state of California. *American Journal of Obstetrics and Gynecology.* 2020.
52. Rice WS, Goldfarb SS, Brisendine AE, Burrows S, Wingate MS. Disparities in Infant Mortality by Race Among Hispanic and Non-Hispanic Infants. *Maternal and Child Health Journal.* 2017;21(7):1581–1588. [PubMed: 28197819]
53. Anderson RN, Copeland G, Hayes JM. Linkages to improve mortality data for American Indians and Alaska Natives: a new model for death reporting? *Am J Public Health.* 2014;104 Suppl 3:S258–262. [PubMed: 24754614]
54. Arias E, Heron M, Hakes J. The validity of race and Hispanic-origin reporting on death certificates in the United States: An update. National Center for Health Statistics. *Vital Health Stat.* 2(172).

55. Joshi S, Weiser T, Warren-Mears V. Drug, opioid-involved, and heroin-involved overdose deaths among American Indians and Alaska Natives — Washington, 1999–2015. *MMWR Morb Mortal Wkly Rep.* 67:1384–1384.
56. Jim MA, Arias E, Seneca DS, et al. Racial misclassification of American Indians and Alaska Natives by Indian Health Service Contract Health Service Delivery Area. *Am J Public Health.* 2014;104 Suppl 3:S295–302. [PubMed: 24754617]
57. Petersen EE, Davis NL, Goodman D, et al. Vital Signs: Pregnancy-Related Deaths, United States, 2011–2015, and Strategies for Prevention, 13 States, 2013–2017. *MMWR Morb Mortal Wkly Rep.* 2019;68(18):423–429. [PubMed: 31071074]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

AJOG at a Glance:**Why was this study conducted?**

This study aimed to compare pregnancy-related mortality across and within urban and rural counties in the United States by race and ethnicity and age.

Key findings

During 2011–2016, rural counties had the highest pregnancy-related mortality ratio, whereas urban counties had the lowest. Urban-rural disparities in pregnancy-related mortality ratio increased with age. Racial and ethnic disparities mirrored national disparities.

What does this add to what is known?

This study reported on urban-rural disparities in pregnancy-related deaths and used the most recent national data on pregnancy-related deaths from the Pregnancy Mortality Surveillance System. Other studies were subnational, based solely on death record data with documented challenges or hospital discharge data. Finally, the urban-rural disparity was greater among older women than among younger women.




FIGURE 1.

Pregnancy-related mortality ratios by urban-rural category

The PRMR is the number of pregnancy-related deaths per 100,000 live births. The PRMR is highest in rural areas. Rural areas include micropolitan (urban cluster with a population of 10,000–49,999) and noncore (nonmetropolitan counties that did not qualify as micropolitan; most rural areas). Urban areas include large metro (“MSA,” with a population of at least 1 million), medium metro (MSA with a population of 250,000–999,999), and small metro (MSA with a population of less than 250,000).

MSA, metropolitan statistical area; PRMR, pregnancy-related mortality ratio.

Merkt et al. Urban-rural differences in pregnancy-related deaths. *Am J Obstet Gynecol* 2021.




FIGURE 2.

Pregnancy-related mortality ratios by age group and urban-rural category

The PRMR is the number of pregnancy-related deaths per 100,000 live births. The PRMR increased in rural areas and among older age groups. Rural areas include micropolitan (urban cluster with a population of 10,000–49,999) and noncore (nonmetropolitan counties that did not qualify as micropolitan; most rural areas). Urban areas include large metro (“MSA,” with a population of at least 1 million), medium metro (MSA with a population of 250,000–999,999), and small metro (MSA with a population of less than 250,000). MSA, metropolitan statistical area; PRMR, pregnancy-related mortality ratio.

Merket et al. Urban-rural differences in pregnancy-related deaths. *Am J Obstet Gynecol* 2021.

Table 1. Distribution of pregnancy-related deaths by county urban-rural category, United States, 2011–2016

Characteristics	n (%) ^a				
	Total	Large metro	Medium metro	Small metro	Metropolitan Noncore
Overall	3,747	1,956 (52)	797 (21)	344 (9)	281 (7)
Race and ethnicity ^b					
White	1,524	589 (39)	365 (24)	177 (12)	218 (14)
Black	1,384	849 (61)	262 (19)	111 (8)	92 (7)
American Indian or Alaska Native	65	10 (15)	9 (14)	12 (18)	15 (23)
Asian or Pacific Islander	204	144 (71)	36 (18)	9 (4)	9 (4)
Hispanic	570	364 (64)	125 (22)	35 (6)	35 (6)
Age group (y)					
15–19	175	69 (39)	38 (22)	27 (15)	24 (14)
20–24	614	283 (46)	147 (24)	69 (11)	70 (11)
25–29	863	432 (50)	205 (24)	83 (10)	73 (8)
30–34	904	513 (57)	186 (21)	80 (9)	74 (8)
35–39	808	446 (55)	154 (19)	52 (6)	89 (11)
40–44	383	213 (56)	67 (17)	33 (9)	39 (10)

The definitions are as follows: large metro, an MSA with a population of at least 1 million; medium metro, an MSA with a population of 250,000 to 999,999; small metro, an MSA with a population of less than 250,000; micropolitan, an urban cluster with a population of 10,000 to 49,999; and noncore, nonmetropolitan counties that did not qualify as micropolitan (most rural areas).

MSA, metropolitan statistical area.

^aPercentages may not add to 100 because of rounding;

^bWomen identified as White, Black, American Indian or Alaska Native, or Asian or Pacific Islander were not Hispanic. Hispanic women could be of any race.

Merkt et al. Urban-rural differences in pregnancy-related deaths. Am J Obstet Gynecol 2021.

Table 2. Distribution of live births by county urban-rural category, United States, 2011–2016

Characteristics	n (%) ^a				
	Total	Large metro	Medium metro	Small metro	Noncore
Overall	23,500,220	13,427,170 (57)	4,905,002 (21)	2,125,202 (9)	1,860,624 (8)
Race and ethnicity ^b					
White	12,738,577	6,198,217 (49)	2,836,817 (22)	1,461,336 (11)	1,352,185 (11)
Black	3,500,990	2,313,354 (66)	650,231 (19)	244,927 (7)	174,346 (5)
American Indian or Alaska Native	225,937	53,941 (24)	44,277 (20)	30,643 (14)	46,463 (21)
Asian or Pacific Islander	1,562,250	1,198,684 (77)	242,727 (16)	70,743 (5)	39,222 (3)
Hispanic	5,472,466	3,662,974 (67)	1,130,950 (21)	317,553 (6)	248,408 (5)
Age group (y)					
15–19	1,587,734	747,334 (47)	368,323 (23)	168,658 (11)	179,540 (11)
20–24	5,244,997	2,506,440 (48)	1,207,345 (23)	569,330 (11)	568,859 (11)
25–29	6,773,408	3,628,200 (54)	1,491,353 (22)	665,686 (10)	594,682 (9)
30–34	6,269,593	3,884,903 (62)	1,239,884 (20)	492,814 (8)	401,491 (6)
35–39	2,971,425	1,981,441 (67)	541,435 (18)	193,815 (7)	158,398 (5)
40–44	653,142	449,665 (69)	113,272 (17)	38,159 (6)	31,999 (5)

Births by race and ethnicity groups were estimated by multiplying county-level number of women and stratified by race and ethnicity and year and state fertility rates. Births by age group were obtained from the National Vital Statistics System via CDC WONDER and do not sum to the overall births. The definitions are as follows: large metro, an MSA with a population of at least 1 million; medium metro, an MSA with a population of 250,000 to 999,999; small metro, an MSA with a population of less than 250,000; micropolitan, urban cluster with a population of 10,000 to 49,999; and noncore, nonmetropolitan counties that did not qualify as micropolitan (most rural areas).

CDC WONDER, Centers for Disease Control and Prevention, Wide-ranging Online Data for Epidemiologic Research; MSA, metropolitan statistical area.

^a Percentages may not add to 100 because of rounding;

^b Race and ethnicity groups were defined as Hispanic births (regardless of race) and non-Hispanic White, Black, American Indian or Alaska Native, or Asian or Pacific Islander births.

MerkT et al. Urban-rural differences in pregnancy-related deaths. Am J Obstet Gynecol 2021.

Table 3.

Pregnancy-related mortality ratio by county urban-rural category and sociodemographic characteristics, United States, 2011–2016

Characteristics	PRMR ^a (95% Confidence Interval)					
	Total	Large metro	Medium metro	Small metro	Metropolitan	Noncore
Overall	15.9 (15.4–16.5)	14.6 (13.9–15.2)	16.2 (15.1–17.4)	16.2 (14.5–18.0)	19.8 (17.9–22.0)	23.8 (21.1–26.7)
Race and ethnicity ^b						
White	12.0 (11.4–12.6)	9.5 (8.8–10.3)	12.9 (11.6–14.3)	12.1 (10.4–14.0)	16.1 (14.1–18.4)	19.7 (16.9–22.8)
Black	39.5 (37.5–41.7)	36.7 (34.3–39.3)	40.3 (35.6–45.5)	45.3 (37.3–54.6)	52.8 (42.5–64.7)	59.3 (46.2–74.9)
American Indian or Alaska Native	28.8 (22.2–36.7)	18.5 (8.9–34.1)	20.3 (9.3–38.6)	39.2 (20.2–68.4)	32.3 (18.1–53.2)	37.5 (22.6–58.6)
Asian or Pacific Islander	13.1 (11.3–15.0)	12.0 (10.1–14.1)	14.8 (10.4–20.5)	12.7 (5.8–24.2)	22.9 (10.5–43.6)	- ^c
Hispanic	10.4 (9.6–11.3)	9.9 (8.9–11.0)	11.1 (9.2–13.2)	11.0 (7.7–15.3)	14.1 (9.8–19.6)	9.8 (4.9–17.5)
Age group (y)						
15–19	11.0 (9.4–12.8)	9.2 (7.2–11.7)	10.3 (7.3–14.2)	16.0 (10.5–23.3)	13.4 (8.6–19.9)	13.7 (8.0–22.0)
20–24	11.7 (10.8–12.7)	11.3 (10.0–12.7)	12.2 (10.3–14.3)	12.1 (9.4–15.3)	12.3 (9.6–15.5)	11.4 (8.4–15.3)
25–29	12.7 (11.9–13.6)	11.9 (10.8–13.1)	13.7 (11.9–15.8)	12.5 (9.9–15.5)	12.3 (9.6–15.4)	17.8 (13.9–22.5)
30–34	14.4 (13.5–15.4)	13.2 (12.1–14.4)	15.0 (12.9–17.3)	16.2 (12.9–20.2)	18.4 (14.5–23.1)	20.4 (15.2–26.8)
35–39	27.2 (25.3–29.1)	22.5 (20.5–24.7)	28.4 (24.1–33.3)	26.8 (20.0–35.2)	56.2 (45.1–69.1)	69.5 (53.9–88.3)
40–44	58.6 (52.9–64.8)	47.4 (41.2–54.2)	59.1 (45.8–75.1)	86.5 (59.5–121.5)	121.9 (86.7–166.6)	154.6 (105.1–219.5)

The definitions are as follows: large metro, an MSA with a population of at least 1 million; medium metro, an MSA with a population of 250,000 to 999,999; small metro, an MSA with a population of less than 250,000; metropolitan, urban cluster with a population of 10,000 to 49,999; and noncore, nonmetropolitan counties that did not qualify as metropolitan (most rural areas).

CDC WONDER, Centers for Disease Control and Prevention, Wide-ranging Online Data for Epidemiologic Research; MSA, metropolitan statistical area; PRMR, pregnancy-related mortality ratio.

^aPRMR denominators (births) for race and ethnicity groups were estimated by multiplying county-level number of women, stratified by race and ethnicity and year and state fertility rates. PRMR denominators (births) for age groups were obtained from the National Vital Statistics System (NVSS) via CDC WONDER;

^bWomen identified as White, Black, American Indian or Alaska Native, or Asian or Pacific Islander were not Hispanic. Hispanic women could be of any race;

^cEm dash indicates a PRMR based on fewer than 8 deaths; these results were suppressed because ratios might be unreliable.

Merkt et al. Urban-rural differences in pregnancy-related deaths. Am J Obstet Gynecol 2021.

Disparity ratios within urban-rural category by race and ethnicity, United States, 2011–2016

Table 4.

Urban-rural category	Black-to-White disparity ratio	AI/AN-to-White disparity ratio	A/PI-to-White disparity ratio	Hispanic-to-White disparity ratio
Large metro	3.9 ^a	1.9	1.3 ^a	1.0
Medium metro	3.1 ^a	1.6	1.1	0.9
Small metro	3.7 ^a	3.2 ^a	1.0	0.9
Metropolitan	3.3 ^a	2.0 ^a	1.4	0.9
Noncore	3.0 ^a	1.9 ^a	– ^b	0.5 ^a

Disparity ratios were calculated by dividing the PRMR in 1 race and ethnicity group by the PRMR in a reference group. Non-Hispanic White women were the reference group, as they represented the largest racial and ethnic group. The definition are as follows: large metro, an MSA with a population of at least 1 million; medium metro, an MSA with a population of 250,000 to 999,999; small metro, an MSA with a population of less than 250,000; micropolitan, urban cluster with a population of 10,000 to 49,999; and noncore, nonmetropolitan counties that did not qualify as micropolitan (most rural areas). Women identified as White, Black, American Indian or Alaska Native, or Asian or Pacific Islander were not Hispanic. Hispanic women could be of any race.

MSA, metropolitan statistical area; PRMR, pregnancy-related mortality ratio.

^aDenotes statistical significance, P<.05;

^bEm dash indicates a PRMR based on fewer than 8 deaths; these results were suppressed because ratios might be unreliable.

Merkt et al. Urban-rural differences in pregnancy-related deaths. Am J Obstet Gynecol 2021.

Disparity ratios within urban-rural category by age group, United States, 2011–2016

Table 5.

Age group (y)					
Urban-rural category	15–19 to 30–34 disparity ratio	20–24 to 30–34 disparity ratio	25–29 to 30–34 disparity ratio	35–39 to 30–34 disparity ratio	40–44 to 30–34 disparity ratio
Large metro	0.7 ^a	0.9	0.9	1.7 ^a	3.6 ^a
Medium metro	0.7	0.8 ^a	0.9	1.9 ^a	3.9 ^a
Small metro	1.0	0.7	0.8	1.7 ^a	5.3 ^a
Metropolitan	0.7	0.7 ^a	0.7 ^a	3.1 ^a	6.6 ^a
Noncore	0.7	0.6 ^a	0.9	3.4 ^a	7.6 ^a

Disparity ratios were calculated by dividing the PRMR in 1 age group by the PRMR in a reference group. Women aged 30 to 34 years were the reference group, as they represented the largest age group. The definitions are as follows: large metro, an MSA with a population of at least 1 million; medium metro, an MSA with a population of 250,000 to 999,999; small metro, an MSA with a population of less than 250,000; metropolitan, urban cluster with a population of 10,000 to 49,999; and noncore, nonmetropolitan counties that did not qualify as metropolitan (most rural areas).

MSA, metropolitan statistical area; PRMR, pregnancy-related mortality ratio.

^aDenotes statistical significance, P<.05.

MerkT et al. Urban-rural differences in pregnancy-related deaths. Am J Obstet Gynecol 2021.