

## EVALUATION OF PULMONARY REACTIONS IN HARD METAL WORKERS

G. CHIAPPINO • M. Tomasini • G. Rivolta • A. Forni  
• E. Sabbioni\* • R. Pietra\*

Research Center on the Biological Effects of Inhaled Dusts  
Institute of Occupational Health, University of Milan, Italy, and  
\*Radioc hemistry Division, Commission of the European Communities  
Joint Research Center, Ispra (VA), Italy

### INTRODUCTION

The production of hard metal components of equipments for different technological uses is presently actively expanding in several countries.<sup>9,10</sup> Consequently, the number of workers exposed to risk of inhaling hard metal particles is increasing.

Lung disorders from exposure to hard metals have been known since almost 50 years, and recently cobalt has been recognized as the major responsible agent.<sup>3,8,9</sup> Both asthma and more severe deep lung involvement can occur, with possible evolution in fibrosis.<sup>2,8,9</sup>

In the last two years we had the opportunity to thoroughly investigate two groups of workers exposed to inhalation of hard metals in completely different occupational settings and to evaluate their pulmonary reactions by bronchoalveolar lavage (BAL). The main findings are reported in this paper.

### MATERIALS AND METHODS

#### Subjects

Group A consisted of 26 workers, 25 males and one female, 18 smokers and 8 non-smokers, aged from 21 to 64 years (mean 32.9 years), engaged in the production of hard metal tools for 2 to 49 years (mean 10.4 years), and therefore exposed to tungsten carbide, titanium, tantalum and cobalt.

Group B consisted of 5 male workers, aged from 21 to 26 years, all but one non-smokers, engaged only in grinding hard metal tool edges by means of high-speed diamond-cobalt wheels with oily coolants for 1 to 9 years (mean 5 years) in a small workshop, with no proper protective equipments.

Two workers of group A, one non-smoker and one smoker, both 30 years of age, exposed for 7 and 13 years respectively, complained of work-related asthma, but were in remission at the time of study. Two workers of group B, non-smokers, aged 21 and 26, suffered from severe exertional dyspnea and complained of weight loss; they had been exposed for 6 and 5 years respectively, and were no longer exposed since 6 months when first studied.

#### Methods

The 31 workers were hospitalized and were investigated by the following protocol: personal and occupational history; routine clinical and laboratory checkup; chest X-rays; lung

function tests; patch tests for cobalt and tungsten, and skin tests for common inhalant allergens; bronchoalveolar lavage (BAL); elemental determination by neutron activation analysis in BAL, blood, urine, toenails and pubic hair.

BAL was performed according to the method currently used in our institute, and cytologically examined as described elsewhere.<sup>4</sup> Lymphocyte typing in BAL was performed on cytocentrifuge preparations by monoclonal antibodies in an immunoperoxidase assay, as described in detail elsewhere.<sup>5</sup>

Elemental determination in biological specimens was carried out by neutron activation analysis, as described elsewhere.<sup>7</sup>

### RESULTS AND DISCUSSIONS

In group A, no significant alterations were observed at chest X-rays. Lung function tests showed no significant impairment of the ventilatory function, whereas one subject with asthma and 4 additional workers presented a slight impairment of the pulmonary diffusion of gas.

On the contrary, BAL cytology showed a marked increase of total cells with a high percentage of lymphocytes in the non-smoker asthmatic subject (Table I). Increased percentage of lymphocytes (17%) with total cells somewhat above the highest normal value for smokers was detected also in one of the asymptomatic workers.

A moderate relative or absolute increase of lymphocytes was present in 8 additional workers, one with asthma (Table II). The increase of BAL lymphocytes was mainly due to an increase of suppressor (OKT8+) cells, with consequent inversion or marked reduction of the OKT4/OKT8 ratio, independent from the smoking habits.

Among the 5 young workers of group B, exposed to the fine aerosol spray originating from high-speed wet-grinding of hard metal edges, the two with severe dyspnea showed quite similar clinical features, consisting in diffuse irregular opacities at chest X-rays, associated with a significant restrictive impairment of the ventilatory function.

The cytology of BAL (Table III) showed in both cases an alveolitis of high intensity, characterized by an enormous increase of total cells, with numerous bizarre giant cells, marked eosinophilia, increased absolute number of lym-

**Table I**  
**Bronchoalveolar Lavage Cytology in the 26 Hard Metal Workers of**  
**Group A (mean values and ranges)**

Cases	No.	Total cells /ml x 10 <sup>-3</sup>	Macro- phages	Lympho- cytes	% Neutro- phils	Eosino- phils
Asthma	1 NS	1800	61	36	0	3
	1 S	640	91	7	2	0
No symptoms	7 NS	173 (95-235)	76 (51-91)	15 (7-27)	3 (<1-6)	<1 (0-2)
	17 S	700 (235-1530)	91 (80-99)	6 (<1-17)	3 (<1-5)	<1 (0-1)

NS, non-smoker

S, smoker

**Table II**  
**Bronchoalveolar Lavage Lymphocyte Behavior in Hard Metal Workers of Group A**

No. of cases		BAL lymphocyte number	Ratio OKT4/OKT8
2	1 asthma	markedly increased	0.15
	1 no symptoms	increased	0.2
8	1 asthma		0.35
	7 no symptoms	slightly increased	0.2-0.8
16	no symptoms	normal	0.9 (1 case)

Table III  
Bronchoalveolar Lavage Cytology in the 5 Hard Metal Workers of  
Group B (individual values)

Cases	No.	Total cells /ml x 10 <sup>-3</sup>	Macro- phages	Lympho- cytes	%	Neutro- phils	Eosino- phils
Interstitial lung disease	NS	3000	78	3	1	18	
	NS*	2475	62	32**	3	3**	
No symptoms	2 NS	120-210	68-72	23-27	4-5	0-1	
	1 S	375	87	10	2	<1	

NS, non smoker      S, smoker

\* under steroid treatment since two months

\*\* BAL pre-treatment at another institution: Lymphocytes 5%, Eosinophils 60%

phocytes with inverted OKT4/OKT8 ratio (0.52 and 0.58 respectively). The lymphocyte subpopulations in blood were normal.

The BAL findings correlated strictly with the histological features observed at open lung biopsy, performed in one case. In fact, histology showed alveolar filling by large mononuclear cells with occasional giant cells, involvement of the alveolar wall (edema, infiltration by lymphocytes and eosinophils) and hyperplastic alveolar lining, and was therefore typical for a desquamative interstitial pneumonitis.

Patch tests for cobalt were positive only in one case.

The 3 asymptomatic workers of group B showed only a slight increase of lymphocytes at BAL, with decreased OKT4/OKT8 ratios (Table III).

As far as concerns the concentrations of cobalt, tungsten and tantalum in biological specimens, evaluated by neutron activation analysis and presently available only for 11 subjects (including the two with asthma of group A and one of group B with the severe subacute lung disorder), no relation was observed between the presence of disease and the amount of any of the metals. The concentrations of cobalt and tungsten were in most instances one to three orders of magnitude compared to normal mean values (data not shown).

The highest values of all three means were observed, independently from signs and symptoms, in the workers with exposure of longer duration, and this was particularly true for concentrations at sites of deposit such as hair and nails. The behavior of the metals in the various specimens was constantly different and indicative of a relatively high mobility of inhaled cobalt. In fact, cobalt concentrations were the

lowest in BAL and the highest in nails, hair and blood, whereas tungsten levels were the highest in BAL and the lowest in blood (median values for cobalt in BAL 4.6 ng/ml, in nails 8000 ng/g, in hair 5550 ng/g and in blood 6.2 ng/ml; median values for tungsten in BAL 348 ng/ml, in nails 5540 ng/g, in hair 875 ng/g and in blood 0.3 ng/ml).

Our findings support the hypothesis that immunological mechanisms play a fundamental role in the pathogenesis of respiratory disorders from exposure to hard metals. The occurrence of asthmatic or interstitial alterations, in fact, seems based on individual hypersusceptibility and not on the degree of occupational exposure.

The relatively low prevalence of respiratory symptoms in group A (2 among 26 workers with severe and often prolonged exposure) also points to individual factors in the pathogenesis of overt disease, and is in agreement with data from the literature.<sup>8</sup>

A lymphocytic alveolitis at BAL with reduced OKT4/OKT8 ratio has been occasionally reported in subjects with hard metal disease or cobalt lung.<sup>1,3,6</sup> However, our finding of a tendency to an increase of lymphocytes of deep lung, with an imbalance of T-lymphocyte subpopulations, in several asymptomatic workers does not seem to have been reported so far, and suggests a more frequent effect of the exposure on the local immune system, whose pathogenetic significance cannot at present be stated.

In group B, the lung pathological changes were more severe and more frequent (2 out of 5 workers). These observations are in agreement with data of the literature, which indicate a higher risk in operations on hard metals involving the use of coolant fluids and producing fine aerosol sprays.<sup>9</sup> In this

group too, however, individual hypersusceptibility seems important, but the type of lung changes suggests different pathogenetic mechanisms. In fact, in agreement with the observations of Davison et al.,<sup>2</sup> the histological and cytological features are not those typical for hypersensitivity pneumonitis.

Even though immunological hypersusceptibility seems to be the most probable pathogenetic factor, the specific immune reactions directly involved in the induction of the different disorders are still poorly understood. Therefore, it is impossible at present to propose sensitive and specific tests for the early identification of hypersusceptible individuals.


In the medical surveillance of exposed workers, the aim should be to detect early respiratory alterations. Lung function tests and BAL in selected cases should be considered at the present time the methods of choice.

## REFERENCES

1. Cassina, G., Migliori, M., Michetti, G., Argenti, G., Seghizzi, P.: Un Caso di Interstiziopatia da Cobalto: Considerazioni Patogenetiche e Prognostiche. *Med. Lav.* 78:224-234 (1984).
2. Davison, A.G., Haslam, P.L., Corrin, B., Coutts, I.I., Dewar, A., Riding, W.D., Studdy, P.R., Newman-Taylor, A.J.: Interstitial Lung Disease and Asthma in Hard-Metal Workers: Bronchoalveolar Lavage, Ultrastructural, and Analytical Findings and Results of Bronchial Provocation Tests. *Thorax* 38:119-128 (1983).
3. Demedts, M., Gheysens, B., Nagels, J., Verbeken, E., Laweryns, J., van den Eeckhout, A., Lahaye, D., Gyselen, A.: Cobalt Lung in Diamond Polishers. *Am. Rev. Respir. Dis.* 130:130-135 (1984).
4. Forni, A., Guerreri, M.C., Chiappino, G.: Nuovi Metodi di Indagine nelle Pneumopatie Professionali: il Lavaggio Broncoalveolare. *Med. Lav.* 76:11-16, (1985).
5. Forni, A., Ortisi, E., Rivolta, G., Chiappino, G.: Bronchoalveolar Lavage T-Lymphocyte Subpopulations in Occupational and Non-occupational Lung Diseases. In: *New Frontiers in Cytology. Modern Aspects of Research and Practice*. K. Goertler, G. Feichter and S. Witte, Eds. Springer, Heidelberg, New York, Tokyo (1988) (in press).
6. Gaucher, L., Grunbec, N., Haloun, A., Ordronneau, J.: Lavage Alvéolaire et Fibrose Pulmonaire aux Métaux Durs. *Presse Med.* 15:440 (1986).
7. Nicolaou, G., Pietra, R., Sabbioni, E., Mosconi, G., Cassina, G., Seghizzi, P.: Multielemental Determination of Metals in Biological Specimens of Hard Metal Workers: A study carried out by Neutron Activation Analysis. *J. Trace Elem. Electrolytes Health Dis.* 1:73-77 (1987).
8. Parkes, W.R.: *Occupational Lung Disorders*, 2nd Ed., pp 464-466. Butterworth, London (1982).
9. Sjögren, J., Hillerdal, G., Andersson, A., Zetterström, O.: Hard Metal Lung Disease: Importance of Cobalt in Coolants. *Thorax* 35:653-659 (1980).
10. Sprince, N.L., Chamberlin, R.I., Hales, C.A., Weber, A.L., Kazemz, H.: Respiratory Disease in Tungsten Carbide Production Workers. *Chest* 86:549-557 (1984).

*Proceedings of the VIIth International Pneumoconioses Conference*  
*Transactions de la VIIe Conférence Internationale sur les Pneumoconioses*  
*Transacciones de la VIIa Conferencia Internacional sobre las Neumoconiosis*

Parte I  
Tome I  
Parte I



Pittsburgh, Pennsylvania, USA—August 23–26, 1988  
Pittsburgh, Pennsylvanie, Etats-Unis—23–26 aout 1988  
Pittsburgh, Pennsylvania EE. UU—23–26 de agosto de 1988



**U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES**  
Public Health Service  
Centers for Disease Control  
National Institute for Occupational Safety and Health

**CDC**  
CENTERS FOR DISEASE CONTROL

## **Sponsors**

International Labour Office (ILO)  
National Institute for Occupational Safety and Health (NIOSH)  
Mine Safety and Health Administration (MSHA)  
Occupational Safety and Health Administration (OSHA)  
Bureau of Mines (BOM)

September 1990

## **DISCLAIMER**

Sponsorship of this conference and these proceedings by the sponsoring organizations does not constitute endorsement of the views expressed or recommendation for the use of any commercial product, commodity, or service mentioned.

The opinions and conclusions expressed herein are those of the authors and not the sponsoring organizations.

**DHHS (NIOSH) Publication No. 90-108 Part I**