

## SIMILARITIES IN LUNG CANCER AND RESPIRATORY DISEASE MORTALITY OF VERMONT AND NEW YORK STATE TALC WORKERS

S.H. LAMM, M.D. • J.A. Starr, MSc

Consultants in Epidemiology and Occupational Health, Inc., Washington, DC 20007

### ABSTRACT

The risks from malignant and non-malignant respiratory deaths of New York State and Vermont State talc workers with at least one year of employment have been compared for both miners and millers. The mortality patterns are similar. In both areas, the talc miners have a 4.5 fold risk of lung cancer, and the talc miners have no increased risk of lung cancer. In both areas, all workers appear to have an increased risk of non-infectious, non-neoplastic respiratory disease (NNRD) mortality, although only the Vermont millers show a statistically significantly elevated risk (7.9 fold). Thus, although the New York talc has been described as asbestiform talc and the Vermont talc as non-asbestiform talc, the mortality patterns of the workers appear to be inconsistent with that classification in that their lung cancer mortality rates are no different and only the Vermont talc millers show a significantly increased NNRD mortality.

### INTRODUCTION

Studies of talc miners and millers in the New York and Vermont talc industry include analyses of mortality, morbidity, industrial hygiene, and mineralogy. Mineralogical differences between the two talcs have been highlighted. The upstate New York talc contains an elongated particulate not found in the Vermont talc that is considered by scientists at the National Institute for Occupational Safety and Health (NIOSH) as tremolitic asbestos and by scientists at the Bureau of Mines and at the company that owns the plant as true talc particulates and as prismatic non-asbestiform tremolite. NIOSH has called the New York State talc asbestiform talc and the Vermont talc non-asbestiform talc. Leaving the question of the mineralogical label of these particulates to the mineralogists, we have elected to examine the respiratory health outcomes of the employees at these two talc industries.

### MATERIALS

The initial shaft of the New York State talc plant was sunk in 1947. Mining and milling operations started in 1948. The mortality experience (1947 through 1978) of all persons hired at the plant between 1947 and 1977 has been reported.<sup>1</sup> Mortality analysis was restricted to the 705 male employees (all caucasian). None of the 36 women employees had died of a respiratory condition. Sixty percent of the men worked at the plant for at least one year; twenty percent for two months to one year; and twenty percent for less than two months. Mortality analysis was reported separately for the 280 white male employees employed at the talc plant for less than one year and for the 425 white male employees employed for at least one year. That report<sup>1</sup> suggested that prior employment jobs accounted for the lung cancer rate.

In-plant job records and prior employment histories on the job applications were analyzed. Employees were classified from the inplant job records as miners (187 worked exclusively in the mine), millers (152 worked exclusively in the mill), and others (34 worked in both the mine and the mill, 11 worked neither in the mine or the mill, and 41 had uninformative records).

The cohort of white male employees of the Vermont talc industry was developed from the records of the Vermont State Health Department's annual radiographic survey of employees of the dusty trades, begun in 1937. Selevan et al. of the National Institute for Occupational Safety and Health (NIOSH) defined the Vermont talc study cohort<sup>2</sup> as all white males in the Vermont talc industry on or after January 1, 1940 with at least one year of talc employment prior to January 1, 1970. Individuals who had at least two radiographs in the file and who had worked for any of five talc companies in three geographic areas of Vermont were eligible for the study. Mortality follow-up was continued through 1975 of the 392 men determined to belong to the cohort.

Health Department and company records were scrutinized to determine their job assignments, and each cohort member was classified as a miner after having had one year of exposure in the mine and/or as a miller after having had one year of exposure in the mill. 225 workers were classified as miners; 163 workers were classified as millers (of whom 47 had also been classified as miners); and 51 were not classifiable.

### METHODS

This report compares standardized mortality ratios (SMRs)

for malignant and non-malignant respiratory causes of death for miners and millers with at least one year of experience in the Upstate New York talc (said to be asbestiform) industry with those in the Vermont State talc (said to be non-asbestiform) industry. Comparison is reasonable, despite the differences in classification variables between the two studies.

## RESULTS

The risks of lung cancer and of non-infectious, non-neoplastic respiratory disease (NNRD) for employees with at least one year in the mines or mills of New York State or Vermont State talc industries are presented, analyzed, and discussed below.

### Respiratory Mortality of New York and Vermont Talc Workers

	Observed/Expected Ratios		Standardized Mortality Ratios	
	New York	Vermont	New York	Vermont
<b>Lung Cancer</b>				
Millers	1/1.41	2/1.96	071	102
Miners	5/1.15	5/1.09	460*	435*
Others	0/0.55	0/0.61	---	---
Total	6/3.11	7/3.66	1.92	1.91
<b>NNRD</b>				
Millers	2/0.74	7/0.89	270	787*
Miners	2/0.49	2/0.56	408	357
Others	2/0.38	2/0.34	528	588
Total	6/1.61	11/1.79	373*	615*

\* =  $p < 0.05$ , two-tailed Poisson test

The risk of malignant disease of the lung (lung/respiratory cancer) is not increased for millers but is significantly increased (4.5 fold) in talc miners both in New York (4.60) and in Vermont (4.35). No difference in risk is seen between miners and millers of New York and of Vermont (Figure 1). These data are sufficiently strong to rule out with eighty percent confidence an underlying relative risk for New York miners vs. Vermont miners of 1.7 and with about ninety five percent certainty an underlying risk of greater than 2.0.



Figure 1. Respiratory or lung cancer mortality risk for miners and millers of New York State and Vermont State talc.

The risk of non-malignant respiratory disease (excluding pneumonia and influenza), i.e., NNRD has a significantly increased risk (almost eight-fold) for Vermont talc millers but not for New York talc millers (risk of 2.7, not significant). The risks for NNRD for miners are calculated to be 4.1 and 3.6 (both non-significant) for those from New York and Vermont, respectively (Figure 2).

As for other respiratory system deaths, influenza or pneumonia caused the death of one New York State talc worker (0.9 expected) but no Vermont talc miner (0.7 expected) or miller (0.8 expected). Mesothelioma caused the death of one New York State talc man (15 years after hire which followed 28 years in mining and construction) and of one Vermont talc man.

## DISCUSSION

We have attempted to assemble similarly defined cohorts of New York State and Vermont State talc workers in order to compare the respiratory mortality risks of their miners and millers. The exposures of millers generally exceed that of miners by a factor of two to six. Nonetheless, both groups demonstrate a similar excess lung cancer risk only for their millers and not for their miners. The similar lung cancer risks of the two groups of talc workers exposed to the differently described talcs suggest that the elongated particulates seen in the New York State talc have not introduced an increased lung cancer risk. We further observe that the risk of non-infectious, non-neoplastic respiratory death, while apparently increased in all groups, is significantly elevated only among the Vermont millers.

Standardized mortality ratios (SMRs) were calculated for each group based on age-specific, calendar time-specific, cause-specific mortality rates for white males. The New York State study SMRs had been calculated using U.S. rates with death certificates coded according to the eighth revision of the International Classification of Diseases (ICD). The Vermont State study SMRs were first calculated using U.S. rates and then recalculated by its authors using Vermont State rates for non-malignant respiratory disease and respiratory cancer




Figure 2. Non-infectious, non-malignant respiratory disease (NNRD) mortality risks for miners and millers of New York State and Vermont State talc.

COMPARATIVE LUNG MORTALITY RISKS of VERMONT and NEW YORK STATE TALC WORKERS with at least one year experience at Talc Plant

	Vermont		New York		New York		New York	
	O/E	SMR	O/E	SMR	O/E	SMR	O/E	SMR
<b>All Emp &gt;1 yr.</b>								
All Causes	44/37.15	118	64/49.83	128	118/83.58	141*	54/33.75	160*
All Cancers			15/9.55	157	26/15.7	165*	9/6.15	146
Lung Cancer	6/3.61	163	6/3.11	193	12/5.01	240*	6/1.90	316*
NNRD	11/1.79	615*	6/1.61	372*	6/2.64	227	0/1.03	---
<b>Millers</b>								
All Causes			20/21.74	92	35/30.97	113	15/9.23	163
All Cancers			3/4.23	71	6/5.94	101	3/1.71	175
Lung Cancer	2/1.96	102	1/1.41	71	1/1.92	52	0/0.51	---
NNRD	7/1.89	787*	2/0.74	270	2/1.02	196	0/0.28	---
<b>Miners</b>								
All Causes			31/16.76	185*	50/26.32	190*	19/9.56	199
All Cancers			10/3.23	310*	15/5.00	300*	5/1.77	282
Lung Cancer	5/1.15	435*	5/1.09	460*	9/1.66	543*	4/0.57	701*
NNRD	2/0.56	357	2/0.49	408	2/0.77	260	0/0.28	---
<b>Others</b>								
All Causes			13/11.33	115	33/26.29	126	20/14.96	134
All Cancers			2/2.09	96	5/4.76	105	3/2.67	112
Lung Cancer	0/0.55	---	0/0.61	---	2/1.43	140	2/0.82	244
NNRD	2/0.34	588.0	2/0.38	526	2/0.85	235	0/0.47	---

## COHORT DEFINITION

LUNG CANCER

Cohort Variable	NEW YORK	VERMONT	NEW YORK	
Gender	Male	Male	VERMONT	
Race	White	White	Observed/Expected	
Employment Dates	1947-1977	1940-1969	Millers	1/1.41      2/1.96
Employment Duration	One Year +	One Year +	Miners	5/1.15      5/1.09
Mortality Dates	1947-1978	1940-1975	Others	0/0.55      0/0.61
<b>Cohort Numbers</b>				
Miners	152	163		
Millers	187	225	SMR	
			Millers	71      102
			Miners	<b>460      435</b>
			Others	---      ---

4/7/88

COMPARATIVE LUNG MORTALITY RISKS of  
VERMONT and NEW YORK STATE TALC WORKERS  
with at least one year experience at Talc Plant

All Emp >1 yr. LATENCY (Years)	Vermont		New York		New York		New York	
	O/E	SMR	O/E	SMR	O/E	SMR	O/E	SMR
					Ever employed		< one Year	
0-4			0/0.27	---	0/0.42	---	0/0.15	---
5-9			0/0.31	---	0/0.49	---	0/0.18	---
10-14			1/0.45	224	1/0.69	145.0	0/0.24	---
15-19			2/0.60	331	2/0.98	205.0	0/0.38	---
20-24			3/0.79	378	8/1.29	623*	5/0.50	1000
25-29			0/0.65	---	1/1.09	92.0	1/0.44	227
30+			0/0.04	---	0/0.05	---	0/0.01	---
Total			6/3.11	193	12/5.01	240*	6/1.90	316*
0-9			0/0.58	---	0/0.91	---	0/0.33	---
10-19			3/1.05	285	3/1.67	180	0/0.62	---
20-29			3/1.44	208	9/2.38	378*	6/0.94	638*
30+			0/0.04	---	0/0.05	---	0/0.01	---
Total			6/3.11	193	12/5.01	240*	6/1.90	316*
0-4			0/0.27	---	0/0.42	---	0/0.15	---
5-14			1/0.76	132	1/1.18	85	0/0.42	---
15-24			5/1.39	360*	10/2.27	441*	5/0.88	568*
25+			0/0.69	---	1/1.14	87	1/0.45	222
Total			6/3.11	193	12/5.01	240*	6/1.90	316*

4/7/88

COMPARATIVE LUNG MORTALITY RISKS of  
VERMONT and NEW YORK STATE TALC WORKERS  
with at least one year experience at Talc Plant

		Vermont		New York	
		O/E	SMR	O/E	SMR
All Causes	Emp >1	44/37.15	118.0	64/49.83	128
All Cancers	Emp >1			15/9.55	157
Lung Cancer	Emp >1	6/3.61	163	6/3.11	193
NRPD	Emp >1	11/1.79	615	6/1.61	372
Pneumonia/Influ	Emp >1	0/1.89	000	1/0.9	109
All Causes	Millers			20/21.74	92
All Causes	Miners			31/16.76	185
All Causes	Others			13/11.33	115
All Cancers	Millers			3/4.23	71
All Cancers	Miners			10/3.23	310
All Cancers	Others			2/2.09	96
Lung Cancer	Millers	2/1.96	102	1/1.41	71
Lung Cancer	Miners	5/1.15	435	5/1.09	460
Lung Cancer	Others	0/0.55	---	0/0.61	---
NRPD	Millers	7/1.89	787	2/0.74	270
NRPD	Miners	2/0.56	357	2/0.49	408
NRPD	Others	2/0.34	588	2/0.38	526
4/7/88	Bold =	p <0.05			
Pneumonia/Influ	Millers	0/.83	000		
Pneumonia/Influ	Miners	0/.67	000		
Pneumonia/Influ	Others	0/.39	000		

Standardized Mortality Ratios		
	Vermont	New York
<b>Lung Cancer</b>		
Millers	102	71
Miners	435	460
Others	---	---
<b>NRPD</b>		
Millers	787	270
Miners	357	408
Others	588	526

**NON-INFECTIOUS, NON-  
MALIGNANT  
RESPIRATORY DISEASE**

VERMONT	NEW YORK	
Observed/Expected		
Miller	2/0.74	7/0.89
Miners	2/0.49	2/0.56
Others	2/0.38	2/0.34
 SMR		
Millers	270	787
Miners	408	357
Others	526	588

with death certificates coded according to the seventh revision of the ICD. This report bases the SMRs on the U.S. rates.

The New York State study reports lung cancer as their measure of malignant respiratory disease and NNRD (non-infectious, non-neoplastic respiratory disease) as their measure of non-malignant respiratory disease. The Vermont State study reports respiratory cancer as their measure of malignant respiratory disease and ONMRD (other non-malignant respiratory disease) as their measure of non-malignant respiratory disease. Both NNRD and ONMRD are terms for total non-malignant respiratory disease, excluding influenza and pneumonia. We have used the labels of lung cancer and NNRD to represent the malignant and non-malignant respiratory disease measures.

Twelve of the thirteen respiratory cancers among the New York State talc workers were lung cancers. The thirteenth case was a man whose five years at the plant included three months as a laborer/oiler in the talc mill and ended with death from mediastinal cancer. Re-analysis of the New York State data as respiratory cancer rather than lung cancer would have reduced the SMR estimates by about 5% but not have altered the comparison between the miners and millers. Both the

New York and the Vermont data are compared against U.S. mortality rates.

The Vermont data included persons with experience in both the mine and the mill in each category; the New York data separated them out. There were only 34 such New York workers with experience in both the mine and the mill. Less than 0.1 lung cancer and less than 0.1 NNRD deaths were expected among them, and none were observed. Including this group among the miners and the millers of New York State would not have affected the results.

Studies of both cohorts lack full information on smoking history. Each indicates that most of the lung cancer cases were known to be cigarette smokers, but data on smoking appears to be inadequate for both cohorts. There is no evidence that miners and millers differ in their smoking habits. Thus, it is unlikely that the differences observed in these comparisons could be due to differences in smoking between groups.

The mortality of the experienced employees of the New York and Vermont cohort who worked other than in the mine or the mill for a year were also examined. There were no lung cancer deaths. Each group had two NNRD deaths, yielding non-significant risks of 5.9 for those from Vermont and 5.3 for those from New York.

While the NNRD mortality may be due to dust exposures at the talc plants, the etiology of the lung cancer is less clear. The NIOSH authors<sup>2</sup> concluded that talc dust was unlikely to be the cause of the respiratory cancer, since the risk was seen only in the miners and not seen among the millers, a group with probable higher dust exposure. Radon daughter measurements in the New York mine do not explain the finding. The presence of a particulate in New York dust and not in Vermont talc dust cannot explain the difference.


The CEOH study<sup>1</sup> had supported the hypothesis of risk from prior employments as the explanation for the lung cancer risk of the New York State talc workers, however, that hypothesis has not been examined for the Vermont talc workers. Further study of both cohorts should be undertaken to explain the mortality patterns seen. The small number of cases in either group will probably be a hindrance to a full and clear explanation. Both cohorts should probably be extended to include later employees and the period of follow-up should be brought more current by at least a decade. A four-fold risk of lung cancer seen in two different studies of talc miners (but not millers) cries for an explanation.

#### REFERENCES

1. Lamm, S.H., Levine, M.S., Starr, J.A., Tirey, S.L.: Analysis of Excess Lung Cancer Risk in Short-Term Employees. *Am. J. Epi.* 127(6):1202-1209 (1988).
2. Selevan, S.G., Dement, J.M., Waggoner, J.K., Froines, J.R.: Mortality Patterns among Miners and Millers of Non-Asbestiform Talc: Preliminary Report. *J. Environ. Path. and Tox.* 2:273-284, 1979.

*Proceedings of the VIIth International Pneumoconioses Conference* Part  
*Transactions de la VIIe Conférence Internationale sur les Pneumoconioses* Tome  
*Transaciones de la VIIa Conferencia Internacional sobre las Neumoconiosis* Parte

**II**



Pittsburgh, Pennsylvania, USA—August 23–26, 1988  
Pittsburgh, Pennsylvanie, Etats-Unis—23–26 aout 1988  
Pittsburgh, Pennsylvania EE. UU—23–26 de agosto de 1988



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES  
Public Health Service  
Centers for Disease Control  
National Institute for Occupational Safety and Health



## **Sponsors**

**International Labour Office (ILO)**  
**National Institute for Occupational Safety and Health (NIOSH)**  
**Mine Safety and Health Administration (MSHA)**  
**Occupational Safety and Health Administration (OSHA)**  
**Bureau of Mines (BOM)**

**November 1990**

## **DISCLAIMER**

Sponsorship of this conference and these proceedings by the sponsoring organizations does not constitute endorsement of the views expressed or recommendation for the use of any commercial product, commodity, or service mentioned.

The opinions and conclusions expressed herein are those of the authors and not the sponsoring organizations.

**DHHS (NIOSH) Publication No. 90-108 Part II**