

ORIGINAL ARTICLE

The validation of work-related self-reported asthma exacerbation

Aimee R Bolen, Paul K Henneberger, Xiaoming Liang, Susan R Sama, Peggy A Preusse, Richard A Rosiello, Donald K Milton

Occup Environ Med 2007;**64**:343–348. doi: 10.1136/oem.2006.028662

See end of article for authors' affiliations

Correspondence to: P K Henneberger, National Institute for Occupational Safety and Health/CDC, M/ S H2800, 1095 Willowdale Road, Morgantown, WV 26501, USA; pkh0@cdc.gov

Accepted 28 October 2006
Published Online First
19 December 2006**Objective:** To determine the validity of work-related self-reported exacerbation of asthma using the findings from serial peak expiratory flow (PEF) measurements as the standard.**Methods:** Adults with asthma treated in a health maintenance organisation were asked to conduct serial spirometry testing at home and at work for 3 weeks. Self-reported respiratory symptoms and medication use were recorded in two ways: a daily log completed concurrently with the serial PEF testing and a telephone questionnaire administered after the PEF testing. Three researchers evaluated the serial PEF records and judged whether a work relationship was evident.**Results:** 95 of 382 (25%) working adults with asthma provided adequate serial PEF data, and 13 of 95 (14%) were judged to have workplace exacerbation of asthma (WEA) based on these data. Self-reported concurrent medication use was the most valid single operational definition, with a sensitivity of 62% and a specificity of 65%.**Conclusions:** A work-related pattern of self-reported asthma symptoms or medication use was usually not corroborated by serial PEF testing and failed to identify many people who had evidence of WEA based on the serial PEF measurements.

Work-related asthma (WRA) is the most common non-asbestos related occupational respiratory disease in the US and in most industrialised nations.^{1–4} It accounts for about 25% of all reported occupational lung diseases in the UK⁵ and an estimated 15% of asthma cases in the US are work related.^{4 6–8} In the US, the Sentinel Event Notification System for Occupational Risks (SENSOR) case-classification scheme for WRA encompasses both new-onset asthma and work-aggravated asthma.⁹ Work-aggravated asthma is defined as the aggravation or exacerbation of pre-existing asthma due to occupational exposures. Based on SENSOR data of four states from 1993 to 1995, 19.1% of WRA cases were in the aggravation category.⁹ These cases had similar adverse outcomes as the new-onset cases, such as persistent breathing problems and the need for emergency care.¹⁰

Questionnaires that gather information on symptoms and exposures are frequently used to assess WRA.¹¹ Questionnaires are inexpensive and feasible for large population-based studies, but patients must be able to clearly perceive their respiratory status for their responses to be useful. Unfortunately, asthma patients' perceptions of dyspnoea are not always consistent with measured levels of air flow obstruction.¹² One study showed that 15% of people with asthma failed to be aware of any discomfort after a 20% reduction in their forced expiratory volume in one second.¹³ The inability of a patient to accurately perceive substantial changes in his or her lung function results in inaccurate reporting of symptoms due to either blunted perception or over perception.¹⁴ Therefore, self-reported symptoms could not be effective surrogates for more objective diagnostic tools in determining the contribution of exposures at work to the onset or exacerbation of asthma.

Serial measurements of peak expiratory flow (PEF) are an established objective aid in the diagnosis of WRA.^{15 16} The test allows for simple, inexpensive measurement of lung function outside of a clinic or laboratory.^{15 17–19} Serial PEF measurements have a high sensitivity and even higher specificity for WRA.^{19–21} For example, when using specific inhalation challenges as the

standard for occupational asthma, serial PEF measurements were found to have a sensitivity of 70–75% and a specificity of 94–100%.^{20 21}

The primary objective of this investigation was to validate work-related self-reported exacerbation of asthma using the findings from serial PEF measurements as the standard.

METHODS

The protocol for the study was reviewed and approved by the National Institute for Occupational Safety and Health (NIOSH) Human Subjects Review Board, and the Human Subjects Review Board of the contracted Health Maintenance organization (HMO), Fallon Clinic Research Department in Eastern and Central Massachusetts, Worcester, Massachusetts, USA. All information was de-identified before it was sent to NIOSH.

Participants

The current validation study was part of the Workplace Exacerbation of Asthma (WEA) Research Project, which was initiated by NIOSH in 2000 to investigate the frequency, causes and consequences of work-related exacerbation of asthma. NIOSH contracted with researchers at the Fallon Clinic in Massachusetts to conduct data collection. Detailed methods for the baseline study are presented elsewhere.²² In brief, both electronic and paper medical records were reviewed to identify adults aged 18–44 years with diagnosed asthma who had been enrolled in the Fallon Community Health Plan for at least 6 months.

From the 598 baseline study participants, candidates for the current validation study were identified. However, 192 baseline study participants were not invited to take part in the validation

Abbreviations: HMO, Health Maintenance organization; NIOSH, National Institute for Occupational Safety and Health; OASYS-2, Occupational Asthma System 2; PEF, peak expiratory flow; SENSOR, Sentinel Event Notification System for Occupational Risks; WEA, workplace exacerbation of asthma; WRA, work-related asthma

study because they were no longer enrolled in Fallon Community Health Plan. Research staff sent materials, including an invitation to participate and an explanation of the study, to the remaining 406 people.

Materials

Office spirometry was accomplished using an OMI Sensormedics 922 spirometer (Occupational Marketing, Houston, Texas, USA). Collection of serial PEF measurements was achieved through the use of ndd EasyOne portable spirometers (ndd Medical Technologies, Andover, Massachusetts, USA). Both the office and portable spirometers were calibrated to meet American Thoracic Society standards.²³ The portable devices were self-recording, capable of storing large numbers of spirometric measurements and included an alarm that sounded at regular intervals to remind participants to test themselves. In addition, participants were prompted at each trial to complete a daily log that recorded symptoms and medication use. Participants entered their answers by means of a keypad built into the side of the spirometer. All the information from the spirometer was date and time stamped. Information was also used from two other survey instruments: the baseline study questionnaire and a telephone questionnaire conducted after the serial PEF testing.

Serial spirometry

Before the start of serial testing, participants were asked to complete an office spirometry session, where they were given instruction on using the portable spirometers. Measurements of PEF were performed for 3 weeks at home and at work. The participants were asked to conduct at least five trials each day with at least three forced expiratory manoeuvres performed at each trial.^{17 18 20 24} The best of the three measurements was used for analysis. On completion of both the office spirometry and at least 16 days of serial testing, the participants were given \$200 as compensation for the time it took to complete these tests.

Self-reporting

Work-related self-reported exacerbation of asthma was assessed in two ways. First, symptoms and medication use concurrent with serial testing were gathered from the daily log. Questions included (1) In the last 2 h, did you have a cough

attack, wheeze, chest tightness or shortness of breath? (2) In the last 2 h, did you use your fast-acting inhaler? Responses to each of these questions were compared between work days and non-work days. Participants with a greater percentage of positive responses on all work days than on all non-work days were classified as having work-related self-reported exacerbation of asthma. Two different operational definitions were derived, shown below.

(1) Reported symptoms more often on work days versus non-work days.

(2) Reported medication use more often on work days versus non-work days.

Second, symptoms and medication use were collected by a post-test telephone questionnaire that asked about the previous 3 weeks of testing. Those who reported that their symptoms or medication use varied by whether they were at or away from work were asked the following questions: (1) Did your asthma symptoms get better or worse away from work? (2) Did you use your inhaler or nebuliser more or less on work days? The following operational definitions were based on answers to the post-test questions:

(1) Asthma symptoms improved away from work.

(2) Medication use increased on work days.

Validation standard

The participants returned the portable spirometer to the HMO after completion of testing. Research staff entered the data into the Occupational Asthma System 2 (OASYS-2), which is a computer-assisted diagnostic aid that was developed to detect asthma-work relationships using serial PEF data.^{25 26} The computer program generates plots of daily maximum, mean and minimum PEF values, and a work-relatedness score ranging from 1 to 4. The probability of WRA increases as the score increases, and a score of >2.5 has a sensitivity of 75% and a specificity of 94% when using specific inhalation challenge as the standard for occupational asthma.²⁰

Three researchers familiar with the interpretation of serial PEF measurements reviewed all the OASYS-2 records. These records were judged to have inadequate data for evaluating a work relationship if participants completed <2 weeks of ≥ 4 trials a day, provided an insufficient number (ie, <2) of work-day/rest-day complexes, or had a respiratory infection during the testing period. Infections were reported to the field staff, who in turn, recorded the information and provided it to the researchers who analysed the data. A complex was defined as a series of at least three work days that was preceded and followed by at least two rest days, or at least two rest days preceded and followed by at least three work days.




Figure 1 Flowchart of the participants

Table 1 Demographics of employed, potential participants based on completion of office and serial spirometry

	Completed neither office nor serial spirometry	Completed office spirometry Did not complete serial spirometry	Completed serial spirometry
n	204	83	95
Female (%)	64	66	71
Mean age (years)	33.5	33.6	34.2
White (%)	95	93	94
Ever smoked (%)	49	28*	34*
Atopy (%)	64	63	68
Moderate asthma (%)	37	35	51*

* $p < 0.05$ for comparison with those who completed neither office nor serial spirometry.

Table 2 Comparison between participants positive and negative for workplace exacerbation of asthma based on serial peak expiratory flow

	WEA based on serial PEF	
	Positive	Negative
n	13	82
Female (%)	85	68
Mean age (years)	33.7	34.3
Ever smoked (%)	31	34
With atopy (%)	54	71

PEF, peak expiratory flow; WEA, workplace exacerbation of asthma.

The three reviewers independently judged the remaining OASYS-2 records to be either positive or negative for WEA based on serial PEF (WEA-PEF). Reviewers were unaware of any work-related respiratory problems that participants might have reported in the baseline and validation studies. The primary features considered when judging the relationship to work were the OASYS-2 score, lower PEF values while working and higher PEF values while away from work and increased diurnal variation on work days. Diurnal variation was defined as the difference between maximum PEF and minimum PEF divided by the mean PEF for the whole period. The category agreed upon by the majority of reviewers was used for analysis.

Data analysis

All statistical analyses were accomplished using SAS 9.1 software. The analysis included comparisons of demographic features and validity calculations. Data from the baseline study questionnaire were used for gender, age, race, smoking status and presence of atopy. Participants were considered positive for atopy if they had ever been told by a doctor that they had hay fever or nasal allergies. To test for statistical significance when making comparisons, the Student's *t* test was used for continuous variables, and the continuity corrected χ^2 statistic and Fisher's exact test were used for categorical variables. The Fisher's exact test was used when expected cell counts were <5. A significance level of 0.05 was applied for all tests.

Validity measurements were calculated to characterise the ability of self-reported symptoms and medication use to identify who did and who did not have work-related disease, using WEA-PEF status as the standard. Contingency tables were created for single operational definitions and either-or/both-and combinations of the operational definitions (*n* = 66). Sensitivity, specificity and Youden's Index were calculated for each table. Youden's Index is the sum of the sensitivity and specificity (expressed as fractions) minus one.²⁷ When sensitivity and specificity are equal to one, Youden's Index is zero, and the operational definition used was no better than random chance.²⁸

RESULTS

Of the 406 adults with asthma who were asked to participate in the validation study, 382 were employed at the time of testing. Figure 1 summarises the disposition of these 382 adults. In all, 178 of 382 (47%) participants completed office spirometry; the other 204 declined to participate or did not respond. Of those who completed office spirometry, 40 did not complete at least 2 weeks of serial spirometry testing, whereas 138 met the 2-week requirement. Forty-three of 138 did not have reviewable records due to an insufficient number of work days or too few consecutive work days (*n* = 28), an insufficient number of rest days (*n* = 6), skipped days mid-testing (*n* = 5) or the presence of a respiratory infection during testing (*n* = 4). In total, 95 of 382 (25%) participants completed serial spirometry testing and provided adequate data to be evaluated for WEA-PEF.

We compared demographic features among the 204 employed candidates for participation who refused or never responded to the invitation, the 83 who completed office spirometry but not serial testing and the 95 who completed both office and serial spirometry. There was little variation in gender, age, race or atopy by participation status (table 1). However, the two groups which completed office spirometry had a lower percentage of ever-smokers than the non-participants (*p* < 0.05) and those groups that completed the serial spirometry had a higher percentage of moderate asthma. The 24 adults with asthma who were not invited due to unemployment were more likely to be female than those who were invited (*p* = 0.01), but were approximately of the same

Table 3 Prevalence of workplace exacerbation of asthma based on serial peak expiratory flow within occupational categories

Occupational code*	n	Occupational category	Workplace exacerbation of asthma, n (%)
001-359	46	Management, professional and business occupations	6 (13)
001-099	11	Management, business and financial	0 (-)
100-219	15	Professional and related	1 (7)
220-299	14	Education, training, media and library	4 (29)
300-359	6	Healthcare, practitioner and technical	1 (17)
360-469	13	Service occupations	3 (23)
360-369	5	Healthcare support	2 (40)
430-469	8	Food preparation, personal care and child care	1 (13)
470-599	27	Sales and office occupations	2 (7)
470-499	6	Sales, retail and related occupations	1 (17)
500-599	21	Office and administrative support	1 (5)
620-979	8	Construction, extraction and maintenance occupations	1 (13)
No code	1	Student	1 (-)
001-999	95	All occupations	13 (14)

*Occupational codes based on 2000 US Census.²⁹

age. The 192 adults with asthma who were not invited owing to insurance compliance were similar to those who were invited with respect to gender and age (data not shown).

Upon evaluation of the OASYS-2 records, 13 of 95 (14%) participants were positive for WEA-PEF. Those with WEA-PEF were more likely to be female and less likely to be atopic, although these differences were not statistically significant (table 2). Also, age and smoking status did not vary by WEA-PEF status.

Twenty-five of 95 (26%) participants changed occupations from baseline to the time of testing. Three of these people were performing the same tasks at a different location. The prevalence of WEA-PEF was higher among those who changed employment (18%) compared to those who did not (12%), but the difference was not statistically significant ($p = 0.49$). The broad occupational category with the highest prevalence (23%) of WEA-PEF was service occupations, whereas the lowest prevalence (7%) was in the sales and office occupations (table 3). The highest prevalences by specific categories were 40% (2 of 5) for healthcare support (in service occupations) and 29% (4 of 14) for education, training, media and library (in management, professional and business occupations).

Fulfillment of the operational definitions of work-related symptoms and medication use was about three times more common when based on reports concurrent with the serial testing as compared with reports from the post-test telephone interview. Specifically, concurrent and post-test reports were 48% vs 14% for symptoms and 39% vs 12% for medication use, respectively.

Table 4 Sensitivity, specificity and Youden's Index for operational definitions based on self-reported symptoms and medication use*

Operational definitions	Sensitivity (%)	Specificity (%)	Youden's Index
Symptoms and medication use reported concurrent with serial testing			
Symptoms	62	54	0.16
Medication use	62	65	0.27
Symptoms and medication use reported post-test			
Symptoms	15	87	0.02
Medication use	15	89	0.04
Combinations of self-reported operational definitions †			
Combinations of two			
Both Ct	54	78	0.32
Ct Rx or post Rx	69	61	0.30
Combinations of three			
Both Ct or post Sx	62	68	0.30
Both Ct or post Rx	62	71	0.33
Either Sx and Ct Rx	54	76	0.30
(Ct Rx or post Sx) and Ct Sx	54	76	0.30
Either Rx and Ct Sx	54	77	0.31
Both post or Ct Rx	69	62	0.31
Combinations of four			
Both Ct or both Sx	54	76	0.30
Both Ct or both post	62	73	0.35
Both Ct or (Ct Sx and post Rx)	54	77	0.31
Both Ct or (Ct Rx and post Sx)	54	76	0.30
(Ct Sx or post Sx) and (Ct Rx or post Rx)	62	73	0.35
(Ct Sx or post Rx) and (Ct Rx or post Sx)	62	71	0.33

Ct, concurrent; post, post-test; Rx, medication use; Sx, symptoms.

*With findings from serial peak expiratory flow as the standard for workplace exacerbation of asthma.

†Combinations of operational definitions were included if Youden's Index ≥ 0.30 .

Sensitivity, specificity and Youden's Index were calculated for operational definitions based on both types of self-reported measures. Table 4 shows the single operational definitions, and also all combinations of definitions that had a Youden's Index ≥ 0.30 (top 1/5 of all the combinations).

None of the operational definitions for work-related self-reported exacerbation of asthma had very high values for both sensitivity and specificity (table 4). The highest specificity was observed for post-test symptoms (87%) and medication use (89%), but both had a very low sensitivity of 15%. The highest Youden's Index for a single self-reported operational definition was 0.27 for concurrent medication use, with a sensitivity of 62% (8 of 13 cases identified) and a specificity of 65% (53 of 82 cases with serial PEF negative for WEA). When the operational definitions were expanded to include either concurrent or post-test medication use, the sensitivity increased to 69%, which was the highest level attained. Nine of the 13 true cases were identified, and the specificity (61%) was nearly the same as that observed for concurrent medication use by itself. A sensitivity of 69% was also achieved for the operational definition of both post-test symptoms and medication use or concurrent medication use.

DISCUSSION

Serial PEF testing by adults with asthma

Most study candidates did not provide useful serial spirometry data, even though they were offered money as compensation for their time. Only 138, or about one-third of the 382 eligible, completed at least 2 weeks of serial testing. In addition, 34 of 138 had data that could not be analysed by the OASYS-2 software owing to too few work or rest days, or too few consecutive work days. This last group was the result of part-time or non-traditional work schedules. Lack of compliance and inadequate data collection have been observed in studies limited to workers with suspected WRA.^{30 31} The low compliance in this study suggests that serial PEF testing is of limited usefulness in population-based or quasi-population-based studies, at least when trying to determine work-relatedness. This problem could possibly be overcome by offering higher monetary incentives that exceed merely compensating participants for their time, and by initiating more frequent (perhaps daily) contact with the participants to keep them on track with the study.

The adults with asthma who completed serial testing were more likely to have moderate asthma rather than mild asthma and less likely to have ever smoked than those who completed neither office nor serial spirometry. However, neither characteristic was associated with a positive serial test result. At the same time, the adults with asthma who did complete the serial testing were very similar to those who did not with respect to gender, age, race and atopic status. Thus, the findings based on the 95 participants are probably representative of the entire group of candidates for participation. Moreover, we determined that 14% of the participants had WEA-PEF, which is similar to the prevalence estimate for WRA in the American Thoracic Society statement.⁸ Still, we cannot entirely rule out that selection bias could have occurred, especially given the low rate of compliance, and this could have biased the findings in either direction.

A major criticism of serial self-testing has been the potential for inaccurate recording of PEF measurements by participants.^{24 32} In this study, participants used self-recording portable spirometers, making the recording of PEF measurements more accurate. A learning effect has also been reported while people are becoming accustomed to using the spirometers. The current participants were oriented to the portable spirometers in person, and the devices provided coaching

messages each time they were used. These attributes helped to assure that the PEF readings were acceptable and accurately recorded.

Overall, this study shows that serial PEF measurements, once collected, can be used to determine work-related patterns of asthma. Serial spirometry is an inexpensive method of screening for WRA that is more objective than self-report and readily available in many countries. A drawback is that our standard has limitations with regard to measuring WRA, with a sensitivity of approximately 70–75% for occupational asthma when compared with specific inhalation challenge.^{20–21} If the same limitation of serial spirometry applies to WEA, then our estimates of sensitivity and/or specificity for self-reports could be inaccurate.

Occupations with a higher prevalence of WEA

The highest prevalence of WEA–PEF was found in the occupation subcategory of healthcare support that includes nursing aides, home health aides, dental assistants and so on. In a review of SENSOR data from 1993 to 1997, exploring the association of WRA and cleaning products, nurse's aides had the second highest prevalence of WRA (20%).³³ The other agents associated with WRA among healthcare workers were latex and indoor air pollution.³⁴ The second highest prevalence of WEA was found in the occupational subcategory of education, training, media and library. In a review of SENSOR data from Massachusetts (1993–2004), educational services industry had the second highest prevalence of WRA.³⁵

The value of self-reports in the identification of work-exacerbated asthma

In some other studies that have examined the value of self-reported work-related symptoms, the participants were workers with asthma seeking compensation.³⁶ Participants in this study were adults with asthma who were being treated in an HMO. In this setting, the false positives based on self-reports are unlikely to represent intentional falsification to gain compensation, and are more likely due to misperception of causation or the presence of work-related symptoms in the absence of significant changes in the PEF. False positives were common in this study—for example, of the 37 participants who fulfilled the operational definition based on concurrent medication use, 29 (78%) were negative for WEA based on PEF. Consistent with this observation, some researchers have suggested that work-related exacerbation of asthma is often characterised by changes in symptoms without changes in the underlying physiology or pathology of the disease.³⁷

The reports of symptoms and medication use that the participants recorded, concurrent with PEF self-testing, were indications of events that had occurred in the past 2 h. In contrast, the post-test questions about symptoms and medication use required the participants to recall what had occurred weeks earlier, and to determine whether there had been a difference between work days and non-work days. The approach based on concurrent reports had a higher sensitivity for detecting work-exacerbated asthma cases than the approach based on post-test reports (62% vs 15%, respectively).

As clinical evaluation for suspected work-exacerbated asthma is often limited to those who have self-identified based on their perceived symptom or medication use patterns, many people who would probably qualify for compensation are not being identified. This same problem would diminish the number of cases reported by treating doctors to surveillance systems. Although concurrent reports of medication use were more effective at identifying WEA–PEF than post-test reports, this approach still resulted in many false positives and false negatives.

Main messages

- Self-reported asthma symptoms and medication use recorded daily were more sensitive indicators of work-related peak expiratory flow (PEF) patterns than symptoms and medication use reported retrospectively
- The majority of adults with asthma who had work-related self-reported asthma symptoms or medication use did not have similar findings with serial PEF
- A sizeable minority of the adults with asthma, whose serial PEF measurements indicated a work-related pattern, did not report work-related asthma symptoms or medication use

Policy implications

- These findings suggest that self-reported symptoms and medication use fail to identify many people with work-related asthma exacerbations as determined by serial peak expiratory flow measurements

CONCLUSIONS

Serial PEF measurements provided objective evidence for WEA but participation was poor in this cohort of adults with asthma. At the same time, those who completed the serial spirometry testing seemed to be representative of the potential participants, and 14% of them were determined to have WEA–PEF. With WEA–PEF status as the standard for WEA, self-reports of symptoms and medication use were inaccurate, resulting in both false positives and false negatives. If clinical evaluation for suspected WEA is limited to those who have self-identified, then many people with this condition who would potentially qualify for compensation are not being identified. The overall performance of self-reports in identifying work-related patterns of PEF among adults with asthma illuminates the need for further research in the area of development and validation of the questionnaire.

ACKNOWLEDGEMENTS

We thank Nancy Sahakian (NIOSH) for her contribution to the review of the serial PEF records.

Authors' affiliations

Aimee R Bolen, Paul K Henneberger, Xiaoming Liang, Division of Respiratory Disease Studies, National Institute for Occupational Safety and Health, Morgantown, West Virginia, USA

Susan R Sama, Donald K Milton, Harvard School of Public Health, Boston, Massachusetts, USA

Peggy A Preusse, Richard A Rosiello, Fallon Clinic Research Department, Worcester, Massachusetts, USA

Funding: The funding source was the Centers for Disease Control and Prevention.

Competing interests: None.

The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the National Institute for Occupational Safety and Health (NIOSH). Mention of any company or product does not constitute endorsement by NIOSH.

REFERENCES

- 1 **Bardana E**. Occupational asthma. In: Slavov R, Reisman R, eds. *Asthma*. 2nd edn. Philadelphia: Hanley & Belfus, 2002:173–90.

- 2 **Henneberger PK**, Hoffman C, Magid D, et al. Work-related exacerbation of asthma. *Int J Occup Environ Health* 2002;**8**:291–6.
- 3 **Marabini A**, Siracusa A, Stopponi R, et al. Outcome of occupational asthma in patients with continuous exposure. *Chest* 2003;**124**:2372–6.
- 4 **Timmer S**, Rosenman K. Occurrence of occupational asthma. *Chest* 1993;**104**:816–20.
- 5 **Godnic-Cvar J**. How to confirm occupational asthma. *Int Arch Occup Environ Health* 1995;**67**:84.
- 6 **Blanc PD**, Cisternas M, Smith S, et al. Occupational asthma in a community-based survey of adult asthma. *Chest* 1996;**109**:56S–7S.
- 7 **Malo JL**. Assessment of peak expiratory flow in asthma. *Curr Opin Pulm Med* 1996;**2**:75–80.
- 8 **Balmes J**, Becklake M, Blanc P, et al. American Thoracic Society Statement: occupational contribution to the burden of airway disease. *Am J Respir Crit Care Med* 2003;**167**:787–97.
- 9 **Jajosky R**, Harrison R, Reinisch F, et al. Surveillance of work-related asthma in selected U.S. states using surveillance guidelines for state health departments—California, Massachusetts, Michigan, New Jersey, 1993–1995. *Morb Mortal Wkly Rep* 1999;**48**:1–20.
- 10 **Goe SK**, Henneberger PK, Reilly MJ, et al. A descriptive study of work aggravated asthma. *Occup Environ Med* 2004;**61**:512–17.
- 11 **Becklake M**, Malo JL, Chan-Yeung M. Epidemiological approaches in occupational asthma. In: Bernstein IL, Chan-Yeung M, Malo JL, Bernstein DL, eds. *Asthma in the workplace*. 2nd edn. New York: Marcel Dekker, 1999:27–65.
- 12 **Jain P**, Kavuru MS, Emerman CL, et al. Utility of peak expiratory flow monitoring. *Chest* 1998;**114**:861–76.
- 13 **Rubinfeld A**, Pain M. Perception of asthma. *Lancet* 1976;**1**:882–4.
- 14 **Rietveld S**. Symptom perception in asthma: a multidisciplinary view. *J Asthma* 1998;**35**:137–46.
- 15 **Lee HS**. Serial peak expiratory flow rate monitoring—a useful tool in epidemiological studies on occupational asthma. *Ann Acad Med Singapore* 1994;**23**:725–30.
- 16 **Liss GM**, Tarlo SM. Peak expiratory flow rates in possible occupational asthma. *Chest* 1991;**100**:63–9.
- 17 **Malo JL**, Cote J, Cartier A, et al. How many times per day should peak expiratory flow rates be assessed when investigating occupational asthma? *Thorax* 1993;**48**:1211–17.
- 18 **Moscato G**, Godnic-Cvar J, Maestrelli P. Statement on self-monitoring of peak expiratory flows in the investigation of occupational asthma. *Eur Respir J* 1995;**8**:1605–10.
- 19 **Anees W**. Use of pulmonary function tests in the diagnosis of occupational asthma. *Ann Allergy Asthma Immunol* 2003;**90**(Suppl 2):47–51.
- 20 **Anees W**, Gannon PF, Huggins V, et al. Effect of peak expiratory flow data quantity on diagnostic sensitivity and specificity in occupational asthma. *Eur Respir J* 2004;**23**:730–4.
- 21 **Leroyer C**, Perfetti L, Trudeau C, et al. Comparison of serial monitoring of peak expiratory flow and FEV₁ in the diagnosis of occupational asthma. *Am J Respir Crit Care Med* 1998;**158**:827–32.
- 22 **Henneberger PK**, Derk SJ, Sama SR, et al. The frequency of workplace exacerbation among health maintenance organization members with asthma. *Occup Environ Med* 2006;**63**:551–7.
- 23 **American Thoracic Society**. Standardization of spirometry: 1994 update. *Am J Respir Crit Care Med* 1995;**152**:1107–36.
- 24 **Bright P**, Burge PS. Occupational lung disease-8: the diagnosis of occupational asthma from serial measurements of lung function at and away from work. *Thorax* 1996;**51**:857–63.
- 25 **Baldwin DR**, Gannon P, Bright P, et al. Interpretation of occupational peak flow records: level of agreement between expert clinicians and OASYS-2. *Thorax* 2002;**57**:860–4.
- 26 **Burge PS**, Pantin CF, Newton DT, et al. Development of an expert system for the interpretation of serial peak expiratory flow measurements in the diagnosis of occupational asthma. *Occup Environ Med* 1999;**56**:758–64.
- 27 **Pearce N**, Beasley R, Burgess C, et al. *Asthma epidemiology: principles and methods*. New York: Oxford University Press, 1998.
- 28 **Pekkanen J**, Pearce N. Defining asthma in epidemiological studies. *Eur Respir J* 1999;**14**:951–7.
- 29 **US Census Bureau**. 2000 Census of population and housing. <http://www.census.gov/prod/cen2000/doc/sf3.pdf> (Accessed 7 March 2007).
- 30 **Cullen MR**, Redlich CA, Beckett WS, et al. Feasibility study of respiratory questionnaire and peak flow recordings in autobody shop workers exposed to isocyanate-containing spray paint: observations and limitations. *Occup Med* 1996;**46**:197–204.
- 31 **Henneberger PK**, Stanbury MJ, Trimboth LS, et al. The use of portable peak flowmeters in the surveillance of occupational asthma. *Chest* 1991;**100**:1515–21.
- 32 **Malo JL**, Cartier A, Ghezzi H, et al. Compliance with peak expiratory flow readings affect the within- and between-reader reproducibility of interpretation of graphs in subjects investigated for occupational asthma. *J Allergy Clin Immunol* 1996;**98**(part 1):1132–4.
- 33 **Rosenman K**, Reilly MJ, Schill DP, et al. Cleaning products and work-related asthma. *J Occup Environ Med* 2003;**45**:556–63.
- 34 **Pechter E**, Davis L, Tumpowsky C, et al. Work-related asthma among health care workers: surveillance data from California, Massachusetts, Michigan, and New Jersey, 1993–1997. *Am J Ind Med* 2005;**47**:265–75.
- 35 **Massachusetts Department of Public Health**. *Occupational lung disease bulletin*. Boston, MA: MDPH, December, 2005.
- 36 **Baur X**, Huber H, Degens PO, et al. Relation between occupational asthma case history, bronchial methacholine challenge, and specific challenge test in patients with suspected occupational asthma. *Am J Ind Med* 1998;**33**:114–22.
- 37 **Vandenplas O**, Malo J-L. Definitions and types of work-related asthma: a nosological approach. *Eur Respir J* 2003;**21**:706–12.

Keep up to date: sign up for our alerting services

Find out automatically when an article is published on a specific topic or by a particular author. We can also alert you when an article is cited or if an eletter or correction is published. You can also choose to be alerted when a new issue is published online [and when we post articles Online First]. Check out the New Content Alerts and Citation tracker from the Online tools section on the home page.