

Lung Cancer Incidence in Never Smokers

Heather A. Wakelee, Ellen T. Chang, Scarlett L. Gomez, Theresa H. Keegan, Diane Feskanich, Christina A. Clarke, Lars Holmberg, Lee C. Yong, Laurence N. Kolonel, Michael K. Gould, and Dee W. West

ABSTRACT

Purpose

Lung cancer is a leading cause of cancer death worldwide. Although smoking remains the predominant cause of lung cancer, lung cancer in never smokers is an increasingly prominent public health issue. However, data on this topic, particularly lung cancer incidence rates in never smokers, are limited.

Methods

We reviewed the existing literature on lung cancer incidence and mortality rates among never smokers and present new data regarding rates in never smokers from the following large, prospective cohorts: Nurses' Health Study; Health Professionals Follow-Up Study; California Teachers Study; Multiethnic Cohort Study; Swedish Lung Cancer Register in the Uppsala/Örebro region; and First National Health and Nutrition Examination Survey Epidemiologic Follow-Up Study.

Results

Truncated age-adjusted incidence rates of lung cancer among never smokers age 40 to 79 years in these six cohorts ranged from 14.4 to 20.8 per 100,000 person-years in women and 4.8 to 13.7 per 100,000 person-years in men, supporting earlier observations that women are more likely than men to have non-smoking-associated lung cancer. The distinct biology of lung cancer in never smokers is apparent in differential responses to epidermal growth factor receptor inhibitors and an increased prevalence of adenocarcinoma histology in never smokers.

Conclusion

Lung cancer in never smokers is an important public health issue, and further exploration of its incidence patterns, etiology, and biology is needed.

J Clin Oncol 25:472-478. © 2007 by American Society of Clinical Oncology

INTRODUCTION

In the United States, lung cancer incidence and mortality rates have been steadily declining over the past decade, following the well-documented decline in the prevalence of tobacco smoking.¹⁻³

However, in the United States, lung cancer remains the leading cause of cancer death, killing more patients than breast, colon, and prostate cancers combined.⁴ Although tobacco smoke is the predominant risk factor for development of lung cancer, there is a distinct group of patients who develop the disease without a history of tobacco smoking. Clinical observations suggest that the percentage of never smokers among lung cancer patients may be increasing; however, it is unclear whether this apparent trend represents an increase in lung cancer incidence among never smokers or the increasing prevalence of never smokers in the general population. The growing number of never smokers in the United States and other countries underscores the importance of

understanding the epidemiology and biology underlying lung cancer in this population.

Are lung cancer rates among never smokers increasing? Although we can take only the first step towards answering this question, in this article, we review the current knowledge regarding the incidence patterns and biology of non-smoking-associated lung cancer and suggest future research directions to improve understanding of this disease in the growing at-risk population. To this end, we summarize the existing literature on lung cancer incidence rates among never smokers and present new data on rates from selected large, prospective cohorts. Finally, we present evidence suggesting biologic and genetic differences between smoking-related and non-smoking-related lung cancers and posit important new research questions.

In the United States, the most widely used resource for documenting cancer trends in population groups is the National Cancer Institute's Surveillance, Epidemiology, and End Results (SEER) cancer registries.⁵ However, SEER and most other

From the Division of Oncology, Department of Medicine and the Department of Health Research and Policy, Stanford University School of Medicine, Stanford; Northern California Cancer Center, Fremont; Veterans Affairs Palo Alto Health Care System, Palo Alto, CA; Channing Laboratory, Department of Medicine, Brigham and Women's Hospital; Harvard Medical School, Boston, MA; Division of Surveillance, Hazard Evaluations and Field Studies, National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention, Cincinnati, OH; Cancer Epidemiology Program, Cancer Research Center of Hawaii, University of Hawaii, Honolulu, HI; and Regional Oncologic Center, Uppsala University Hospital, Uppsala, Sweden.

Submitted May 4, 2006; accepted September 22, 2006.

Supported in part by Grant No. R01 CA77398 from the National Cancer Institute and Contract No. 97-10500 from the California Cancer Research Fund.

The findings and conclusions in this article are those of the authors and do not necessarily represent the views of the National Institute for Occupational Safety and Health.

Authors' disclosures of potential conflicts of interest and author contributions are found at the end of this article.

Address reprint requests to Heather Wakelee, MD, Division of Medical Oncology, Stanford Clinical Cancer Center, 875 Blake Wilbur Dr, Stanford, CA 94305-5826; e-mail: hwakelee@stanford.edu.

© 2007 by American Society of Clinical Oncology

0732-183X/07/2505-472/\$20.00

DOI: 10.1200/JCO.2006.07.2983

cancer registries do not collect information on patient smoking history, which precludes the examination of cancer incidence patterns by smoking status. Furthermore, information on the prevalence of current and former smokers within subgroups of age, sex, and race/ethnicity in the general population (data necessary for detailed incidence calculations) is difficult to obtain for many populations. SEER data have been linked with population-based tobacco use information,⁶ but this approach allows only for ecologic correlations between the population-level prevalence of smoking and incidence of lung cancer in broad demographic subgroups; individual patient-level data on smoking status are still needed for more specific inference on patterns of smoking-related cancer.⁷

Previous studies reporting incidence rates of lung cancer in never smokers cannot easily be compared because of dissimilar population standards for age adjustment of incidence rates. We present updated and previously unpublished incidence data, age adjusted to a common population, for lung cancer in never smokers from six cohorts. We also provide distributions of lung cancer histology and age at diagnosis to illustrate differences in the characteristics of lung cancer according to smoking status. It is important to emphasize that, although these data add to the sparse body of knowledge on incidence rates of lung cancer in never smokers, they cannot contribute to an understanding of whether rates have changed over time, primarily because cohort recruitment in all of these studies took place over a circumscribed period of time.

METHODS

Using data from six large cohort populations (Table 1), truncated (ie, limited to age 40 to 79 years, rather than all ages),⁸ age-adjusted incidence rates were

calculated based on the 2000 United States standard population, with the proportions of 5-year age groups between 40 and 79 years recalculated to summate to 1.0. The age-adjusted incidence rates, with exact Poisson 95% CIs,¹⁸ were limited to adults age 40 to 79 years to facilitate comparison among the cohorts. We also examined the percentage, with exact binomial 95% CI,¹⁸ of lung cancer patients age 40 to 79 years at diagnosis with adenocarcinoma histology and the median age at diagnosis of current, former, and never smokers in each cohort (Table 2). Histology was ascertained through medical records or linkage to cancer registries for all cohorts for which such information was available, as detailed in Table 1.

Details about the methodology of the Nurses' Health Study (NHS),^{9,10} Health Professionals Follow-Up Study (HPFS),^{9,11} California Teachers Study (CTS),¹² Multiethnic Cohort (MEC) Study,^{13,14} Swedish Uppsala/Örebro Lung Cancer Register (U/OLCR),¹⁵ and First National Health and Nutrition Examination Survey Epidemiologic Follow-Up Study (NHEFS)¹⁶ have been previously reported and are listed in Table 1. These six cohorts were selected because they were prospective, provided prediagnosis data on smoking status, observed cohort members for validated diagnoses of incident cases of lung cancer, enrolled adults age 40 to 79 years, and provided diversity both geographically and demographically.

RESULTS

Table 2 presents the age-adjusted incidence rates of lung cancer in never, former, and current smokers in all six cohorts. These rates per 100,000 person-years for never smokers age 40 to 79 years were 15.2 (female) in the NHS, 11.2 (male) in the HPFS, 20.8 (female) in the CTS, 13.7 (male) and 20.7 (female) in the MEC, 4.8 (male) and 14.4 (female) in the U/OLCR, and 12.7 (male) and 19.3 (female) in the NHEFS. By comparison, age-adjusted rates in current smokers are roughly 12 to 30 times higher. Overall rates were comparable between the US population-based cohorts, including MEC and NHEFS, and

Table 1. Characteristics of the NHS, HPFS, CTS, MEC, U/OLCR, and NHEFS Cohort Members Included in This Analysis

Characteristic	NHS ^{9,10}	HPFS ^{9,11}	CTS ¹²	MEC ^{13,14}	U/OLCR ¹⁵	NHEFS ¹⁶
Dates of follow-up	1976 to 2002	1986 to 2002	1995-1996 to 2002	1993-1996 to 2001	2003	1971-1975 to 1992
Age at baseline, years	30-55	40-75	33-79*	45-75	40-79	25-74
Population at risk, No.						
Male	—	51,529	—	82,460	438,966†	5,075
Female	121,700	—	108,329	101,359	447,603†	7,637
Incident lung cancer patients, 40-79 years at diagnosis, No.						
Male	—	528	—	1,078‡	273	160
Female	1,817	—	393	805‡	250	75
Region	United States	United States	California	California/Hawaii	Uppsala/Örebro, Sweden	United States
Ethnicity	Mostly white	Mostly white	Mostly white	Multiple	Mostly white	Multiple
Follow-up for lung cancer	Biennial questionnaires and medical records, if possible	Biennial questionnaires and medical records, if possible	Linkage to cancer registry	Linkage to cancer registry	Linkage to cancer registry	Questionnaires approximately every 5 years and inpatient records or death certificates§
Smoking data	Biennial questionnaires	Biennial questionnaires	Baseline questionnaire	Baseline questionnaire	Questionnaire for at-risk population; medical records or clinical assessment for patients	Baseline and follow-up questionnaire

Abbreviations: NHS, Nurses' Health Study; HPFS, Health Professionals Follow-Up Study; CTS, California Teachers Study; MEC, Multiethnic Cohort Study; U/OLCR, Swedish Uppsala/Örebro Lung Cancer Register; NHEFS, First National Health and Nutrition Examination Survey Epidemiologic Follow-Up Study.

*Age range of eligible cohort members was restricted to those potentially age 40 to 79 years during follow-up.

†Population at risk based on 2003 census count of males and females age 40 to 79 years in Uppsala/Örebro (Statistics Sweden: www.scb.se)¹⁷; smoking status in population at risk based on a survey of 68,000 randomly selected Uppsala/Örebro residents (64% of whom completed the survey) at the beginning of 2004.

‡Age range of incident lung cancer patients was 45 to 79 years at diagnosis.

§The NHEFS lacked information on tumor histology.

Table 2. No. of Lung Cancer Patients and Lung Cancer AAIR Per 100,000 Person-Years Among Adults Age 40 to 79 Years, Percentage of Patients With Adenocarcinoma Histology, and Median Age at Diagnosis by Smoking Status in the Cohorts

Cohort	Smoking Status		
	Never	Former	Current
NHS, female			
Incident lung cancer patients, No.	168	711	938
Age-truncated AAIR	15.2	76.9	293.3
95% CI	9.1 to 24.5	63.7 to 93.2	266.7 to 322.9
Adenocarcinoma, %	70	50	42
95% CI	62 to 78	46 to 54	39 to 46
Median age at diagnosis, years	64	68	64
HPFS, male			
Incident lung cancer patients, No.	43	312	173
Age-truncated AAIR	11.2	67.6	304.5
95% CI	6.5 to 19.0	56.4 to 82.5	277.4 to 334.7
Adenocarcinoma, %	54	51	30
95% CI	37 to 71	45 to 57	23 to 39
Median age at diagnosis, years	67	71	68
CTS, female			
Incident lung cancer patients, No.	91	179	123
Age-truncated AAIR	20.8	65.6	264.4
95% CI	13.5 to 31.2	54.8 to 79.4	240.0 to 291.7
Adenocarcinoma, %	64	54	34
95% CI	53 to 74	46 to 61	26 to 43
Median age at diagnosis, years	67	70	67
MEC, male*			
Incident lung cancer patients, No.	47	520	511
Age-truncated AAIR	13.7	80.0	314.6
95% CI	9.0 to 21.5	67.1 to 96.1	284.6 to 347.7
Adenocarcinoma, %	53	42	31
95% CI	38 to 68	37 to 46	27 to 36
Median age at diagnosis, years	72	72	69
MEC, female*			
Incident lung cancer patients, No.	142	270	393
Age-truncated AAIR	20.7	65.2	233.7
95% CI	13.5 to 31.1	53.5 to 80.2	208.5 to 261.8
Adenocarcinoma, %	58	46	32
95% CI	49 to 66	40 to 52	28 to 37
Median age at diagnosis, years	72	70	67
U/OLCR, male			
Incident lung cancer patients, No.	10	124	139
Age-truncated AAIR	4.8	59.8	173.7
95% CI	2.2 to 10.6	48.1 to 74.5	153.4 to 197.0
Adenocarcinoma, %	67	36	34
95% CI	30 to 93	28 to 45	26 to 42
Median age at diagnosis, years	64	71	64
U/OLCR, female			
Incident lung cancer patients, No.	37	68	145
Age-truncated AAIR	14.4	51.4	149.4
95% CI	8.2 to 23.6	39.9 to 66.0	129.4 to 172.4
Adenocarcinoma, %	64	46	38
95% CI	46 to 79	33 to 58	30 to 46
Median age at diagnosis, years	67	66	63
NHEFS, male†			
Incident lung cancer patients, No.	4	45	111
Age-truncated AAIR	12.7	141.4	362.7
95% CI	10.2 to 18.2	124.9 to 161.0	334.8 to 393.6
Median age at diagnosis, years	78	72	69

(continued on following page)

Table 2. No. of Lung Cancer Patients and Lung Cancer AAIR Per 100,000 Person-Years Among Adults Age 40 to 79 Years, Percentage of Patients With Adenocarcinoma Histology, and Median Age at Diagnosis by Smoking Status in the Cohorts (continued)

Cohort	Smoking Status		
	Never	Former	Current
NHEFS, female†			
Incident lung cancer patients, No.	15	10	50
Age-truncated AAIR	19.3	69.1	168.8
95% CI	14.2 to 27.5	57.2 to 84.1	146.1 to 194.5
Median age at diagnosis, years	71	67	62

Abbreviations: AAIR, age-adjusted incidence rate for invasive lung cancer (per 100,000 person-years) standardized to US 2000 standard million population between age 40 and 79 years; NHS, Nurses' Health Study; HPFS, Health Professionals Follow-Up Study; CTS, California Teachers Study; MEC, Multiethnic Cohort Study; U/OLCR, Swedish Uppsala/Örebro Lung Cancer Register; NHEFS, First National Health and Nutrition Examination Survey Epidemiologic Follow-Up Study.

*MEC participants and lung cancer patients were restricted to age 45 to 79 years; the incidence rate of lung cancer in those 40 to 44 years old was assumed to be half that among those 45 to 49 years old for calculation of AAIRs.

†The NHEFS lacked information on tumor histology.

the cohorts of highly selected populations, including NHS, HPFS, and CTS, although rates in the U/OLCR were consistently lower than in the US cohorts. In 2002, the age-adjusted (standardized to the 2000 US standard population) incidence rates of lung cancer in all males and females in Sweden overall were 29.9 and 20.0 per 100,000 person-years, respectively,¹⁹ whereas the rates among white males and females in the entire United States were 72.5 and 49.9 per 100,000 person-years, respectively.²⁰

As expected, lung cancer age-adjusted incidence rates in all six cohorts were significantly lower in never smokers than former or current smokers. Adenocarcinoma was more common in never smokers than in former or current smokers in all cohorts for which information about histology was available, although the small number of patients who never smoked resulted in wide CIs (Table 2). Although never smokers were slightly older at lung cancer diagnosis than current smokers in two population-based cohorts (MEC and NHEFS), this difference was not observed in the majority of cohorts evaluated (NHS, HPFS, CTS, and U/OLCR; Table 2).

Among female never smokers, the incidence rates in CTS and MEC were slightly, albeit nonsignificantly, higher than the rate in NHS, which was established nearly two decades earlier than the other two cohorts. Likewise, the incidence rate of non-smoking-associated lung cancer was slightly but nonsignificantly higher among males in MEC than males in the earlier established HPFS, although the discrepancy could be a result of racial/ethnic differences in risk of lung cancer in never smokers. Although a higher relative risk of lung cancer associated with smoking has been demonstrated in racial/ethnic groups that are more prevalent in MEC than in HPFS,¹⁴ data on rates of non-smoking-associated lung cancer by racial/ethnic group are limited.

Evaluating comparable groups of males and females, rates of non-smoking-associated lung cancer were consistently higher among females in the NHS, MEC, U/OLCR, and NHEFS cohorts compared with males in the HPFS, MEC, U/OLCR, and NHEFS cohorts. The higher rate among females suggests sex-based differences in either susceptibility or exposure to risk factors (such as secondhand smoke) for non-smoking-associated lung cancer.

DISCUSSION

Our analysis of recent cohort data finds truncated age-adjusted incidence rates of lung cancer in never smokers ranging from 4.8 to 20.8

per 100,000 person-years in men and women 40 to 79 years old. Because the effects of time cannot be separated from those of aging in these cohorts, we cannot assess secular trends in the incidence rate of non-smoking-associated lung cancer in these cohorts, and we also cannot compare these rates to historical data to evaluate incidence changes over time. However, establishing the current incidence rates, as we have done, is an important step in better understanding this distinct disease subset. To put the problem of lung cancer in never smokers in perspective, the rates we report are similar to age-adjusted rates for myeloma (13.2 per 100,000) in men or cervical (15.4 per 100,000) or thyroid cancer (17.3 per 100,000) in women age 40 to 79 years old diagnosed in the United States between 1998 and 2002.²⁰

Better understanding of the incidence rate and etiology of lung cancer in never smokers is important because of the implications for therapeutic trials and epidemiologic studies of lung cancer. Differences in lung cancer biology between never smokers and smokers are illustrated by findings from several studies. One of the most striking distinctions is the observed differential response to drugs that target the epidermal growth factor receptor (EGFR). Compared with current or former smokers diagnosed with lung cancer, never smoker patients treated with these agents have higher response rates to treatment and better survival.^{21,22} In a randomized phase III trial with the EGFR kinase inhibitor gefitinib in refractory, advanced lung cancer patients, never smokers treated with gefitinib compared with placebo had a reduced risk of death from lung cancer (relative risk for survival analysis, hazard ratio [HR] = 0.67, $P = .012$), whereas the HR of lung cancer death in former/current smokers did not differ between the gefitinib and placebo arms (HR = 0.92, $P = .242$).²¹ In the registration trial (BR.21) for the EGFR kinase inhibitor erlotinib, the overall response rate to erlotinib was 24.7% for never smokers and 3.9% ($P < .001$) for former/current smokers.²² Of all the variables tested, only a history of never smoking was a significant independent predictor of improved survival with erlotinib therapy.²²

The biology underlying the differential response to treatment with EGFR inhibitors is an area of active investigation and helps to illustrate why lung cancer in never smokers may behave differently. Mutations in the *EGFR* are seen more often in tumors from never smokers.²³⁻²⁶ Differences in EGFR protein expression may also contribute to differences in treatment response,²³ with a distinct EGFR pathway immunohistochemical profile seen in never versus current smokers.²⁷ Other analyses have also demonstrated distinct

mutational or expression patterns in *KRAS* TP53, p53, and nitrotyrosine (a marker of nitric oxide protein damage) in tumors of never smokers compared with smokers.^{26,28}

Most reports show a modest survival benefit for non-small-cell lung cancer (NSCLC) patients who are never smokers compared with smokers, regardless of therapy. This was seen for never smokers in the BR.21 trial with erlotinib (HR = 0.8, $P = .048$ compared with current/former smokers regardless of therapy)²² and in a review of 12,000 Southern California NSCLC patients (comparing current/former with never smokers, HR = 1.09, $P = .045$).²⁹ Additionally, a single-institution review of 650 patients with NSCLC found the 5-year overall survival rate to be 16% for current smokers versus 23% for never smokers ($P = .004$).³⁰ Another review of 311 patients with early-stage lung cancer found that the relative risk of death was 0.45 ($P = .042$) comparing never smokers with current smokers.³¹ Finally, among 61 patients with screen-detected lung cancer in Japan, the mean tumor volume doubling time was twice as long in 31 never smokers (607 \pm 392 days) as it was in 30 current smokers (292 \pm 297 days, $P = .001$).³² The implications of these results for epidemiologic studies are clear; improvements in lung cancer survival over time might be a result of an increasing proportion of never smokers among lung cancer patients rather than improved therapies.

Other evidence for a biologic difference in lung cancer between smokers and never smokers comes from differences in histology. Adenocarcinomas seem to be more common in never smokers, light smokers, and former smokers, whereas squamous cell or other histologic types are more common in heavy smokers and current smokers.^{27,33,34} Furthermore, the prevalence of adenocarcinoma among lung cancer patients increases with years since quitting smoking.³⁵ Likewise, our data show a higher proportion of adenocarcinoma among never smokers than among former or current smokers (Table 2).

As our data show, lung cancer rates in never smokers are comparable to the incidence rates of cervical cancer or myeloma in the United States, yet the etiology of this disease is not well understood. Identifying risk factors for lung cancer among never smokers has been an area of active inquiry. Secondhand smoke has been established as a major risk factor among never smokers.³⁶⁻³⁸ Occupational exposures, such as asbestos, chromium, arsenic, and others, also play a role, although more so in smokers.³⁹⁻⁴¹ Domestic radon exposure may also contribute to the risk of lung cancer in never smokers,^{42,43} although some controversy remains,⁴⁴ and arsenic in drinking water has also been implicated.^{45,46} Other factors including indoor pollutants (cooking oil vapors and coal burning),⁴⁷ previous lung disease,⁴⁸⁻⁵⁰ dietary factors,^{51,52} family history,^{37,53,54} and genetic factors may also affect lung cancer development.⁵⁵⁻⁶⁰

Overall, lung cancer incidence rates in the U/OLCR cohort were consistently lower than in the US cohorts. Although the lower prevalence of smoking in Sweden compared with the United States likely contributes to the lower overall incidence rate of lung cancer in Sweden,^{19,20} perhaps in part as a result of lower exposure to secondhand smoke among never smokers, it does not entirely explain our finding of lower rates of lung cancer among never smokers only or among smokers only. Instead, the discrepancies between the countries even within strata of smoking status suggest differences in smoking patterns among smokers and in the prevalence of environmental or genetic cofactors for lung cancer among both smokers and never smokers. The cumulative risk of lung cancer among Swedish male smokers is also considerably lower than that among men from other European countries.⁶¹

The biologic differences in lung cancer in never smokers versus current/former smokers are apparent primarily in differential response to specific therapies (most notably EGFR inhibitors) and in distribution of histology (increased adenocarcinoma in never smokers), as supported by our data. Our data also support the observation that women are more likely than men to have non-smoking-associated lung cancer, which is in contrast to the finding that men had a higher mortality rate from non-smoking-associated lung cancer than women in the American Cancer Society Cancer Prevention Study cohorts.⁶² This discrepancy could be, in part, a result of better survival among women than men with non-smoking-associated lung cancer, although data on this subject are lacking. The literature does support a survival benefit for women versus men with lung cancer overall.⁶³ Clearly, more research is needed regarding the intriguing etiology, prognosis, treatment, and outcomes of non-smoking-associated lung cancer.

Despite the emergence of clinical and epidemiologic studies focused on identifying biologic and genetic differences between smoking- and non-smoking-associated lung cancers and risk factors for non-smoking-associated lung cancer, it remains uncertain whether the incidence of lung cancer in never smokers is increasing. Evaluation of secular trends is possible in longitudinal studies with open enrollment over a long span of time, such that incidence changes caused by aging are distinguishable from secular changes. To our knowledge, there have been only two published examples of data of this type. The first is a linkage of the nationwide Swedish construction workers' health care program to the national cancer registry, which documented an increase in the age-adjusted (standardized to the 2000 World Standard Population) incidence rate of non-smoking-related lung cancer between 1976 to 1980 (1.5 per 100,000) and 1991 to 1995 (5.4 per 100,000).⁶⁴ The second is a comparison of two American Cancer Society Cancer Prevention Study cohorts, in which the age-adjusted (standardized to the combined age distribution of the two cohorts and, therefore, not directly comparable to the Swedish construction workers' study) mortality rate of non-smoking-associated lung cancer in women increased slightly, but statistically significantly, from 12.3 per 100,000 in 1959 to 1972 to 14.7 per 100,000 in 1982 to 2000, with most of the increase occurring among women age 70 to 84 years.⁶² The mortality rate among men, however, did not change over time.

In a comparison of two large, hospital- and community-based case-control studies conducted in the United Kingdom in 1950 and 1990, the percentage of never smokers among the male lung cancer patients was 0.5% in both studies, whereas the percentage of never smokers among male controls increased from 4.5% to 19.0%.⁶⁵ Among women, the percentage of lung cancer patients who never smoked was 37.0% in 1950 and 7.6% in 1990, whereas the percentage of never-smokers among female controls decreased less dramatically from 54.6% to 50.3%. These results suggest that the proportion of lung cancer patients who never smoked does not necessarily reflect population-level changes in smoking prevalence.

In the United States, a study of 100 NSCLC patients seen at a single institution in the late 1980s determined smoking status through questionnaires and medical record review and found that 11% of patients were never smokers.³³ In a large case series of 11,969 NSCLC patients from three Southern California counties (1995 to 2003), investigators estimated that 9.7% of patients were never smokers.²⁹ In this study, a nonsignificant increase in the prevalence of never smokers with NSCLC was noted in 1999 to 2003 versus 1995 to 1999.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

The authors indicated no potential conflicts of interest.

AUTHOR CONTRIBUTIONS

Conception and design: Heather A. Wakelee, Ellen T. Chang, Scarlett L. Gomez, Theresa H.M. Keegan, Christina A. Clarke, Michael K. Gould, Dee W. West

Provision of study materials or patients: Ellen T. Chang, Diane Feskanich, Christina A. Clarke, Lars Holmberg, Lee C. Yong, Laurence N. Kolonel, Dee W. West

Collection and assembly of data: Heather A. Wakelee, Ellen T. Chang, Diane Feskanich, Christina A. Clarke, Lars Holmberg, Lee C. Yong, Laurence N. Kolonel

Data analysis and interpretation: Heather A. Wakelee, Ellen T. Chang, Scarlett L. Gomez, Theresa H.M. Keegan, Diane Feskanich, Christina A. Clarke, Lars Holmberg, Lee C. Yong, Laurence N. Kolonel, Michael K. Gould, Dee W. West

Manuscript writing: Heather A. Wakelee, Ellen T. Chang, Scarlett L. Gomez, Theresa H.M. Keegan

Final approval of manuscript: Heather A. Wakelee, Ellen T. Chang, Scarlett L. Gomez, Theresa H.M. Keegan, Diane Feskanich, Christina A. Clarke, Lars Holmberg, Lee C. Yong, Laurence N. Kolonel, Michael K. Gould, Dee W. West

REFERENCES

- Giovino GA, Schooley MW, Zhu BP, et al: Surveillance for selected tobacco-use behaviors—United States, 1900-1994. *MMWR CDC Surveill Summ* 43:1-43, 1994
- Jemal A, Chu KC, Tarone RE: Recent trends in lung cancer mortality in the United States. *J Natl Cancer Inst* 93:277-283, 2001
- Jemal A, Ward E, Thun MJ: Contemporary lung cancer trends among U.S. women. *Cancer Epidemiol Biomarkers Prev* 14:582-585, 2005
- Jemal A, Siegel R, Ward E, et al: Cancer statistics, 2006. *CA Cancer J Clin* 56:106-130, 2006
- Glaser SL, Clarke CA, Gomez SL, et al: Cancer surveillance research: A vital subdiscipline of cancer epidemiology. *Cancer Causes Control* 16:1009-1019, 2005
- Wingo PA, Ries LA, Giovino GA, et al: Annual report to the nation on the status of cancer, 1973-1996, with a special section on lung cancer and tobacco smoking. *J Natl Cancer Inst* 91:675-690, 1999
- Morgenstern H: Ecologic studies in epidemiology: Concepts, principles, and methods. *Annu Rev Public Health* 16:61-81, 1995
- Gondos A, Parkin DM, Chokunonga E, et al: Calculating age-adjusted cancer survival estimates when age-specific data are sparse: An empirical evaluation of various methods. *Br J Cancer* 94:450-454, 2006
- Bain C, Feskanich D, Speizer FE, et al: Lung cancer rates in men and women with comparable histories of smoking. *J Natl Cancer Inst* 96:826-834, 2004
- Colditz GA, Manson JE, Hankinson SE: The Nurses' Health Study: 20-year contribution to the understanding of health among women. *J Womens Health* 6:49-62, 1997
- Rimm EB, Stampfer MJ, Colditz GA, et al: Effectiveness of various mailing strategies among nonrespondents in a prospective cohort study. *Am J Epidemiol* 131:1068-1071, 1990
- Bernstein L, Allen M, Anton-Culver H, et al: High breast cancer incidence rates among California teachers: Results from the California Teachers Study (United States). *Cancer Causes Control* 13:625-635, 2002
- Kolonel LN, Henderson BE, Hankin JH, et al: A multiethnic cohort in Hawaii and Los Angeles: Baseline characteristics. *Am J Epidemiol* 151:346-357, 2000
- Haiman CA, Stram DO, Wilkens LR, et al: Ethnic and racial differences in the smoking-related risk of lung cancer. *N Engl J Med* 354:333-342, 2006
- Lamberg K, Wagenius G, Garma H, et al: Lung cancer in the Uppsala/Orebro region: Register for lung cancer: Account of materials for 1995-2003. Uppsala, Sweden, Regional Oncological Center, 2004
- Yong LC, Brown CC, Schatzkin A, et al: Intake of vitamins E, C, and A and risk of lung cancer: The NHANES I epidemiologic followup study. *Am J Epidemiol* 146:231-243, 1997
- Statistics Sweden: Statistical database: Population. <http://www.scb.se>
- Daly L: Simple SAS macros for the calculation of exact binomial and Poisson confidence limits. *Comput Biol Med* 22:351-361, 1992
- Swedish Cancer Registry: Cancer Incidence in Sweden 2002. Stockholm, Sweden, National of Health and Welfare, Centre for Epidemiology, 2003
- Surveillance, Epidemiology, and End Results (SEER) Program Seer*Stat Database: Incidence - SEER 13 Regs Public-Use, Nov 2004 Sub for Expanded Races (1992-2002). Bethesda, MD, National Cancer Institute, Division of Cancer Control and Population Sciences, Surveillance Research Program, Cancer Statistics Branch, released April 2005, based on the November 2004 submission
- Thatcher N, Chang A, Parikh P, et al: Gefitinib plus best supportive care in previously treated patients with refractory advanced non-small-cell lung cancer: Results from a randomised, placebo-controlled, multicentre study (Iressa Survival Evaluation in Lung Cancer). *Lancet* 366:1527-1537, 2005
- Shepherd FA, Rodrigues Pereira J, Ciuleanu T, et al: Erlotinib in previously treated non-small-cell lung cancer. *N Engl J Med* 353:123-132, 2005
- Tsao MS, Sakurada A, Cutz JC, et al: Erlotinib in lung cancer: Molecular and clinical predictors of outcome. *N Engl J Med* 353:133-144, 2005
- Pao W, Miller V, Zakowski M, et al: EGF receptor gene mutations are common in lung cancers from "never smokers" and are associated with sensitivity of tumors to gefitinib and erlotinib. *Proc Natl Acad Sci U S A* 101:13306-13311, 2004
- Pham D, Kris MG, Riely GJ, et al: Use of cigarette-smoking history to estimate the likelihood of mutations in epidermal growth factor receptor gene exons 19 and 21 in lung adenocarcinomas. *J Clin Oncol* 24:1700-1704, 2006
- Tam IY, Chung LP, Suen WS, et al: Distinct epidermal growth factor receptor and KRAS mutation patterns in non-small cell lung cancer patients with different tobacco exposure and clinicopathologic features. *Clin Cancer Res* 12:1647-1653, 2006
- Dutu T, Michiels S, Fouret P, et al: Differential expression of biomarkers in lung adenocarcinoma: A comparative study between smokers and never-smokers. *Ann Oncol* 16:1906-1914, 2005
- Le Calvez F, Mukeria A, Hunt JD, et al: TP53 and KRAS mutation load and types in lung cancers in relation to tobacco smoke: Distinct patterns in never, former, and current smokers. *Cancer Res* 65:5076-5083, 2005
- Zell JA, Ou SH, Ziogas A, et al: Epidemiology of bronchioloalveolar carcinoma: Improvement in

survival after release of the 1999 WHO classification of lung tumors. *J Clin Oncol* 23:8396-8405, 2005

30. Nordquist LT, Simon GR, Cantor A, et al: Improved survival in never-smokers vs current smokers with primary adenocarcinoma of the lung. *Chest* 126:347-351, 2004

31. Sardari Nia P, Weyler J, Colpaert C, et al: Prognostic value of smoking status in operated non-small cell lung cancer. *Lung Cancer* 47:351-359, 2005

32. Hasegawa M, Sone S, Takashima S, et al: Growth rate of small lung cancers detected on mass CT screening. *Br J Radiol* 73:1252-1259, 2000

33. Sridhar KS, Raub WA Jr: Present and past smoking history and other predisposing factors in 100 lung cancer patients. *Chest* 101:19-25, 1992

34. Brownson RC, Loy TS, Ingram E, et al: Lung cancer in nonsmoking women: Histology and survival patterns. *Cancer* 75:29-33, 1995

35. Muscat JE, Wynder EL: Lung cancer pathology in smokers, ex-smokers and never smokers. *Cancer Lett* 88:1-5, 1995

36. Brennan P, Buffler PA, Reynolds P, et al: Secondhand smoke exposure in adulthood and risk of lung cancer among never smokers: A pooled analysis of two large studies. *Int J Cancer* 109:125-131, 2004

37. Gorlova OY, Zhang Y, Schabath MB, et al: Never smokers and lung cancer risk: A case-control study of epidemiological factors. *Int J Cancer* 118:1798-1804, 2006

38. Vineis P, Airolidi L, Veglia P, et al: Environmental tobacco smoke and risk of respiratory cancer and chronic obstructive pulmonary disease in former smokers and never smokers in the EPIC prospective study. *BMJ* 330:277, 2005

39. Alberg AJ, Brock MV, Samet JM: Epidemiology of lung cancer: Looking to the future. *J Clin Oncol* 23:3175-3185, 2005

40. Gottschall EB: Occupational and environmental thoracic malignancies. *J Thorac Imaging* 17:189-197, 2002

41. Neuberger JS, Field RW: Occupation and lung cancer in nonsmokers. *Rev Environ Health* 18:251-267, 2003

42. Krewski D, Lubin JH, Zielinski JM, et al: A combined analysis of North American case-control studies of residential radon and lung cancer. *J Toxicol Environ Health A* 69:533-597, 2006

43. Darby S, Hill D, Auvinen A, et al: Radon in homes and risk of lung cancer: Collaborative analysis of individual data from 13 European case-control studies. *BMJ* 330:223, 2005

44. Sandler DP, Weinberg CR, Shore DL, et al: Indoor radon and lung cancer risk in Connecticut and Utah. *J Toxicol Environ Health A* 69:633-654, 2006

45. Chen CL, Hsu LI, Chiou HY, et al: Ingested arsenic, cigarette smoking, and lung cancer risk: A

follow-up study in arseniasis-endemic areas in Taiwan. *JAMA* 292:2984-2990, 2004

46. Ferreccio C, Gonzalez C, Milosavljevic V, et al: Lung cancer and arsenic concentrations in drinking water in Chile. *Epidemiology* 11:673-679, 2000

47. Boffetta P, Nyberg F: Contribution of environmental factors to cancer risk. *Br Med Bull* 68:71-94, 2003

48. Littman AJ, Thornquist MD, White E, et al: Prior lung disease and risk of lung cancer in a large prospective study. *Cancer Causes Control* 15:819-827, 2004

49. Mayne ST, Buenconsejo J, Janerich DT: Previous lung disease and risk of lung cancer among men and women nonsmokers. *Am J Epidemiol* 149:13-20, 1999

50. Wu AH, Fontham ET, Reynolds P, et al: Previous lung disease and risk of lung cancer among lifetime nonsmoking women in the United States. *Am J Epidemiol* 141:1023-1032, 1995

51. Wakai K, Ando M, Ozasa K, et al: Updated information on risk factors for lung cancer: Findings from the JACC Study. *J Epidemiol* 15:S134-S139, 2005 (suppl 2)

52. Feskanich D, Ziegler RG, Michaud DS, et al: Prospective study of fruit and vegetable consumption and risk of lung cancer among men and women. *J Natl Cancer Inst* 92:1812-1823, 2000

53. Wu AH, Fontham ET, Reynolds P, et al: Family history of cancer and risk of lung cancer among lifetime nonsmoking women in the United States. *Am J Epidemiol* 143:535-542, 1996

54. Brownson RC, Alavanja MC, Caporaso N, et al: Family history of cancer and risk of lung cancer in lifetime non-smokers and long-term ex-smokers. *Int J Epidemiol* 26:256-263, 1997

55. Raimondi S, Boffetta P, Anttila S, et al: Metabolic gene polymorphisms and lung cancer risk in non-smokers: An update of the GSEC study. *Mutat Res* 592:45-57, 2005

56. Wenzlaff AS, Cote ML, Bock CH, et al: CYP1A1 and CYP1B1 polymorphisms and risk of lung cancer among never smokers: A population-based study. *Carcinogenesis* 26:2207-2212, 2005

57. Taioli E, Gaspari L, Benhamou S, et al: Polymorphisms in CYP1A1, GSTM1, GSTT1 and lung cancer below the age of 45 years. *Int J Epidemiol* 32:60-63, 2003

58. Bonner MR, Bennett WP, Xiong W, et al: Radon, secondhand smoke, glutathione-S-transferase M1 and lung cancer among women. *Int J Cancer* 119:1462-1467, 2006

59. Wenzlaff AS, Cote ML, Bock CH, et al: GSTM1, GSTT1 and GSTP1 polymorphisms, environmental tobacco smoke exposure and risk of lung cancer among never smokers: A population-based study. *Carcinogenesis* 26:395-401, 2005

60. Jung CY, Choi JE, Park JM, et al: Polymorphisms in the hMSH2 gene and the risk of primary

lung cancer. *Cancer Epidemiol Biomarkers Prev* 15:762-768, 2006

61. Crispo A, Brennan P, Jockel KH, et al: The cumulative risk of lung cancer among current, ex- and never-smokers in European men. *Br J Cancer* 91:1280-1286, 2004

62. Thun MJ, Henley SJ, Burns D, et al: Lung cancer death rates in lifelong nonsmokers. *J Natl Cancer Inst* 98:691-699, 2006

63. Donington JS, Le QT, Wakelee HA: Lung cancer in women: Exploring sex differences in susceptibility, biology, and therapeutic response. *Clin Lung Cancer* 8:22-29, 2006

64. Boffetta P, Jarvholm B, Brennan P, et al: Incidence of lung cancer in a large cohort of non-smoking men from Sweden. *Int J Cancer* 94:591-593, 2001

65. Peto R, Darby S, Deo H, et al: Smoking, smoking cessation, and lung cancer in the UK since 1950: Combination of national statistics with two case-control studies. *BMJ* 321:323-329, 2000

66. Nyberg F, Agudo A, Boffetta P, et al: A European validation study of smoking and environmental tobacco smoke exposure in nonsmoking lung cancer cases and controls. *Cancer Causes Control* 9:173-182, 1998

67. Wells AJ, English PB, Posner SF, et al: Misclassification rates for current smokers misclassified as nonsmokers. *Am J Public Health* 88:1503-1509, 1998

68. Caraballo RS, Giovino GA, Pechacek TF, et al: Factors associated with discrepancies between self-reports on cigarette smoking and measured serum cotinine levels among persons aged 17 years or older: Third National Health and Nutrition Examination Survey, 1988-1994. *Am J Epidemiol* 153:807-814, 2001

69. Klebanoff MA, Levine RJ, Clemens JD, et al: Serum cotinine concentration and self-reported smoking during pregnancy. *Am J Epidemiol* 148:259-262, 1998

70. Cope GF, Battersby N: Smoking verification and the risk of myocardial infarction. *J Epidemiol Community Health* 58:156, 2004

71. University of California, Los Angeles Center for Health Policy Research CDoHS, Public Health Institute: California Health Interview Survey (CHIS). www.chis.ucla.edu

72. Ciotti M, Giuliani L, Ambrogi V, et al: Detection and expression of human papillomavirus oncogenes in non-small cell lung cancer. *Oncol Rep* 16:183-189, 2006

73. Chen YC, Chen JH, Richard K, et al: Lung adenocarcinoma and human papillomavirus infection. *Cancer* 101:1428-1436, 2004

74. Mornex JF, Thivolet F, De las Heras M, et al: Pathology of human bronchioloalveolar carcinoma and its relationship to the ovine disease. *Curr Top Microbiol Immunol* 275:225-248, 2003

Acknowledgment

We thank Alison Canchola, Inna Feldman, Lisa Kahle, Sarah Marshall, Philip Taylor, and Lynne Wilkens for their assistance with data from the various cohorts.