

7, 28 and 56 days post-exposure. In bronchoalveolar lavage (BAL) studies, polymorphonuclear leukocytes (PMNs) were assessed to index pulmonary inflammation, BAL fluid lactate dehydrogenase (LDH) activity was measured as a marker of cytotoxicity, and BAL fluid albumin was determined as a marker of the lung air-blood barrier integrity. MWCNT exposure induced dose- and time-dependent changes, with maximum changes occurring at 7 days post-exposure for all three BAL markers. At 7 days post-exposure, mice exposed to 40 µg/mouse MWCNT had increased BAL PMNs (724-fold), BAL fluid LDH activity (2.6-fold) and BAL fluid albumin (2.4-fold) versus vehicle-exposed controls. At 56 days post-exposure, mice exposed to 40 µg/mouse MWCNT still had increased BAL PMNs (22.3-fold), BAL fluid LDH activity (1.9-fold) and BAL fluid albumin (1.6-fold) versus vehicle-exposed controls. Thus, relative to 7 days post-exposure, these BAL markers had decreased, but were still significantly elevated above vehicle-exposed controls at 56 days post-exposure. At 28 and 56 days post-exposure, histopathology confirmed persistent interstitial inflammation and indicated fibrosis. In summary, these data indicate that exposure to MWCNT results in dose- and time-dependent changes in pulmonary inflammation and damage, suggesting that MWCNT may pose an occupational health hazard.

PS 2197 NEUROINFLAMMATION AND BLOOD-BRAIN BARRIER CHANGES FOLLOWING EXPOSURE TO ENGINEERED NANOMATERIALS.

K. Sriram, D. W. Porter, A. M. Jefferson, G. X. Lin, M. G. Wolfarth, B. T. Chen, W. McKinney, D. G. Frazer and V. Castranova. *Centers for Disease Control & Prevention (CDC)-NIOSH, Morgantown, WV.*

The unique physico-chemical properties of engineered nanomaterials influence their ability to aerosolize, and thus inhalation exposure is of major occupational concern. Inhaled nanoparticles can potentially translocate to the brain via olfactory sensory neurons or through systemic circulation and cause irreversible damage to the nervous system. To determine if engineered nanomaterials pose a neurological risk, we evaluated the effects of multi-walled carbon nanotubes (MWCNT) in a murine model. Male C57BL/6J mice were exposed to MWCNT either by pharyngeal aspiration (single dose of 10-80µg/mouse; 1, 7, 28, or 56d post-exposure) or whole-body inhalation (10mg/m³ x 5h/d x 2, 4, 8 or 12d; 1d post-exposure). MWCNT exposure elicited neuroinflammation, altered blood-brain barrier (BBB) integrity and induced cellular stress in discrete brain areas. Specifically, MWCNT induced (2 to 16-fold) the mRNA expression of several proinflammatory chemokines (Ccl2, Ccl3, Ccl4, Cxcl2), cytokines (Il-1β, Il-6, Tnfα), selectins (Sele, Selp) and markers of cellular stress (Hspb2, Mt1, Mt2). Exposure to MWCNT also decreased the expression of BBB-related markers (Edn2, Vegfa), suggestive of alterations in BBB integrity. In the hippocampus, MWCNT altered the expression of certain Alzheimer's-related genes (Aplp2, Apha2, Apha3, Bace1, Bace2, Ctsc, Cttd), which interact with or are involved in the processing of amyloid-precursor protein. The neurotoxic responses were comparable between the two routes of exposure and some of the effects persisted until 56d post-exposure. Taken together, our findings suggest that exposure to an engineered nanomaterial like MWCNT, can elicit neuroinflammation, disrupt BBB integrity and cause cellular/molecular changes that could potentially culminate in neurodegeneration.

PS 2198 NANOPARTICLE DISPERSION METHOD USING NATURAL LUNG SURFACTANT.

L. Wang^{1,2}, V. Castranova^{1,2}, R. R. Mercer^{1,2}, N. Wu², T. Li³, S. Li³, J. Hall¹, M. Li² and Y. Rojanasakul². ¹National Institute for Occupational Safety and Health (NIOSH), Morgantown, WV, ²West Virginia University, Morgantown, WV and ³National Institute of Occupational Health and Poison Control, CDC, Beijing, China.

Nanomaterials, as a class of small-scale (<100 nm) substances with unique mechanical, optical and electrical properties, are increasingly being used in a wide range of industries. Their unique properties present new challenges to understanding the toxicity of these materials to humans and the environment. Lung is the major target organ for airborne nanoparticles. *In vitro* and *in vivo* exposure studies often rely on the use of suspended nanoparticle preparations. However, nanoparticles suspended in culture medium or physiologic saline solution tend to form micrometer-sized aggregates. Increasing evidence indicates that the degree of dispersion of nanoparticles has a strong influence on their biological activities. In this study, we test a new method of nanoparticle dispersion using natural lung surfactant, Survanta[®], as a dispersing agent. Dose dependence studies of Survanta[®] were performed on single-walled carbon nanotube (SWCNT) dispersion. Our results show that Survanta[®] at a concentration of 150 µg/ml, which is comparable to that found in normal rodent lungs, was optimal in dispersing SWCNT (0.1 mg/ml), producing well dispersed preparations as analyzed by microscopic and light scattering methods. This dose of Survanta[®] was found to be non-toxic and non-inflammatory *in vivo* and *in vitro* when used alone, and did not mask the bioactivity of SWCNT.

We also found that the dispersed form of SWCNT was more effective in inducing cytotoxicity and lung fibrosis than the non-dispersed form, indicating the importance of nanoparticle dispersion on biological activities. Since Survanta[®] is commercially available and its one step nanoparticle dispersion is simple and rapid, this method provides major advantages over existing methods of nanoparticle dispersion. Furthermore, our stability studies showed that Survanta[®]-dispersed nanoparticles remain well dispersed for months and upon dilution with aqueous medium.

PS 2199 BIODEGRADATION OF SINGLE WALLED CARBON NANOTUBES THROUGH PEROXIDASE CATALYSIS.

N. V. Konduru¹, I. I. Vlasova¹, B. L. Allen², E. R. Kisin³, A. R. Murray³, A. A. Kapralov¹, A. Star², A. A. Shvedova³ and V. E. Kagan¹. ¹Environmental and Occupational Health, University of Pittsburgh, Pittsburgh, PA, ²Department of Chemistry, University of Pittsburgh, Pittsburgh, PA and ³Pathology/Physiology Research Branch, Health Effects Laboratory Division (HELD), NIOSH, Morgantown, WV.

Management of exposure to bioengineered single walled carbon nanotubes (SWCNT) is becoming a major environmental and health challenge. Because of unique characteristics, SWCNT have found increased applications in various fields of science and technology. With increased risk of human and environmental exposure, strategies to degrade SWCNT are also garnering interest. Here, we show - for the first time - that two peroxidases, namely myeloperoxidase and horseradish peroxidase are effective in catalyzing the biodegradation of SWCNT. A systematic characterization of resulting products of degradation was performed using transmission electron microscopy (TEM), dynamic light scattering (DLS), gel electrophoresis, mass spectrometry, UV-Vis-NIR spectroscopy and thermogravimetric analysis (TGA). We also demonstrate that unlike naïve non-biodegraded nanoparticles, the peroxidase degraded particles did not elicit pulmonary inflammatory response in mice as evidenced by the release of pro-inflammatory cytokines and the content of neutrophils (PMN) in bronchoalveolar lavage. These results mark a novel approach to employ peroxidase catalysis for directed biodegradation of carbon nanotubes in biofluids/tissues as well as in environmental settings. Supported by NIOSH OH008282, NORA 927000Y, National Heart, Lung and Blood Institute Grant HL-70755, The Swedish Research Council, the Swedish Council for Working Life and Social Research, The Human Frontier Science Program (HFSP) and the 7th Framework Program of the European Commission.

PS 2200 PULMONARY AND SYSTEMIC INHALATION TOXICITY OF MULTI-WALLED AND SINGLE WALLED CARBON NANOTUBES.

J. D. McDonald¹, L. Mitchell^{1,2}, S. Burchiel² and A. Gigliotti¹. ¹Lovelace Respiratory Research Institute, Albuquerque, NM and ²University of New Mexico, Albuquerque, NM.

Inhalation of multiwalled carbon nanotubes (MWCNTs) and single walled carbon nanotubes (SWCNT) at particle concentrations up to 1 mg/m³ did not result in significant lung inflammation or tissue damage, but caused systemic immune function alterations. C57BL/6 adult (10-12 week) male mice were exposed by whole-body inhalation to control air or 0.3 or 1 mg/m³ respirable aggregates of MWCNTs or SWCNTs for 14 days, with either immediate sacrifice or sacrifice of a recovery group 30 days after the end of exposure. Histopathology of lungs from exposed animals showed alveolar macrophages containing significant amounts of black particles; however, there was minimal to no inflammation or tissue damage observed. Bronchial alveolar lavage fluid also demonstrated particle-laden macrophages; however, white blood cell counts were not increased compared to controls. Both types of carbon nanotubes caused systemic immunosuppression after 14 days and after recovery. Immunosuppression was characterized by reduced T-cell-dependent antibody response to sheep erythrocytes as well as T-cell proliferative ability in presence of mitogen, Concanavalin A (Con A).

PS 2201 SINGLE-WALLED CARBON NANOTUBES: SKIN EXPOSURES.

A. R. Murray^{1,2}, E. Kisin¹, S. S. Leonard¹, S. H. Young¹, C. Kommineni¹, V. E. Kagan³, V. Castranova^{1,2} and A. A. Shvedova^{1,2}. ¹PPRB, NIOSH, Morgantown, WV, ²Department of Physiology/Pharmacology, WVU, Morgantown, WV and ³Department of Environmental and Occupational Health, University of Pittsburgh, Pittsburgh, PA.

Nanotechnology is a newly developing field resulting in the development of unique materials with a variety of applications from electronics to engineered tissue. SWCNT are of the most interest because of their unique mechanical and electrical

properties which result from their fibrous shape and small size. The most common technique for manufacturing SWCNT relies on the use of iron as a transition metal catalyst and can result in the presence of up to 30% metal catalyst in raw SWCNT. The low density of SWCNT may lead to inhalation as well as deposition on exposed skin, which provide routes of exposure that are important to consider when evaluating toxicity. We hypothesized that SWCNT are toxic to the skin, and this toxicity is dependent on the ability of SWCNT to interact with the skin and initiate oxidative stress, and the induction of transcription factors leading to inflammation. To test this hypothesis, the effects of SWCNT were assessed both *in vitro* and *in vivo*. Engineered skin exposed to SWCNT showed increased epidermal thickness and accumulation and activation of dermal fibroblasts, which resulted in increased collagen as well as release of pro-inflammatory cytokines. Exposure of JB6 cells to unpurified SWCNT resulted in the production hydroxyl radicals as detected by ESR and caused a significant dose-dependent activation of AP-1 and NFκB, while partially-purified SWCNT activated only NFκB. Topical exposure of SKH-1 mice (5 days, 40, 80, 160 μg/mouse/day) to SWCNT caused oxidative stress, depletion of GSH, oxidation of protein thiols and carbonyls, elevated MPO activity, and an increase of dermal cell numbers resulting in skin thickening. Altogether, these data indicated that topical exposure to unpurified SWCNT induced free radical generation, oxidative stress, and inflammation, leading to dermal toxicity. Acknowledgements: supported by NIOSH OH008282, NIH HL70755, NORA 927000Y, 927Z1LU and EC-FP-7-NANOMMUNE-214281

PS 2202 PULMONARY EFFECTS OF SINGLE-WALLED CARBON NANOTUBES: INHALATION VS ASPIRATION.

E. Kisin¹, A. R. Murray¹, A. F. Hubbs¹, R. R. Mercer¹, P. Keohavong³, N. Sussman³, B. T. Chen¹, G. Deye², V. Castranova¹, P. A. Baron², V. E. Kagan³ and A. A. Shvedova¹. ¹PPRB, NIOSH, Morgantown, WV, ²DART, NIOSH, Cincinnati, OH and ³Department of Environmental and Occupational Health, University of Pittsburgh, Pittsburgh, PA.

Health effects and occupational risk of exposures associated with manufacturing and application of nanoparticles are critical points for the safe and sustainable development of nanotechnology. The toxic effects of nanoscale materials have not been fully characterized and the limited *in vivo* studies indicate the urgent necessity for further toxicological assessments of nanomaterials. Some argue that pharyngeal aspiration – a single exposure to a bolus of SWCNT - is an artificial exposure where the single large dose contributes to the pulmonary response. Moreover, aspiration studies reported thus far have been relatively high dose exposures, which may not be relevant to chronic lower dose seen in occupational settings. Inhalation of SWCNT more closely mimics occupational and environmental venues than the above mentioned administrations providing more dispersed SWCNT structures while bolus effects are avoided. By applying a new technique to aerosolize SWCNT, we obtained stable and uniform SWCNT dispersions with a concentration of 5 mg/m³ and a count mode aerodynamic diameter of 240 nm for the inhalation experiments. In the current study, we utilized non-purified SWCNT containing up to 17.7% of iron for both inhalation (5 mg/m³, 5 hrs/day for 4 days) and aspiration (varying doses of 5-20 μg/mouse) exposures. Pathological events in both exposure routes were realized through qualitatively similar synergized interactions of early inflammatory response and oxidative stress culminating in the development of multifocal granulomatous pneumonia and interstitial fibrosis. Quantitatively, SWCNT inhalation was more effective than aspiration in causing inflammatory response, oxidative stress, collagen deposition and fibrosis as well as mutations of K-ras gene locus in the lung of C57BL/6 mice. Acknowledgements: supported by NIOSH OH008282, NIH HL70755, NORA 927000Y, and EC-FP-7-NANOMMUNE-214281.

PS 2203 SINGLE-WALLED CARBON NANOTUBES INDUCE PULMONARY AND VASCULAR RESPONSE FOLLOWING INTRATRACHEAL INSTILLATION.

M. C. Schladweiler¹, J. H. Shannahan², R. F. Thomas¹, R. Saxena¹, M. Gilmour¹ and U. P. Kodavanti¹. ¹NHEERL/ETD/PTB, U.S. EPA, Durham, NC and ²Curriculum in Toxicology, UNC-Chapel Hill, Chapel Hill, NC.

Carbon-based nanotubes have been shown to induce varying degrees of pulmonary response in rodents influenced by the dose, the extent of agglomeration, and the functional properties. We hypothesized that low concentrations of non-modified or acid-functionalized (AF) single walled carbon nanotubes (SWCNT) will cause distinct pulmonary and aortic effects on markers of inflammation, coagulation, vasoconstriction following pulmonary exposure. Male Wistar Kyoto rats (12 week old) were intratracheally instilled with pre-characterized freshly sonicated suspensions of SWCNT or AF-SWCNT at 0, 100 or 500 μg/kg in saline. Pulmonary injury and inflammatory effects were small as determined by bronchoalveolar lavage fluid (BALF) analysis and were concentration-dependent. Small increases in BALF protein were noted with high dose of both types of tubes while albumin increased only

with AF-SWCNT. Surprisingly, small but significant increases in BALF γ-glutamyl transferase activity (marker of cell membrane damage) were noted only with high concentrations of SWCNT but not AF-SWCNT. LDH activity was increased 4-d at high concentration of both types of SWCNT. Both types of nanotubes moderately increased BALF neutrophils at high concentration while no increase in macrophages occurred (1-d-4-d). Real-time PCR for mRNA markers of oxidative stress, inflammation, vasoconstriction, thrombosis and cell filamentous components in rats exposed to AF-SWCNT revealed pulmonary induction of HO-1, MIP-2, endothelin-1, PAI-1 and β-catenin at both time points (1-d-4-d). However, mRNA expression for any of the biomarkers was not altered in the aorta at either time points. Pulmonary instillation of SWCNT produces acute pulmonary inflammatory, vasoconstrictive and prothrombotic effects in rats, whereas these effects are not evident within 4 days in the aorta. (Does not reflect US EPA policy). Supported in part by EPA SEE Program and EPA/UNC CR833237.

PS 2204 PULMONARY EFFECTS FROM ACUTE EXPOSURE TO AEROSOLIZED SINGLE-WALLED CARBON NANOTUBES.

A. K. Madl, K. Durinick, A. Lam, T. Guo, S. V. Teague, Y. Qu and K. E. Pinkerton. University of California, Davis, Davis, CA.

Although nanotechnology is still an emerging field and the enthusiasm for the potential societal benefits of engineered nanomaterials continues, concerns are being raised about whether our knowledge of possible health risks is keeping pace with products going to market. Single-walled carbon nanotubes (SWCNTs) have gained notable attention for application in a number of industries because of their unique electronic, optical, mechanical, chemical, or even biological properties. Due to the potential for human exposure, toxicological studies are needed to understand the potential health hazards of these nanomaterials. To better understand the biological responses associated with acute SWCNT exposure, Sprague Dawley rats were exposed to either aerosolized SWCNTs (300 or 1000 μg/m³, raw [FeSWCNT or purified cSWCNT) or fresh air via nose-only inhalation for 6 hours for 1 day. Cytotoxicity markers (total protein, lactate dehydrogenase [LDH], and gamma-glutamyl transferase [GGT]) in bronchoalveolar lavage and mucin and collagen staining in lung tissue were used as a means to assess immediate and persistent (0, 1, 3, 7 and 28 d post-exposure [PE]) effects of acute exposure to SWCNTs. Results showed that markers for cytotoxicity (LDH, GGT) were acutely increased up to 7 d PE, whereas mucin staining in proximal airways initially increased and then decreased at 7 and 28 d PE, and collagen staining in the alveoli increased at 7 d PE. Effects in all cases were transient, influenced by dose and particle composition (e.g., iron content), and generally were resolved by 7-28 d PE. While these data suggest that exposure to aerosolized SWCNTs may induce cytotoxic and structural responses in the lungs, further research is needed to evaluate whether these changes are suggestive of precursor events to pathological changes or lung remodeling that might develop under more severe or prolonged exposure conditions and may have implications for human health risk for persons potentially exposed to airborne SWCNTs.

PS 2205 INHALED MULTI-WALLED CARBON NANOTUBES STIMULATE A PLEURAL INFLAMMATORY RESPONSE IN THE LUNGS OF MICE.

J. C. Bonner¹, J. P. Ryman-Rasmussen^{1,4}, M. F. Cesta^{1,4,5}, A. R. Brody², J. K. Shipley-Phillips³, J. Everitt⁶, E. W. Tewksbury⁷, O. R. Moss⁴, B. A. Wong⁴, D. E. Dodd¹ and M. E. Andersen⁴. ¹Environmental and Molecular Toxicology, North Carolina State University, Raleigh, NC, ²Molecular and Biomedical Sciences, North Carolina State University, Raleigh, NC, ³Population Health and Pathobiology, North Carolina State University, Raleigh, NC, ⁴The Hamner Institutes for Health Sciences, Research Triangle Park, NC, ⁵Experimental Pathology, NIEHS, Research Triangle Park, NC and ⁶GlaxoSmithKline, Research Triangle Park, NC.

Carbon nanotubes have recently been reported to have asbestos-like properties since they stimulate the formation of pleural granulomas when injected into the abdominal cavity of mice. This has raised legitimate concerns over the safety of nanotubes because mesothelial granulomas that form on the pleural surface have the potential to develop into mesothelioma, a type of cancer associated with the inhalation of asbestos fibers. Here we report that male C57BL6 mice that inhaled an aerosol of multi-walled carbon nanotubes (6-hr exposure at 100 mg/m³) developed inflammatory foci on the pleural surface of the lung, even though very little inflammation or fibrosis was observed within the lung parenchyma. These foci were primarily monocytic and persisted at 14 days post-exposure. We observed nanotubes dispersed throughout the lung at 1 day post-exposure with some embedded within the pleural wall. Most of the carbon nanotubes (>90%) were contained within macrophages throughout the 14 day study period. Inhalation is the most relevant route of occupational exposure to carbon nanotubes, and our findings are the first to demonstrate that inhaled carbon nanotubes cause pleural inflammation. We emphasize that further work is urgently needed to assess whether longer term, lower concentration nanotube exposures cause similar pleural responses as seen with this

The Toxicologist


Supplement to *Toxicological Sciences*

An Official Journal of the
Society of Toxicology



SOT

Society of
Toxicology



48th Annual Meeting
and ToxExpo™
Baltimore, Maryland



49th Annual Meeting and ToxExpo™
Salt Lake City, Utah

2010

March 7-11, 2010 • Salt Palace Convention Center

SOT | Society of Toxicology

Photos courtesy of Salt Lake Convention and Visitors Bureau. Photographed by Richard Cheski, Steve Greenwood, and Eric Schramm.

Deadline for Proposals for SOT 2010 Annual Meeting Sessions: April 30, 2009

WHY SUBMIT A PROPOSAL?

1. To present new developments in toxicology.
2. To provide attendees an opportunity to learn about state-of-the-art technology and how it applies to toxicological research.
3. To provide attendees an opportunity to learn about the emerging fields and how they apply to toxicology.

SESSION TYPES

Continuing Education—Emphasis on quality presentations of generally accepted, state-of-the-art knowledge in toxicology

Note: CE Courses will be held on Sunday.

Symposia—“Cutting-edge” science; new areas, concepts, or data

Workshops—State-of-the-art knowledge in toxicology

Roundtables—Controversial subjects

Historical Highlights—Review of a historical body of science that has impacted toxicology

Informational Sessions—Scientific planning or membership development

Education-Career Development Sessions—Sessions that provide the tools and resources to toxicologists that will enhance their professional and scientific development

2010 Thematic Approach

The Scientific Program Committee will continue the thematic approach for the 2010 Annual Meeting. All proposal submissions will be reviewed for their relevance under the following themes—*Cell Signaling, Gene-Environment Interactions, Metabolic Disease, Mitochondrial Basis of Disease, Toxicity Testing in the 21st Century*, and *Translational Toxicology* for the 2010 meeting. Please note that while we are actively soliciting proposals for the themes listed above, all proposal submissions will be reviewed under the current criteria for their timeliness and relevance to the field of toxicology.

Please refer to the SOT 2009 *Program*, Scientific Program Overview on the fold-out cover for a list of 2009 sessions highlighted under the thematic approach.

You can now submit your proposal on-line at www.toxicology.org

Preface

This issue of *The Toxicologist* is devoted to the abstracts of the presentations for the continuing education, symposia, workshop, roundtable, platform, and poster discussion sessions of the 48th Annual Meeting of the Society of Toxicology, held at the Baltimore Convention Center, March 15–19, 2009.

An alphabetical Author Index, cross referencing the corresponding abstract number(s), begins on page 469.

The issue also contains a Key Word Index (by subject or chemical) of all the presentations, beginning on page 487.

The abstracts are reproduced as accepted by the Scientific Program Committee of the Society of Toxicology and appear in numerical sequence.

Copies of *The Toxicologist* are available at \$45 each plus \$5 postage and handling (U.S. funds) from:

Society of Toxicology
1821 Michael Faraday Drive, Suite 300
Reston, VA 20190

www.toxicology.org

© 2009 Society of Toxicology

All text and graphics are © 2009 by the Society of Toxicology unless noted. Some Baltimore photos are courtesy of the Baltimore Area Convention and Visitors Association unless otherwise noted. For promotional use only. No advertising use is permitted.

This abstract book has been produced electronically by ScholarOne, Inc. Every effort has been made to faithfully reproduce the abstracts as submitted. The author(s) of each abstract appearing in this publication is/are solely responsible for the content thereof; the publication of an article shall not constitute or be deemed to constitute any representation by the Society of Toxicology or its boards that the data presented therein are correct or are sufficient to support the conclusions reached or that the experiment design or methodology is adequate. Because of the rapid advances in the medical sciences, we recommend that independent verification of diagnoses and drug dosage be made.