

References

1. Rabe KF, Hurd S, Anzueto A, Barnes PJ, Buist SA, Calverley P, Fukuchi Y, Jenkins C, Rodriguez-Roisin R, van Weel C, *et al.* Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* 2007;176:532–555.
2. Potter WA, Olafsson S, Hyatt RE. Ventilatory mechanics and expiratory flow limitation during exercise in patients with obstructive lung disease. *J Clin Invest* 1971;50:910–919.
3. O'Donnell DE, Revill SM, Webb KA. Dynamic hyperinflation and exercise intolerance in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2001;164:770–777.
4. O'Donnell DE, Parker CM. COPD exacerbations. 3: Pathophysiology. *Thorax* 2006;61:354–361.
5. O'Donnell DE, Laveneziana P. The clinical importance of dynamic lung hyperinflation in COPD. *COPD* 2006;3:219–232.
6. O'Donnell DE, Voduc N, Fitzpatrick M, Webb KA. Effect of salmeterol on the ventilatory response to exercise in chronic obstructive pulmonary disease. *Eur Respir J* 2004;24:86–94.
7. Peters MM, Webb KA, O'Donnell DE. Combined physiological effects of bronchodilators and hyperoxia on exertional dyspnoea in normoxic COPD. *Thorax* 2006;61:559–567.
8. Aliverti A, Rodger K, Dellaca RL, Stevenson N, Lo MA, Pedotti A, Calverley PMA. Effect of salbutamol on lung function and chest wall volumes at rest and during exercise in COPD. *Thorax* 2005;60:916–924.
9. Stevenson NJ, Walker PP, Costello RW, Calverley PMA. Lung mechanics and dyspnea during exacerbations of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2005;172:1510–1516.
10. Stevenson NJ, Calverley PM. Effect of oxygen on recovery from maximal exercise in patients with chronic obstructive pulmonary disease. *Thorax* 2004;59:668–672.
11. Enright PL. GOLD stage I is not a COPD risk factor. *Thorax* 2007;62:1107–1109.
12. Hardie JA, Buist AS, Vollmer WM, Ellingsen I, Bakke PS, Morkve O. Risk of over-diagnosis of COPD in asymptomatic elderly never-smokers. *Eur Respir J* 2002;20:1117–1122.
13. Ofir D, Laveneziana P, Webb KA, Lam Y-M, O'Donnell DE. Mechanisms of dyspnea during cycle exercise in symptomatic patients with GOLD Stage I chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2008;177:622–629.
14. Dellacà RL, Duffy N, Pompilio PP, Aliverti A, Koulouris NG, Pedotti A, Calverley PM. Expiratory flow limitation detected by forced oscillation and negative expiratory pressure. *Eur Respir J* 2007;29:363–374.
15. Rutten FH, Moons KGM, Cramer M-JM, Grobbee DE, Zuithoff NPA, Lammers J-WJ, Hoes AW. Recognising heart failure in elderly patients with stable chronic obstructive pulmonary disease in primary care: cross sectional diagnostic study. *BMJ* 2005;331:1379–1385.

DOI: 10.1164/rccm.200712-1792ED

Thinking Longitudinally in a Cross-sectional World

Asbestos exposure is widely recognized to cause serious and often deadly health effects (1). The World Health Organization and the International Labour Organization have called for asbestos bans worldwide, and more than 40 countries have either banned or severely restricted asbestos use (2, 3). The U.S. Congress is currently considering legislation entitled the “Ban Asbestos in America Act of 2007” (4). In this issue of the *Journal* (pp. 630–637), Rohs and colleagues report dose-dependent pleural changes identified in a cross-sectional follow-up study of a cohort of workers exposed to asbestos-contaminated vermiculite originating in Libby, Montana (5). The study has particular relevance to current national and international efforts to control risk from exposure to durable fibers.

Vermiculite is a mineral that expands when rapidly heated. Expanded vermiculite is used in both industrial and consumer products, such as loose-fill attic insulation, gardening and agricultural products, and automobile brakes. Between 1925 and 1990, the majority of the world's vermiculite was mined in Libby, Montana. Amphibole minerals, including tremolite asbestos, contaminated the Libby vermiculite products (6).

In 1980, following reports of bloody pleural effusions at one industrial site, Lockey and colleagues examined 513 workers exposed to Libby vermiculite; 2.2% showed radiographic evidence of either interstitial fibrosis or pleural changes consistent with asbestos effects (7). This was an important finding at the time, especially since exposure to vermiculite had been thought by many to be essentially benign and knowledge of the contamination of Libby vermiculite was limited. The Rohs and colleagues' study examines the same cohort of workers 25 years later. The proportion of affected workers has risen to 28.7% for pleural changes and 2.9% for fibrosis, despite no known interim exposure to asbestos. These findings are consistent with the well-known capability of asbestos to cause both malignant and non-malignant changes years after an original exposure—effects that increase as residence time of the inhaled, biopersistent fibers lengthens (1).

The current study makes important new contributions to our understanding of the consequences of exposure to durable

fibers. The investigators estimated cumulative fiber exposures for each worker based on job history and historical records of plant industrial hygiene evaluations. As expected, workers with the most exposure had the greatest probability of pleural abnormalities. What is striking, however, is the level of adverse effects in the low-exposed group: 13.9% of workers with cumulative exposures estimated between 0.25 and 0.74 fibers/cc-years showed pleural abnormalities. This compares with a current legal permissible exposure limit (PEL) in the United States that limits asbestos exposures to 0.45 fibers/cc-years over a 45-year working lifetime (8), indicating that a significant number of workers exposed at the current exposure limit would experience pleural abnormalities. Although arguments persist about the extent to which pleural abnormalities are inherently harmful, it is clear that the presence of pleural changes demonstrates a physiologic response to durable fibers and is associated with increased risk of pulmonary fibrosis, lung cancer, and mesothelioma (1).

A notable aspect of this study is the contemporary classification of the fibers contaminating the Libby vermiculite. In 1980, these fibers were classified as “tremolite asbestos” and, as such, fit within the definition of regulated asbestos fibers (8). (U.S. regulations define asbestos as one of six specific mineral fibers that are or have been of commercial value.) The current study notes that the primary durable fibers contaminating the vermiculite are winchite and richterite with perhaps 10% tremolite asbestos; however, winchite and richterite are not explicitly regulated as asbestos. This investigation does not prove that the health effects of these fibers are identical to tremolite; however, their chemical and physical properties are virtually identical to tremolite, indicating the importance of evaluating potentially harmful fibers not only on whether they meet a mineralogical definition of asbestos but also on their biologically relevant features of durability, biopersistence, shape, size, and surface characteristics. The Ban Asbestos Act explicitly expands the definition of asbestos to include winchite and richterite (4). The U.S. National Institute for Occupational Safety and Health has circulated a draft “Roadmap for Scientific Research” proposing

studies that would be useful in better understanding the nature of hazardous fibers (9).

The increase in fiber-related disease from exposure to the contaminated vermiculite is not limited to the group of industrial workers examined recently by Rohs and colleagues and previously by Lockey and coworkers. Following the original report by Lockey and colleagues, investigators studied the morbidity and mortality of vermiculite miners and millers in Libby, Montana, and found increased risk of lung cancer and nonmalignant respiratory disease (10, 11). Consistent with the Rohs report, a recent update of that cohort mortality study confirmed significant excess deaths from asbestosis, lung cancer, cancer of the pleura, and mesothelioma at fiber exposure levels lower than in the original study (12).

These findings, as a whole, highlight the importance of interpreting and responding to cross-sectional research findings not only according to what they say about the present but also in the context of what they indicate about the future. The initial Lockey investigation found a relatively modest 2.2% prevalence of pleural abnormalities; Rohs and colleagues found over 10 times that level, despite the fact that contaminated vermiculite had been removed from the production process by 1980. In Libby, Montana, miners and millers continued to be exposed to the contaminated vermiculite until mining operations ceased in 1990, and their current mortality experience, after continuing exposure and longer latency, reflects this as well (12). Consumers of vermiculite originating in Libby have also been exposed to potentially hazardous fibers (13).

Anticipation of potential risk is the keystone of occupational disease and injury prevention (14). Had the early cross-sectional investigations of disease from contaminated vermiculite stimulated assessment of future risk, it is possible that the number of people exposed to hazardous fibers, their level of exposure, and disease risk could have been reduced.

The opportunity for prevention of fiber-related morbidity and mortality remains. Recent investigations have demonstrated strong relationships between national asbestos consumption rates and levels of asbestos-associated mortality after a lengthy lag (15). Although the United States has successfully restricted asbestos exposure and is considering a ban, consumption in some countries with rapidly developing economies is continuing to expand (16). Bans and effective use restrictions worldwide have the potential to prevent decades of future death and disease.

Wayne Gretzky, an exceptional ice hockey player, famously attributed his athletic accomplishments to the ability to "skate to where the puck is going to be." Through more effective response to cross-sectional scientific findings based on understanding their future implications, particularly for long-latency diseases, the medical and public health communities have the opportunity to emulate Gretzky's success.

Conflict of Interest Statement: G.R.W. has no financial relationship with a commercial entity that has an interest in the subject of this manuscript.

GREGORY R. WAGNER, M.D.
National Institute for Occupational Safety and Health
Washington, D.C.

The findings and conclusions expressed in this editorial are those of the author and do not necessarily represent the views of the National Institute for Occupational Safety and Health.

References

1. American Thoracic Society. Diagnosis and initial management of non-malignant diseases related to asbestos: official statement of the American Thoracic Society. *Am J Respir Crit Care Med* 2004;170:691-715.
2. World Health Organization. Elimination of asbestos-related diseases, World Health Organization 2006 [Internet] [accessed Nov 14, 2007]. Available from: http://www.who.int/occupational_health/publications/asbestosrelateddiseases.pdf
3. Kazan-Allen L. Current asbestos bans and restrictions: July 2007 [Internet] [accessed Nov 14, 2007]. Available from: http://www.ibasecretariat.org/lka_alpha_asb_ban_280704.php
4. Ban Asbestos in America Act of 2007, H.R. 3285, 110th Cong, 1st Sess. (2007).
5. Rohs AM, Lockey JE, Dunning KK, Shukla R, Fan H, Hilbert T, Borton E, Wiot J, Meyer C, Shipley TR, et al. Low-level fiber-induced radiographic changes caused by Libby vermiculite: a 25-year follow-up study. *Am J Respir Crit Care Med* 2008;177:630-637.
6. National Institute for Occupational Safety and Health. Fact sheet: NIOSH recommendations for limiting potential exposures of workers to asbestos associated with vermiculite from Libby, Montana [Internet]. DHHS (NIOSH) Publication No. 2003-141; May 2003. [Accessed 2008 Jan 23]. Available from: <http://www.cdc.gov/niosh/docs/2003-141>
7. Lockey JE, Brooks SM, Jarabek AM, Khoury PR, McKay RT, Carson A, Morrison JA, Wiot JF, Spitz HB. Pulmonary changes after exposure to vermiculite contaminated with fibrous tremolite. *Am Rev Respir Dis* 1984;129:952-958.
8. Code of Federal Regulations. Codified at 29 CFR 1910.1001: Asbestos [Internet] [accessed 2008 Jan 23]. Available from: http://www.osha.gov/pls/oshweb/owadisp.show_document?p_table=STANDARDS&p_id=9995
9. National Institute for Occupational Safety and Health Mineral Fibers Group. Asbestos and other mineral fibers: a roadmap for scientific research. February 2007 [Internet] [accessed Nov 14, 2007]. Available from: <http://www.cdc.gov/niosh/review/public/099/pdfs/NIOSHAsbestosRoadmap.pdf>
10. Amandus HE, Althouse R, Morgan WK, Sargent EN, Jones R. The morbidity and mortality of vermiculite miners and millers exposed to tremolite-actinolite: part III. Radiographic findings. *Am J Ind Med* 1987;11:27-37.
11. Amandus HE, Wheeler R. The morbidity and mortality of vermiculite miners and millers exposed to tremolite-actinolite: part II. Mortality. *Am J Ind Med* 1987;11:15-26.
12. Sullivan PA. Vermiculite, respiratory disease, and asbestos exposure in Libby, Montana: update of a cohort mortality study. *Environ Health Perspect* 2007;115:579-585.
13. Agency for Toxic Substances and Disease Registry. Vermiculite consumer products. May 2003 [Internet] [accessed Nov 14, 2007]. Available from: <http://www.atsdr.cdc.gov/NEWS/vermiculite051603.html>
14. Levy BS, Wagner GR, Rest KM, Weeks JL, editors. Preventing occupational disease and injury, 2nd ed. Washington, DC: American Public Health Association (APHA); 2005.
15. Lin RT, Takahashi K, Karjalainen A, et al. Ecological association between asbestos-related diseases and historical asbestos consumption: an international analysis. *Lancet* 2007;369:844-849.
16. Virta RL. Worldwide asbestos supply and consumption trends from 1900 through 2003 [Internet] [accessed Nov 14, 2007]. U.S. Geological Survey Circular 1298; 2006. Available from: <http://pubs.usgs.gov/circ/2006/1298/c1298.pdf>

DOI: 10.1164/rccm.200711-1703ED