

RESEARCH

Open Access

Cytogenetic analysis of an exposed-referent study: perchloroethylene-exposed dry cleaners compared to unexposed laundry workers

James D Tucker^{1,2*}, Karen J Sorensen², Avima M Ruder³, Lauralynn Taylor McKernan³, Christy L Forrester⁴, Mary Ann Butler³

Abstract

Background: Significant numbers of people are exposed to tetrachloroethylene (perchloroethylene, PCE) every year, including workers in the dry cleaning industry. Adverse health effects have been associated with PCE exposure. However, investigations of possible cumulative cytogenetic damage resulting from PCE exposure are lacking.

Methods: Eighteen dry cleaning workers and 18 laundry workers (unexposed controls) provided a peripheral blood sample for cytogenetic analysis by whole chromosome painting. Pre-shift exhaled air on these same participants was collected and analyzed for PCE levels. The laundry workers were matched to the dry cleaners on race, age, and smoking status. The relationships between levels of cytological damage and exposures (including PCE levels in the shop and in workers' blood, packyears, cumulative alcohol consumption, and age) were compared with correlation coefficients and t-tests. Multiple linear regressions considered blood PCE, packyears, alcohol, and age.

Results: There were no significant differences between the PCE-exposed dry cleaners and the laundry workers for chromosome translocation frequencies, but PCE levels were significantly correlated with percentage of cells with acentric fragments ($R^2 = 0.488$, $p < 0.026$).

Conclusions: There does not appear to be a strong effect in these dry cleaning workers of PCE exposure on persistent chromosome damage as measured by translocations. However, the correlation between frequencies of acentric fragments and PCE exposure level suggests that recent exposures to PCE may induce transient genetic damage. More heavily exposed participants and a larger sample size will be needed to determine whether PCE exposure induces significant levels of persistent chromosome damage.

Background

Tetrachloroethylene (perchloroethylene; PCE; CAS No. 127-18-4) is a chlorinated solvent with widespread use as a degreasing agent and as a solvent in the dry cleaning industry. Over 1.5 million workers are exposed to this compound every year [1], with historical mean occupational exposure levels of 59 ppm and higher depending on the application [2]. While current mean levels of exposure to PCEs are substantially lower, probably in the range of 1-10 ppm, concerns about

exposures continue because PCE is an established animal carcinogen [3,4] causing leukemias and kidney and liver tumors. IARC considers PCE to be a probable human carcinogen and a definite animal carcinogen [5]. PCE is one of twenty common occupational carcinogens "where evidence of carcinogenicity is substantial but not yet conclusive for humans", and for which more research is recommended [6]. The current Occupational Safety and Health Administration permissible exposure limit is 100 ppm which is based on neurotoxic effects; the current National Institute for Occupational Safety and Health (NIOSH) recommended exposure limit is "the lowest feasible level", which is based on potential carcinogenicity [7].

* Correspondence: jtucker@biology.biosci.wayne.edu

¹Department of Biological Sciences, 2117 Biological Sciences Building, 5047 Gullen Mall, Wayne State University, Detroit, MI 48202-3917 USA
Full list of author information is available at the end of the article

Numerous studies have examined PCE-exposed workers for adverse health effects (reviewed in [1,2]). Increased risks of scleroderma as well as biliary and liver or brain cancer have been observed as a result of exposure, as have neurological impairments, increased reproductive failure among women including spontaneous abortions, and eccentric sperm morphology and motility among men. Recently, neurobehavioral deficits following PCE exposure have also been reported [8]. Human exposure to PCE sometimes occurs in parallel with other agents and these exposures have been associated with a wide variety of cancers [9-11]. However, it has not been possible to determine whether these human tumors are due to PCE alone, or to other chemicals, or both. Prenatal exposures to PCE have also occurred and have been associated with increases in spontaneous abortions [12,13]. Prenatal exposures to PCE have been assessed in a retrospective cohort study of contaminated drinking water, and no significant effects on birth weight or gestational duration were observed [14]. Similarly, prenatal and early postnatal exposure to PCE-contaminated drinking water were not associated with developmental disorders of attention, learning or behavior identified on the basis of questionnaire responses [15].

Two studies have investigated the cytogenetic effects of PCE in occupationally-exposed humans. Ikeda et al. [16] evaluated chromosomal aberrations and sister chromatid exchanges (SCEs) in lymphocytes from 10 participants. No significant dose-related changes were observed in the frequencies of numerical or structural aberrations, SCEs, or any of several measures of cell growth. Seiji et al. [17] evaluated 27 workers exposed to PCE and did not observe a significant increase in SCE frequencies compared to the unexposed controls.

SCEs and chromosome aberrations are markers of early biologic effects of exposure. However neither endpoint shows persistence beyond a few years following acute exposure. Most types of structural aberrations, including dicentrics and acentric fragments, decline significantly with time after exposure because these aberrations kill cells and as a result their frequencies eventually return to baseline [18-20]. Under conditions of chronic exposure, the induction of SCEs or dicentrics and fragments would be in equilibrium with their loss, possibly reaching a plateau above the baseline levels. However, because of this loss of events over time, neither SCEs nor dicentrics and fragments are accurate biomarkers for quantifying chronic exposure, because detection of the total amount of exposure depends on the accumulation of damage over prolonged periods of time. In contrast, chromosome translocations are compatible with cell survival. For this reason they show much greater persistence through cell division [21,22]

and consequently their frequencies are believed to integrate the effect of chronic exposure [23-25]. Thus, while PCE exposure is of concern to human health, investigations of possible cumulative genetic damage resulting from exposure are lacking. For these reasons we sought to evaluate PCE-exposed workers by quantifying translocations, which are not only a persistent biomarker of exposure but also a biomarker of effect because translocations are present in almost all tumors [26]. Here we report the results of a study investigating 18 PCE-exposed female dry cleaners and an unexposed control group comprised of an equal number of female laundry workers who were not exposed to PCE. Cytogenetic analyses for chromosome translocations were performed by whole-chromosome painting in peripheral blood lymphocytes.

Methods

Study participants

This study was approved by the NIOSH Human Subjects Review Board. Study design and methods have been described previously [27,28]. Briefly, 18 dry cleaning workers and 18 laundry workers (unexposed controls) were recruited from seven shops in and around southwest Ohio. All participants were women under age 70 who signed an informed consent form and completed an interview which included occupational, smoking, and drinking histories. None of the participants, when asked, reported having had chemotherapy or radiotherapy. Dry cleaners were machine operators or pressers for one year or more (range 1-19, mean 8, standard deviation 5) in shops with third generation machines containing only refrigerated condensers or fourth generation machines with refrigerated condensers and carbon adsorbers [27]. Laundry workers had been in the industry for at least one year and had never been exposed to PCE. Laundry workers were matched by race (Caucasian or African-American), smoking status, and age (\pm 5 years when possible) to already selected dry cleaners. Personal breathing zone samples were collected from dry cleaners on Wednesday of a typical work week. Two or more area samples and personal breathing zone samples were collected from a minimum of two workers in each laundry. Sampling analysis determined concentration of PCE in air according to NIOSH Method 1003 [29].

Blood collection and air sampling

Venous blood was collected from dry cleaners before work on a Thursday following three consecutive days of PCE exposure, and from launderers on a typical work day. At the same time, pre-shift end-exhaled air was collected and analyzed by NIOSH Method 3704 [30]. The blood specimens were shipped to the PCE-analysis laboratory and refrigerated upon arrival. The average

time between blood drawing and analysis was 32 (range 6 to 64) days. All exposure measurements of air levels were performed in accordance with procedures of the NIOSH Analytical Methods Manual [29]. All pre-shift specimens were collected before the dry cleaners entered their dry cleaning shops. Blood was obtained via venipuncture. Blood samples for measuring PCE were collected in gray-top vacutainers that had been previously processed to remove volatile contaminants [31]. These whole blood samples were kept at refrigerator temperature until analysis. Blood PCE was measured by the method of Ashley et al. [31]. Volatile compounds were separated from the blood matrix by purge and trap concentration. Detection, identification, and quantification of PCE was performed by GC/MS. Quantification was achieved by isotope dilution and reference to commercially available standard compounds [31].

Chromosome painting

For the fluorescent in situ hybridization (FISH) whole chromosome painting analysis, 5 ml blood were obtained in heparinized Vacutainer CPT tubes, and transported at refrigeration temperature to the Lawrence Livermore National Laboratory for arrival within 24 hours of collection. All samples were then immediately set up in culture as described [32] for 52 hours, the last 4 in the presence of 0.1 µg/ml Colcemid. Cells were then swollen with 75 mM KCl as hypotonic, fixed three times in methanol: acetic acid (3:1 v/v), dropped onto clean microscope slides, air dried, then stored in the presence of N₂ gas and CaSO₄ desiccant at -20°C until needed for hybridization. Chromosomes 1, 2, and 4 were painted red and chromosomes 3, 5, and 6 were simultaneously painted green, then counterstained with DAPI (blue) as described [32]. An average of 766 (range 751 to 919) whole genome equivalents was scored from each participant. The structural chromosome aberrations scored included translocations, insertions, dicentric and acentric fragments. Color junctions were also enumerated, and included any chromosome rearrangement that leads to the misunion of chromosome pieces painted in different colors. Most color junctions are the result of translocations, with smaller contributions from insertions, dicentric and acentric fragments. Complete terminology and specific analytical techniques have been described [32,33].

Statistical analyses

SAS[®] (Version 9.1, SAS Institute, Cary, NC, USA) was used to analyze the data. Pearson correlation coefficients were used to determine the unadjusted relationships between the PCE exposure indices (TWA {time-weighted average}, breath, and blood PCE) and the FISH biomarkers (percent of cells with structural chromosome

aberrations). Differences in the percent of cells with aberrations (translocations, insertions, dicentric, acentric fragments, and all exchanges, i.e. color junctions) between exposed and unexposed workers were evaluated with t-tests. We evaluated the percent of cells with each type of aberration, rather than the number of aberrations per cell, because this allowed us to avoid problems with enumerating different types of translocations [22], e.g. differential counting of reciprocal and non-reciprocal exchanges. For the sake of consistency, the other types of aberrations evaluated here are reported in this same manner. Multiple linear regression models were used to determine the adjusted relationships between the natural log blood PCE and each type of aberration. The data were log-transformed to obtain approximate normal distributions. The covariates included in the models were pack years (cumulative smoking index), log cumulative alcohol intake, and age. Positive relationships between exposure and aberration level were considered statistically significant, using one-tailed tests, if $p < 0.05$.

Results and Discussion

The demographics of the study participants are shown in Table 1. There were no significant differences

Table 1 Data Summary

Demographics	Laundry Workers	Dry Cleaners
Facilities	3 ^a	4
Workers	18 ^a	18
Race, Caucasian/African American	12/6	13/5
Current smokers	10	10
Age	39 ± 9 ^b	40 ± 13
Exposure Indices		
PCE TWA (ppm) ^c	<0.02	3.8 ± 5.3
Pre-shift end-exhaled PCE (ppm) ^d	–	0.45 ± 0.33
Post-shift end-exhaled PCE (ppm) ^d	–	1.21 ± 0.87
Blood PCE	0.19 ± 0.44 µg/l	74.81 ± 104.27 µg/l ^e
Pack years ^f	8.44 ± 11.32	13.13 ± 12.0
Lifetime number of drinks ^g	1,106 ± 2,609	6,382 ± 18,144

^aValues are number of facilities that contributed participants to this study, workers who are Caucasian or African American, and number of smokers. 20 laundry workers were recruited but only the 18 who donated blood specimens are included here.

^bAll values are mean ± SD.

^cPCE TWA in laundries was below the level of detection (0.023 ppm).

^dEnd-exhaled breath was collected from laundry workers but only one had a PCE level (4.5 ppb) above the limit of detection.

^eMean is significantly different than the corresponding mean for laundry workers (t-test, $P < 0.05$).


^fPacks/day × smoking years.

^gDrinks ("cans or glasses of beer, glasses of wine, shots of hard liquor")/week × 52 × drinking years.

between the PCE-exposed dry cleaners and the laundry workers (unexposed controls) for race, current cigarette smoking status, or age. PCE TWA in laundries was below the level of detection (0.023 ppm) for both the personal breathing zone and the area monitors. Post-shift exhaled breath PCE levels in dry cleaners were generally higher than their pre-shift levels. Because blood PCE levels were available both for dry cleaners and laundry workers, and are considered the preferred exposure measure, they were used as the occupational exposure independent variable [27].

Regression analyses were performed using age, log of the blood PCE levels, log of the cumulative alcohol consumption, and pack years as the independent variables. A total of five dependent variables were evaluated; these are the percent of cells with chromosome translocations, inversions, dicentrics, fragments, and color junctions. Each dependent variable was regressed against the set of four independent variables. None of the chromosome aberration types showed a significant effect of PCE exposure, or effects due to cigarette smoking or alcohol consumption (data not shown). The absence of an effect of PCE exposure upon translocations is evident in Figure 1. However, as shown in Figure 2, translocations did show a significant effect with age, which is consistent with a recently published report of more than 1900 unexposed participants [34]. Color junctions also showed a significant effect with age but this is likely to be due to the inclusion of translocations in this endpoint as the frequencies of dicentrics, fragments, and insertions showed no age effect.

Correlation analyses were also performed among the variables. PCE blood level, TWA of the blood PCE levels, and exhaled breath were highly correlated [27]. No correlation was observed between the TWA of PCE and the percent of cells with chromosome translocations,



inversions, dicentrics, and color junctions. However, PCE levels were significantly correlated with acentric fragments ($p = 0.0026$, Figure 3). In addition, the dry cleaners had higher percentages of cells with translocations, insertions, color junctions, dicentrics and acentric fragments than did the laundry workers but none of these increases were statistically significant (Table 2).

The dry cleaning plants where the exposed workers were employed had low ambient levels of PCE, and only one of the exposed workers was a full-time machine operator. Since operators have much higher exposures than do other employees, and since many shops have higher ambient PCE levels, this does not rule out a PCE-exposure effect among more heavily exposed workers.

Conclusions

In this population there does not appear to be a strong effect of PCE exposure on chromosome damage, even





Table 2 Cytogenetic comparisons of 18 dry-cleaning workers exposed to tetrachloroethylene and 18 unexposed laundry workers

Variable	PCE exposed?	Percent	Confidence Interval	Pooled t-test	Pr > t *
% cells with translocations	no	0.599	(0.329 - 0.870)	-0.93	0.18
	yes	0.747	(0.546 - 0.948)		
% cells with insertions	no	0.0073	(-0.008 - 0.0229)	-1.18	0.12
	yes	0.0291	(-0.007 - 0.0648)		
% cells with color junctions	no	0.621	(0.338 - 0.905)	-1.13	0.13
	yes	0.811	(0.597 - 1.025)		
% cells with dicentrics	no	0.0365	(0.0064 - 0.0667)	-1.99	0.027
	yes	0.0873	(0.0426 - 0.132)		
% cells with acentric fragments	no	0.161	(0.0641 - 0.258)	-1.30	0.10
	yes	0.253	(0.140 - 0.367)		

*Probability of a positive relationship between exposure and level of aberrations (one-tailed test).

for translocations that measure accumulated exposure. However, the dry cleaners did have non-significant increases in translocations, insertions, fragments, dicentrics and junctions compared to laundry workers. Further work on more heavily exposed participants and with a larger sample size will be needed to determine whether PCE exposure induces significant levels of cytogenetic damage.

List of Abbreviations

PCE: tetrachloroethylene, perchloroethylene; NIOSH: National Institute for Occupational Safety and Health; SCEs: sister chromatid exchanges; FISH: fluorescence in situ hybridization; TWA: time-weighted average.

Acknowledgements

This work was funded by an Interagency Agreement between the National Institute for Occupational Safety and Health and the Lawrence Livermore National Laboratory. We thank David L. Ashley for supplying the special tubes in which the blood for analysis of tetrachloroethylene was collected, and for performing the analyses of the blood tetrachloroethylene levels. We thank Jodie Ghore and Cathy Chu for expert technical assistance in performing the cytogenetic analyses. We acknowledge Lynda Ewers and Ed Burroughs for their technical expertise regarding breath analysis, and thank Donald Booher, Marian Coleman, James Deddens, Kevin L. Dunn, Jensen Groff, Timothy Jiggins, Jun Ju, Xiudong Lei, Sharon Lemire, Barbara MacKenzie, Leroy May, Virginia O'Neill, Susan Welch, Larry Wetzel and Jim Woodfin for their assistance during specimen collection and data analysis. This study was supported in part by the CDC Office of Women's Health. This work was performed in part under the auspices of the U.S. Department of Energy by the Lawrence Livermore National Laboratory under contract number W7405-ENG-48. The findings and conclusions in this manuscript are those of the author(s) and do not necessarily represent the views of the National Institute for Occupational Safety and Health.

Author details

¹Department of Biological Sciences, 2117 Biological Sciences Building, 5047 Gullen Mall, Wayne State University, Detroit, MI 48202-3917 USA. ²Physical and Life Sciences Directorate, L-452, Lawrence Livermore National

Laboratory, Livermore, CA 94550 USA. ³National Institute for Occupational Safety and Health, CDC, 4676 Columbia Parkway, Cincinnati, OH 45226 USA. ⁴National Institute for Occupational Safety and Health, CDC, CDCW Bldg PATRI Room 9271, Washington, DC 20201 USA.

Authors' contributions

JDT oversaw the cytogenetic portion of this work and wrote the first draft of the paper. KJS performed the tissue culturing and the cytogenetics work. LTM contributed to the original protocol, oversaw specimen collection, and conducted breath analyses. AMR contributed to the original protocol, participated in the field study, and did statistical analyses. CLF and MAB contributed to the original protocol and participated in the field study. All authors have read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 2 December 2010 Accepted: 10 March 2011

Published: 10 March 2011

References

- Ruder AM: **Potential health effects of occupational chlorinated solvent exposure.** *Ann N Y Acad Sci* 2006, **1076**:207-227.
- Gold LS, De Roos AJ, Waters M, Stewart P: **Systematic literature review of uses and levels of occupational exposure to tetrachloroethylene.** *J Occup Environ Hyg* 2008, **5**:807-839.
- National Cancer Institute: *In Bioassay of tetrachloroethylene for possible carcinogenicity. Volume 13.* Bethesda, MD: National Cancer Institute; 1977.
- National Toxicology Program: **Toxicology and carcinogenesis studies of tetrachloroethylene (perchloroethylene) (CAS No.127-18-4) in F344/N rats and B6C3F1 mice.** *Technical Report Series* 1986, 1-190.
- International Agency for Research on Cancer: **Tetrachloroethylene.** *IARC Monogr Eval Carcinog Risks Hum* 1995, **63**:159-221.
- Ward EM, Schulte PA, Straif K, Hopf NB, Caldwell JC, Carreon T, DeMarini DM, Fowler BA, Goldstein BD, Hemminki K, Hines CJ, Pursiainen KH, Kuempel E, Lewtas J, Lunn RM, Lynde E, McElvenny DM, Muhle H, Nakajima T, Robertson LW, Rothman N, Ruder AM, Schubauer-Berigan MK, Siemiatycki J, Silverman D, Smith MT, Sorahan T, Steenland K, Stevens RG, Vineis P, Zahm SH, Zeise L, Coglianò VJ: **Research recommendations for selected IARC-classified agents.** *Environ Health Perspect* 2010, **118**:1355-1362.
- National Institute for Occupational Safety and Health: **NIOSH Pocket Guide to Chemical Hazards.** Cincinnati, Ohio: DHHS; 2004.

8. Benignus VA, Boyes WK, Geller AM, Bushnell PJ: **Long-term perchloroethylene exposure: a meta-analysis of neurobehavioral deficits in occupationally and residentially exposed groups.** *J Toxicol Environ Health A* 2009, **72**:824-831.
9. Blair A, Stewart PA, Tolbert PE, Grauman D, Moran FX, Vaught J, Rayner J: **Cancer and other causes of death among a cohort of dry cleaners.** *Br J Ind Med* 1990, **47**:162-168.
10. Ruder AM, Ward EM, Brown DP: **Mortality in dry-cleaning workers: an update.** *Am J Ind Med* 2001, **39**:121-132.
11. Aschengrau A, Ozonoff D, Paulu C, Coogan P, Vezina R, Heeren T, Zhang Y: **Cancer risk and tetrachloroethylene-contaminated drinking water in Massachusetts.** *Arch Environ Health* 1993, **48**:284-292.
12. Doyle P, Roman E, Beral V, Brookes M: **Spontaneous abortion in dry cleaning workers potentially exposed to perchloroethylene.** *Occup Environ Med* 1997, **54**:848-853.
13. Kolstad HA, Brandt LP, Rasmussen K: **[Chlorinated solvents and fetal damage. Spontaneous abortions, low birth weight and malformations among women employed in the dry-cleaning industry].** *Ugeskr Laeger* 1990, **152**:2481-2482.
14. Aschengrau A, Weinberg J, Rogers S, Gallagher L, Winter M, Vieira V, Webster T, Ozonoff D: **Prenatal exposure to tetrachloroethylene-contaminated drinking water and the risk of adverse birth outcomes.** *Environ Health Perspect* 2008, **116**:814-820.
15. Janulewicz PA, White RF, Winter MR, Weinberg JM, Gallagher LE, Vieira V, Webster TF, Aschengrau A: **Risk of learning and behavioral disorders following prenatal and early postnatal exposure to tetrachloroethylene (PCE)-contaminated drinking water.** *Neurotoxicol Teratol* 2008, **30**:175-185.
16. Ikeda M, Koizumi A, Watanabe T, Endo A, Sato K: **Cytogenetic and cytokinetic investigations on lymphocytes from workers occupationally exposed to tetrachloroethylene.** *Toxicol Lett* 1980, **5**:251-256.
17. Seiji K, Jin C, Watanabe T, Nakatsuka H, Ikeda M: **Sister chromatid exchanges in peripheral lymphocytes of workers exposed to benzene, trichloroethylene, or tetrachloroethylene, with reference to smoking habits.** *Int Arch Occup Environ Health* 1990, **62**:171-176.
18. Carrano A: **Chromosome aberrations and radiation induced cell death I. Transmission and survival parameters of aberrations.** *Mutat Res* 1973, **17**:341-353.
19. Carrano A: **Chromosome aberrations and radiation induced cell death II. Predicted and observed cell survival.** *Mutat Res* 1973, **17**:355-366.
20. Tucker JD, Strout CL, Christensen ML, Carrano AV: **Sister chromatid exchange induction and persistence in peripheral blood and spleen lymphocytes of mice treated with ethylnitrosourea.** *Environ Mutagen* 1986, **8**:345-355.
21. Tucker JD, Cofield J, Matsumoto K, Ramsey MJ, Freeman DC: **Persistence of chromosome aberrations following acute radiation: I, PAINT translocations, dicentrics, rings, fragments, and insertions.** *Environ Mol Mutagen* 2005, **45**:229-248.
22. Tucker JD, Cofield J, Matsumoto K, Ramsey MJ, Freeman DC: **Persistence of chromosome aberrations following acute radiation: II, does it matter how translocations are scored?** *Environ Mol Mutagen* 2005, **45**:249-257.
23. Tucker JD, Tawn EJ, Holdsworth D, Morris S, Langlois R, Ramsey MJ, Kato P, Boice JD Jr, Tarone RE, Jensen RH: **Biological dosimetry of radiation workers at the Sellafield nuclear facility.** *Radiat Res* 1997, **148**:216-226.
24. Tucker JD, Sorensen KJ, Chu CS, Nelson DO, Ramsey MJ, Orlando C, Heddl JA: **The accumulation of chromosome aberrations and Dlb-1 mutations in mice with highly fractionated exposure to gamma radiation.** *Mutat Res* 1998, **400**:321-335.
25. Tucker JD: **Low-dose ionizing radiation and chromosome translocations: a review of the major considerations for human biological dosimetry.** *Mutat Res* 2008, **659**:211-220.
26. Frohling S, Dohner H: **Chromosomal abnormalities in cancer.** *N Engl J Med* 2008, **359**:722-734.
27. McKernan LT, Ruder AM, Petersen MR, Hein MJ, Forrester CL, Sanderson WT, Ashley DL, Butler MA: **Biological exposure assessment to tetrachloroethylene for workers in the dry cleaning industry.** *Environ Health* 2008, **7**:12.
28. Torasson M, Butler MA, Ruder A, Forrester C, Taylor L, Ashley DL, Mathias P, Marlow KL, Cheever KL, Krieg E, Wey H: **Effect of perchloroethylene, smoking, and race on oxidative DNA damage in female dry cleaners.** *Mutat Res* 2003, **539**:9-18.
29. National Institute for Occupational Safety and Health: **Manual of Analytical Methods (NMAM®), DHHS (NIOSH) Publication 94-113, 1st Supplement Publication 96-135, 2nd Supplement Publication 98-119, 3rd Supplement 2003-154.** Edited by: Schlecht PC, O'Connor PF, 4 1994.
30. National Institute for Occupational Safety and Health: **Tetrachloroethylene (portable GC) in exhaled breath and air #3704.** In *NIOSH Manual of Analytical Methods*. Edited by: Eller PM, Cassinelli ME. Cincinnati, Ohio: DHHS/PHS/CDC/NIOSH; 1998:1-7.
31. Ashley DL, Bonin MA, Cardinali FL, McCraw JM, Holler JS, Needham LL, Patterson DG Jr: **Determining volatile organic compounds in human blood from a large sample population by using purge and trap gas chromatography/mass spectrometry.** *Anal Chem* 1992, **64**:1021-1029.
32. Matsumoto K, Ramsey MJ, Nelson DO, Tucker JD: **Persistence of radiation-induced translocations in human peripheral blood determined by chromosome painting.** *Radiation Research* 1998, **149**:602-613.
33. Tucker JD, Morgan WF, Awa AA, Bauchinger M, Blakey D, Cornforth MN, Littlefield LG, Natarajan AT, Shasserre C: **A proposed system for scoring structural aberrations detected by chromosome painting.** *Cytogenet Cell Genet* 1995, **68**:211-221.
34. Sigurdson AJ, Ha M, Hauptmann M, Bhatti P, Sram RJ, Beskid O, Tawn EJ, Whitehouse CA, Lindholm C, Nakano M, Kodama Y, Nakamura N, Vorobtsova I, Oestreicher U, Stephan G, Yong LC, Bauchinger M, Schmid E, Chung HW, Darroudi F, Roy L, Voisin P, Barquinero JF, Livingston G, Blakey D, Hayata I, Zhang W, Wang C, Bennett LM, Littlefield LG, Edwards AA, Kleinerman RA, Tucker JD: **International study of factors affecting human chromosome translocations.** *Mutat Res* 2008, **652**:112-121.

doi:10.1186/1476-069X-10-16

Cite this article as: Tucker *et al.*: **Cytogenetic analysis of an exposed-referent study: perchloroethylene-exposed dry cleaners compared to unexposed laundry workers.** *Environmental Health* 2011 **10**:16.

Submit your next manuscript to BioMed Central and take full advantage of:

- **Convenient online submission**
- **Thorough peer review**
- **No space constraints or color figure charges**
- **Immediate publication on acceptance**
- **Inclusion in PubMed, CAS, Scopus and Google Scholar**
- **Research which is freely available for redistribution**

Submit your manuscript at
www.biomedcentral.com/submit

