

Glutathione enzyme and selenoprotein polymorphisms associate with mercury biomarker levels in Michigan dental professionals

Jaclyn M. Goodrich ^a, Yi Wang ^a, Brenda Gillespie ^b, Robert Werner ^{a,c}, Alfred Franzblau ^a, Niladri Basu ^{a,*}

^a Department of Environmental Health Sciences, University of Michigan School of Public Health, 1415 Washington Heights, Ann Arbor, MI 48109, USA

^b Department of Biostatistics, University of Michigan School of Public Health, 1415 Washington Heights, Ann Arbor, MI 48109, USA

^c Department of Physical Medicine and Rehabilitation, University of Michigan, 325 E. Eisenhower Parkway Suite 100, Ann Arbor, MI 48108, USA

ARTICLE INFO

Article history:

Received 28 June 2011

Revised 15 September 2011

Accepted 16 September 2011

Available online 24 September 2011

Keywords:

Elemental mercury

Methylmercury

Polymorphism

Gene-environment

Selenoprotein

Glutathione s-transferase

ABSTRACT

Mercury is a potent toxicant of concern to both the general public and occupationally exposed workers (e.g., dentists). Recent studies suggest that several genes mediating the toxicokinetics of mercury are polymorphic in humans and may influence inter-individual variability in mercury accumulation. This work hypothesizes that polymorphisms in key glutathione synthesizing enzyme, glutathione s-transferase, and selenoprotein genes underlie inter-individual differences in mercury body burden as assessed by analytical mercury measurement in urine and hair, biomarkers of elemental mercury and methylmercury, respectively. Urine and hair samples were collected from a population of dental professionals ($n=515$), and total mercury content was measured. Average urine ($1.06 \pm 1.24 \text{ ug/L}$) and hair mercury levels ($0.49 \pm 0.63 \text{ ug/g}$) were similar to national U.S. population averages. Taqman assays were used to genotype DNA from buccal swab samples at 15 polymorphic sites in genes implicated in mercury metabolism. Linear regression modeling assessed the ability of polymorphisms to modify the relationship between mercury biomarker levels and exposure sources (e.g., amalgams, fish consumption). Five polymorphisms were significantly associated with urine mercury levels (*GSTT1* deletion), hair mercury levels (*GSTP1-105*, *GSTP1-114*, *GSS 5'*), or both (*SEPP1 3'UTR*). Overall, this study suggests that polymorphisms in selenoproteins and glutathione-related genes may influence elimination of mercury in the urine and hair or mercury retention following exposures to elemental mercury (via dental amalgams) and methylmercury (via fish consumption).

© 2011 Elsevier Inc. All rights reserved.

Introduction

Mercury toxicokinetics and toxicodynamics are largely influenced by the chemical form of the metal. Though humans are exposed to mercury through a variety of sources (e.g. vaccines, thermostats, compact fluorescent lights), the main sources of elemental mercury (inorganic) and methylmercury (organic) exposure come from dental amalgams and fish consumption, respectively (ATSDR, 1999; Clarkson and Magos, 2006). Urine is commonly used to measure elemental mercury exposure and blood or hair to measure methylmercury exposure (Berglund et al., 2005; Clarkson et al., 2007). However, each biomarker shows tremendous inter-individual variation, and this complicates risk assessment. For

example, mercury typically accumulates 250 times more in the hair than it does in the blood, though this ratio ranges widely among individuals (ATSDR, 1999; Berglund et al., 2005). The half-life of methylmercury in hair ranges from 33 to 120 days (Birke et al., 1972), and the half-life of inhaled elemental mercury ranges from 35 to 90 days in the whole body (Hersh et al., 1976). Failure to understand such inter-individual variation in the accumulation of mercury, its distribution to various biomarkers and its half-life in the body may lead to risk assessment decisions that fail to protect the most susceptible individuals.

Inter-individual variation in mercury biomarkers may be partly explained by genetic factors. For example, mercury is eliminated following conjugation with glutathione, a process that may involve glutathione s-transferases (GSTs; GST theta 1, GSTT1; GST mu 1, GSTM1; GST pi 1, GSTP1) and indirectly depends on enzymes of the glutathione synthesis and degradation pathways (glutamate-cysteine ligase catalytic and modifier subunits, GCLC, GCLM; glutathione synthetase, GSS; glutathione reductase, GSR; γ -glutamyltransferase 1, GGT1) (Ballatori and Clarkson, 1985; Rooney 2007). Selenoproteins (e.g. selenoprotein pi 1, SEPP1; glutathione peroxidases, GPX) combat the oxidative stress created by mercury and bind the toxicant directly

Abbreviations: CRM, certified reference material; GST, glutathione s-transferase; Hg, mercury; MAF, minor allele frequency; MDA, Michigan Dental Association; NHANES, National Health and Nutrition Examination Survey; ln, natural log; SNP, single nucleotide polymorphism; UTR, untranslated region.

* Corresponding author at: 6634-1415 Washington Heights, Ann Arbor, MI, 48109, USA. Fax: +1 734 936 7283.

E-mail address: niladri@umich.edu (N. Basu).

via selenocysteine residues (Chen et al., 2006). Genetic polymorphisms are ubiquitous among these glutathione and selenoprotein genes that facilitate the distribution, metabolism and elimination of mercury. As documented in recent epidemiological studies, such polymorphisms can influence mercury biomarker levels, which may ultimately improve understanding of inter-individual variability following exposure to methylmercury (Custodio et al., 2004; Engström et al., 2008; Gundacker et al., 2007, 2009) and elemental mercury (Custodio et al. 2005; Gundacker et al., 2009).

The aforementioned studies provide early evidence that genetic polymorphisms in key mercury handling genes may underlie differences in mercury distribution and accumulation in exposure biomarkers. However, discrepancies exist among the studies (see Table 1 for more details, including results of this study). For example, Custodio et al. (2004) found the minor allele (T) of the *GSTP1*-114 polymorphism (rs1138272) to be associated with increased blood mercury levels while Engström et al. (2008) found the opposite in subjects with higher fish consumption. Disparate findings may be related to differences between the studies in terms of biomarkers examined, statistical methodologies utilized, sources of mercury exposure, and polymorphism frequency. Here we aim to address these limitations and help improve understanding of gene-environment interactions influencing mercury biomarker levels by studying both forms of mercury (through analytical measurement of biomarkers in urine and hair), various sources of exposure (dental amalgams and fish as determined from surveys), and additional polymorphisms in glutathione-related (*GSTT1*, *GSTM1*, *GSTP1*, *GSTM3*, *GGT1*, *GSS*, *GSR*, *GCLC*, *GCLM*) and selenoprotein (*SEPP1*, *GPX1*, *GPX4*) genes in a group of dental professionals that are exposed to mercury via occupational and environmental routes. Polymorphisms in the listed genes, which are important in the metabolism and elimination of mercury, are hypothesized to affect individual susceptibility to elemental mercury and/or methylmercury accumulation in the body as assessed by urine and hair biomarkers.

Materials and methods

Study population. A convenience sample of 515 dental professionals (dentists, hygienists, and other related professions) was recruited during the 2009 (n = 231) and 2010 (n = 284) Michigan Dental Association (MDA) Annual Conventions. Institutional Review Board (IRB) approval was obtained for this study from the University of Michigan (HUM00027621).

Survey data. For each participant a self-administered survey provided information on subject demographics (e.g., age, gender, weight), occupational practices (e.g., hours worked per week, number of amalgams removed and placed per week), and personal exposures to elemental mercury (amalgams in subject's mouth) and methylmercury (fish consumption). Subjects provided detailed information on fish consumption (e.g., portion size, type of fish, monthly consumption). Fish data were used to estimate a mercury intake variable (ug per kg body weight per day) using mercury concentration data from the U.S. Food and Drug Administration for each fish species (Bahnik et al., 1994; Mierzykowski and Carr, 2001; U.S. FDA).

Mercury exposure assessment. Spot urine samples, biomarkers of elemental mercury exposure, (>30 mL) were collected from each participant and stored frozen. Hair samples, biomarkers for methylmercury, (>10 mg) were cut from the occipital region of the scalp, and the first 2 cm of hair (proximal end) were used for measurement. Total mercury levels in urine and hair were measured using a direct mercury analyzer (DMA-80, Milestone Inc., CT) according to U.S. EPA Method 7473 as previously described (Basu et al., 2010; Paruchuri et al., 2010). Quality control measures included incorporating one blank, one replicate sample, and a certified reference material (CRM; hair: NIES Japan CRM#13; two urine CRMs: QMEQAS08U-01 and ClinChek; dogfish liver: DOLT, National

Table 1

Summary of results in studies assessing the association of genetic polymorphisms with mercury biomarker levels.

Reference	Population	n	Hg biomarkers	Exposure sources	Polymorphisms assessed	Polymorphisms significantly associated with higher biomarker Hg levels	Polymorphisms significantly associated with lower biomarker Hg levels
This study	Michigan dental professionals	208 to 482	Hair, urine	Dental amalgam, fish	See Table 3	Positive genotype by amalgam interaction in urine Hg: <i>SEPP1</i> 3'UTR CT/TT; positive genotype by fish Hg interaction in hair Hg model: <i>GSS</i> 5' CG/GG	Ln(hair Hg) and <i>GSTP1</i> -105 G; Ln (urine Hg): <i>GSTT1</i> −/−; negative main effect in urine Hg model: <i>SEPP1</i> 3'UTR T and <i>GSTT1</i> −/−; negative main effect in hair Hg model: <i>GSTP1</i> -105 G and <i>GSTP1</i> -114 T; negative genotype by fish Hg interaction in hair Hg model: <i>SEPP1</i> 3'UTR T
Custodio et al. (2004)	Swedish adults	365	Erythrocytes	Fish	<i>GCLC</i> (rs17883901), <i>GCLM</i> (rs41303970), <i>GSTP1</i> -105, −114 (rs1695, rs1138272), <i>GSTA1</i> (rs3957396), <i>GSTT1</i> and <i>GSTM1</i> deletion	<i>GCLC</i> T and/or <i>GSTP1</i> -114 T carriers had steeper regression of erythrocyte Hg on polyunsaturated fatty acids	
Custodio et al. (2005)	Ecuadorian gold miners and buyers	309	whole blood, plasma, urine	gold mining practices	<i>GCLC</i> , <i>GCLM</i> , <i>GSTP1</i> -105, <i>GSTP1</i> -114, <i>GSTA1</i> , <i>GSTT1</i> and <i>GSTM1</i> deletion	<i>GCLM</i> T and elevated blood, plasma, urine Hg in moderately exposed miners	
Gundacker et al. (2007)	Austrian students	174 to 222	whole blood, urine, hair	fish	<i>GSTT1</i> and <i>GSTM1</i> deletion	<i>GSTT1</i> /M1 deletions and hair Hg	
Engström et al. (2008)	Swedish adults	292	erythrocytes	fish	<i>GCLC</i> , <i>GCLM</i> , <i>GSTP1</i> -105, <i>GSTP1</i> -114	<i>GCLM</i> TT and erythrocyte Hg (adjusted for polyunsaturated fatty acids)	
Gundacker et al. (2009)	Austrian students	184 to 304	whole blood, hair, urine	dental amalgam, fish	<i>GCLC</i> , <i>GSTP1</i> -105, <i>GSTP1</i> -114, <i>GSTA1</i> , <i>GSTT1</i> and <i>GSTM1</i> deletion, metallothionein SNPs	<i>GSTP1</i> -105 G and unadjusted median blood and hair Hg, in CATREG analysis on dichotomized Hg levels: <i>GSTP1</i> -114 T and/or <i>GCLC</i> T	
						<i>GSTA1</i> A and unadjusted median blood Hg	

Research Council Canada) in every batch of 10–15 samples. All samples exceeded the theoretical method detection limits, defined as $3 \times$ SD of blanks, (0.03 ug/L mercury for urine and 0.01 ug/L mercury for hair). Mean (\pm SD) recovery of mercury from CRMs was $94.4 \pm 15.4\%$. Mean within day ($3.1 \pm 1.7\%$) and between day ($6.7 \pm 6.8\%$) variability of CRMs were similar to that of duplicated subject samples.

Genotype analyses. To obtain DNA from buccal cells, subjects swabbed the inside of their cheeks with four cotton swabs for 15 s each. Cotton swabs were stored at room temperature in 0.5 mL of buffer (100 mM sodium chloride, 10 mM Tris base, 10 mM EDTA, 0.5% sodium dodecyl sulfate with 0.1 mg/mL of Proteinase K, according to [Min et al., 2006](#)). DNA was purified from buccal cells using Promega SV Wizard Genomic DNA purification kits (Promega Corporation, Madison, WI), and DNA quantification was performed using PicoGreen reagent with relative fluorescence measured by a Perkin Elmer HTS 7000 Plus Bioassay Reader (Shelton, CT).

Over the two study years, fifteen polymorphisms were genotyped in glutathione s-transferases, glutathione synthesizing enzymes, and selenoproteins, but in a given study year ten or eleven polymorphisms were genotyped. Fourteen polymorphisms (rs1695, rs1138272, rs3877899, rs713041, rs7579, rs17883901, rs41303970, rs1050450, rs5751901, rs7483, rs3761144, rs1002149, rs2911678, *GSTT1* deletion) were genotyped using Taqman Allelic Discrimination Assays (Applied Biosystems; [Livak et al., 1995](#)). Results were read and interpreted using an ABI 7700 (Carlsbad, CA). All primers and probes were pre-designed by Applied Biosystems except for the *GSTT1* deletion genotyping which utilized primers and probes designed by [Mordukhovich et al. \(2009\)](#). The *GSTM1* deletion polymorphism was genotyped in the 2009 samples using the PCR-based method of [Lee et al. \(2009\)](#). *GSTT1* and *GSTM1* genotyping resulted in two distinct groups: deletion homozygotes and individuals with at least one intact copy of the gene.

Statistical analysis. All statistical analyses were performed using PASW Statistics v18 (Chicago, IL). The 2009 and 2010 data were grouped together given that demographics, mercury biomarker levels and exposures had similar distributions both years. In all analyses, mercury values were natural log-transformed to achieve normality. The influence of genetic polymorphisms, assuming additivity of variant alleles, on mercury biomarker levels (hair or urine) was studied via simple linear regression models with one ordinal variable per polymorphism (coded 1 = major homozygote, 2 = heterozygote, 3 = minor homozygote) in each model. The association of genetics with urine mercury and hair mercury biomarkers adjusted for predictors was also assessed by multiple linear regression. Base models without genetic variables were first created. All variables (e.g., demographics, fish consumption, occupational exposures) were included in the base model, and insignificant variables ($p > 0.05$) were removed via a manual backward selection process. Final base models consisted of significant predictors of hair or urine mercury ($p < 0.05$ for parameter estimate) unless otherwise noted. The effect of genotype (main effect and interaction with each exposure predictor variable) was then assessed with separate models for each polymorphism. Models with separate dummy variables for heterozygotes and minor homozygotes and interaction terms for each were also run (data not shown). The effect estimates (magnitude, direction, significance) for genotype obtained from this method, which did not assume genetic additivity, were similar to results reported here.

According to the method of [Levine and Fahy \(1945\)](#), which has been shown to reduce variability of metal excretion associated with spot urine samples, urine mercury levels were adjusted for the mean specific gravity (1.017) of the study population ([Mason and Calder, 1994](#)). The linear regression models described were run

with unadjusted and adjusted urine. Parameter estimates did not notably change and significance only changed in a few instances in specific gravity adjusted models. Unadjusted models are reported here unless otherwise noted. In all urine mercury models, eight subjects were excluded due to a history of kidney disease (lithiasis, pyelonephritis, and/or orthostatic proteinuria) that may have influenced urinary excretion of mercury.

Results

Descriptive statistics

The majority of the 515 participants provided urine ($n = 502$) and hair ($n = 505$). Subjects lacking both biomarker samples and demographic information were excluded ($n = 4$). Sixty-one percent of all subjects were female, and the majority of the population ($>90\%$) self-identified racially as Caucasian. [Table 2](#) provides average (\pm SD) demographic and occupational variables for the total population and occupation-stratified population (dentists, $n = 243$ vs. non-dentists, $n = 268$). Genders were unequally distributed in the occupational groups as 24% of dentists were female compared to 95% of non-dentists (e.g., dental hygienists, dental assistants). Stratification of subjects into two occupational groups yielded significant differences ($p < 0.05$) between the groups. On average, dentists were older, removed and placed more dental amalgams in the office, and had more amalgams in their own mouths compared with non-dentists.

Mercury biomarkers

Average mercury biomarker levels are reported in [Table 2](#). Urine mercury (mean \pm SD, 1.04 ± 1.18 ug/L; median 0.66 ug/L) ranged from 0.03 to 9.26 ug/L with a distribution resembling that of the U.S. population according to the National Health and Nutrition Examination Survey (NHANES 2003–2004, median 0.48 ug/L; [CDC, 2009](#)). Dentists had significantly higher urine mercury levels compared to non-dentists (1.37 ± 1.3 ug/L vs. 0.75 ± 0.97 ug/L; $p < 0.001$ for ANOVA comparing natural log-transformed values). The distribution of hair mercury (0.49 ± 0.63 ug/g; range 0.02 to 6.15 ug/g) also resembled NHANES data from 1999 to 2000 (mean 0.47 ug/g; [McDowell et al., 2004](#)). Dentists had significantly higher hair mercury levels compared to non-dentists (0.69 ± 0.81 vs. 0.31 ± 0.33 ug/g; $p < 0.001$) even though the estimated mercury intake from fish consumption was not significantly different between the two groups.

Table 2
Mean (SD) of key demographic and exposure variables in total population and occupation stratified populations.

	Total population	Occupation stratified	
		Dentists	Non-dentists
n	511	243	268
% female	61.3	24.3	94.8
BMI (kg/m ²)	26.4 (4.7)	26.4 (4.0)	26.5 (5.3)
Age (years)	52.1 (12.1)	56.1 (11.6)	48.3 (11.2) ^b
Hours worked/week (in past 6 months)	27.9 (11.6)	29.1 (11.1)	27.0 (11.9) ^a
# Amalgam (personal)	4.1 (4.1)	4.6 (4.2)	3.7 (3.9) ^a
# Amalgam placed/week	12.1 (25.8)	20.4 (31.1)	4.7 (16.7) ^b
# Amalgam removed/week	13.3 (23.1)	23.2 (27.2)	4.4 (13.4) ^b
Hg Intake (ug/kg body weight/day)	0.08 (0.13)	0.09 (0.14)	0.08 (0.12)
Urine Hg (ug/L)	1.04 (1.18)	1.37 (1.3)	0.75 (0.97) ^b
Hair Hg (ug/g)	0.49 (0.63)	0.69 (0.81)	0.31 (0.33) ^b

^{a,b} denote p-value <0.05 and <0.001 , respectively, for ANOVA test comparing dentists and non-dentists. ANOVA comparing urine and hair Hg used natural log-transformed values.

Table 3

Genotype frequencies of 15 polymorphisms in glutathione S-transferases, glutathione synthesizing enzymes, and selenoproteins and simple linear regression assessing the influence of genotype on mean urine and hair Hg levels (natural log-transformed). Significant ($p < 0.05$) β estimates (95% CI) for genotype are reported from models of urine Hg or hair Hg.

Gene	db SNP ID	Alleles	SNP Details	n	Minor allele frequency	Urine Hg	Hair Hg
<i>GSTP1</i>	rs1695	A>G	105 Ile>Val	485 ^a	0.32		-0.22 (-0.36, -0.07)
<i>GSTP1</i>	rs1138272	C>T	114 Ala>Val	475 ^a	0.08		
<i>GCLM</i>	rs41303970	C>T	5' near gene	481 ^a	0.19		
<i>SEPP1</i>	rs3877899	C>T	234 Ala>Thr	485 ^a	0.24		
<i>SEPP1</i>	rs7579	C>T	3' UTR	479 ^a	0.28	-0.16 (-0.3, -0.02)	
<i>GSTT1</i>	N/A	+>−	Deletion	437 ^a	0.41	-0.27 (-0.52, -0.02)	
<i>GPX1</i>	rs1050450	C>T	200 Pro>Leu	223 ^b	0.28		
<i>GPX4</i>	rs713041	C>T	3' UTR	226 ^b	0.45		
<i>GCLC</i>	rs17883901	G>A	5' near gene	230 ^b	0.08		
<i>GSTM1</i>	N/A	->+	Deletion	209 ^b	0.32		
<i>GGT1</i>	rs5751901	T>C	intron	251 ^c	0.36		
<i>GSTM3</i>	rs7483	G>A	224 Val>Ile	243 ^c	0.26		
<i>GSS</i>	rs3761144	C>G	5' near gene	255 ^c	0.40		
<i>GSR</i>	rs1002149	G>T	5' near gene	240 ^c	0.18		
<i>GSR</i>	rs2911678	T>A	intron	250 ^c	0.24		

^a Genotyped in subjects recruited in 2009 and 2010.

^b Genotyped in subjects recruited in 2009 only.

^c Genotyped in subjects recruited in 2010 only.

Genotype and mercury biomarkers

All genotyped polymorphisms achieved Hardy–Weinburg Equilibrium. Minor allele frequencies (MAF) for each polymorphism were generally similar to HapMap reference Caucasian population MAFs (Table 3). In statistical models comparing mercury levels among genotype groups, all polymorphisms were divided into major homozygote, heterozygote, and minor homozygote groups with the exception of *GSTM1* and *GSTT1* deletion polymorphisms which were divided into two genotype groups (double deletion vs. at least one intact copy).

Simple linear regression models were used to test the differences in mean natural log-transformed urine (ug/L) or hair (ug/g) mercury levels among genotype groups for each polymorphism. Three statistically significant findings emerged (β estimate (95% CI) reported in Table 3). Urine levels were lower among individuals with double deletion of *GSTT1* compared to those with the intact gene (-0.27 (95% CI: -0.52, -0.02) ug/L change in ln(urine Hg) with *GSTT1* deletion). Likewise, individuals with the T allele for *SEPP1* 3'UTR (rs7579) had lower urine mercury levels (-0.16 (95% CI: -0.3, -0.02) ug/L change in ln(urine Hg) with each T allele). Hair mercury levels differed by *GSTP1*-105 (rs1695) genotype with minor homozygotes (GG) exhibiting the lowest hair mercury measurements (-0.22 (-0.36, -0.07) ug/g change in ln(hair Hg) with each G allele).

Linear regression modeling of urine biomarker

Linear regression was used to model natural log-transformed urine mercury levels with significant exposure variables. In the final base model, significant predictors were number of dental amalgams in the subjects' mouth ('amalgam'), amalgams removed or placed per week in the dental office ('amalgam handled', an ordinal variable), and occupation (dentist vs. non-dentist) (Table 4). The categories for 'amalgam handled' were defined as follows: reference (0 amalgams handled), low (1 to 20 per week), medium (21 to 45), and high (>45). The effect of genotype for each polymorphism was tested (main effect, interactions between genotype and the three exposure variables). For all models, only the interactions between genotype and 'amalgam' are reported here as amalgams were the dominant predictor of urine Hg levels. The number of subjects in each model ranged from 204 to 466 depending on the polymorphism; not all polymorphisms were genotyped both years, and some subjects did not provide adequate DNA to complete the genotyping.

Urine mercury models with nominally significant ($p < 0.05$) genotype (main effect or interaction) terms are reported in Table 4 (see Supplemental Table 1 for all urine models). Significant main effects of genotype were observed for *SEPP1* 3'UTR (β estimate (95% CI) = -0.26 (-0.43, -0.08) ug/L) and *GSTT1* deletion (-0.25 (-0.5, -0.00) ug/L), though the latter was only observed when urine was

Table 4

Parameter estimates from models of natural log-transformed urine Hg levels (unadjusted and adjusted for specific gravity) with exposure predictor variables, genotype, and genotype-by-amalgam interactions. Models with polymorphisms that have significant genotype variable(s) in at least one urine model are reported. Beta coefficients (95% CI) are listed for each predictor variable. Statistically significant beta coefficients ($p < 0.05$) are in bold.

SNP	n ^a	Adj. R ²	Amalgam	Amalgam handled ^b	Non-dentist	Genotype	Genotype * amalgam
Base model	492	0.23	0.08 (0.06, 0.1)	0.12 (0.04, 0.21)	-0.45 (-0.64, -0.25)		
<i>SEPP1</i> 3'UTR	240, 186, 35	0.25	0.03 (-0.03, 0.08)	0.11 (0.03, 0.2)	-0.44 (-0.64, -0.24)	-0.26 (-0.43, -0.08)	0.04 (0.01, 0.07)
<i>GSTT1</i> deletion	347, 71	0.24	0.08 (0.02, 0.15)	0.12 (0.03, 0.21)	-0.4 (-0.6, -0.19)	-0.18 (-0.48, 0.12)	0 (-0.06, 0.05)
<i>Sp. gravity adjusted urine</i>							
Base model	491	0.22	0.07 (0.06, 0.09)	0.09 (0.02, 0.16)	-0.28 (-0.44, -0.12)		
<i>SEPP1</i> 3'UTR	240, 186, 35	0.23	0.05 (0.01, 0.09)	0.08 (0.01, 0.15)	-0.29 (-0.45, -0.12)	-0.09 (-0.24, 0.05)	0.02 (-0.01, 0.04)
<i>GSTT1</i> deletion	347, 71	0.23	0.07 (0.02, 0.13)	0.08 (0.01, 0.16)	-0.23 (-0.4, -0.07)	-0.25 (-0.5, -0.00)	0 (-0.04, 0.05)

^a Number of subjects in model in each genotype group. Genotype groups are major homozygote, heterozygote, minor homozygote for *SEPP1* 3'UTR and gene present vs. double deletion for *GSTT1*.

^b Ordinal variable for number of amalgams removed and placed per week (see text for category cut-offs).

Table 5

Parameter estimates from models of natural log-transformed hair Hg levels with exposure variable (estimated mercury intake from fish consumption, ug per kg body weight per day), genotype, and genotype-by-fish Hg interactions. Models with significant genotype variables are reported. Beta coefficients (95% CI) are reported for each predictor variable. Statistically significant beta coefficients ($p < 0.05$) are in bold.

SNP	n ^a	Adj. R ²	Fish Hg	Genotype	Genotype*fish Hg
Base model	503	0.19	3.62 (2.96, 4.28)		
<i>GSTP1-105</i>	214, 216, 44	0.2	2.87 (0.95, 4.8)	−0.22 (−0.38, −0.06)	0.42 (−0.73, 1.57)
<i>GSTP1-114</i>	393, 69, 2	0.2	2.91 (0.74, 5.07)	−0.31 (−0.6, −0.03)	0.67 (−1.05, 2.39)
<i>SEPP1 3'UTR</i>	245, 188, 36	0.19	6.22 (4.14, 8.29)	0.15 (−0.02, 0.32)	−1.61 (−2.82, −0.4)
<i>GSS 5'</i>	94, 111, 44	0.15	0.34 (−2.13, 2.81)	−0.15 (−0.35, 0.06)	1.88 (0.44, 3.31)

^a Number of subjects in model in each genotype group. Genotype groups are major homozygote, heterozygote, minor homozygote.

first adjusted for specific gravity. A significant interaction was observed with *SEPP1 3'UTR* T alleles and personal amalgams (β (95% CI) = 0.04 (0.01, 0.07) ug/L), though the interactions with amalgam handled and occupation were not significant (data not shown).

Linear regression modeling of hair biomarker

Hair mercury levels (ln-transformed) were modeled using statistical methods similar to urine mercury modeling. The final base model included the most significant predictor of hair Hg, 'fish Hg' (calculated ug Hg intake per kg body weight per day from fish consumption, β (95% CI) = 3.62 (2.96, 4.28) ug/g, see Table 5). Addition of genotype into the model resulted in two genotypes with significant main effects (*GSTP1-105*, rs1695, β (95% CI) = −0.22 (−0.38, −0.06) ug/g; *GSTP1-114*, rs1138272, −0.31 (−0.6, −0.03) ug/g). Significant interactions between genotype and fish Hg were also observed in models with *SEPP1 3'UTR* and *GSS 5'* (rs3761144). Individuals with the minor allele (T) of *SEPP1 3'UTR* accumulated less mercury in the hair per intake from fish consumption (β (95% CI) = −1.61 (−2.82, −0.4) ug/g change in ln(hair Hg) with each T allele). The opposite relationship was observed with the minor allele of *GSS 5'* (G) and fish Hg (1.88 (0.44, 3.31) ug/g change in ln(hair Hg) per G allele). To highlight the nature of these results, two scatterplots of hair mercury by

mercury intake from fish (both ln-transformed) are shown stratified by *SEPP1 3'UTR* genotype (Fig. 1A) and *GSS 5'* genotype (Fig. 1B). Significant results (main effect or genotype-by-exposure variable interactions) were not observed for any other polymorphism in the models of urine or hair mercury (see Supplemental Tables 1 and 2 for all urine and hair models).

Discussion

Recent epidemiological studies suggest that seven genetic polymorphisms in GSTs and glutathione synthesizing enzymes may influence mercury accumulation in common biomarkers (hair, blood, urine) following exposure to elemental mercury or methylmercury (Custodio et al., 2004, 2005; Engström et al., 2008; Gundacker et al., 2007, 2009). Here, we build upon these studies and assess the influence of fifteen polymorphisms in GSTs, glutathione synthesizing enzymes and selenoproteins on the accumulation of both elemental mercury and methylmercury in urine and hair biomarkers, respectively, in a population of occupationally and environmentally exposed dental professionals. Our results suggest that five polymorphisms may affect accumulation of mercury in urine (*GSTT1* deletion), hair (*GSTP1-105*, *GSTP1-114*, *GSS 5'*) or both (*SEPP1 3'UTR*).

Four nominally significant ($p < 0.05$) polymorphism-mercury biomarker relationships were observed that have not been reported previously, including two SNPs (*GSS 5'*, *SEPP1 3'UTR*) genotyped for the first time in a mercury-gene study. A deletion polymorphism in *GSTT1* was associated with decreased urine mercury in both simple (Table 3) and multivariate linear regression (Table 4), though the latter was significant only after adjusting for specific gravity. Effect modification of *SEPP1 3'UTR* genotype on the relationship between exposure source and mercury biomarker was observed in both urine and hair mercury models. The minor allele, T, of the *SEPP1 3'UTR* SNP was linked to lower urine mercury levels on average (Table 3). However, upon adjusting for elemental mercury exposures, each T allele was associated with greater urine mercury accumulation per amalgam (Table 4). In hair mercury models, *SEPP1 3'UTR* T was associated with lower hair mercury per unit of intake from fish consumption (Fig. 1A, Table 5). Contrariwise, the *GSS 5'* minor allele was associated with increasing hair mercury concentration per unit of fish mercury (Fig. 1B, Table 5). This study also reports associations between minor alleles of two SNPs, *GSTP1-105* and *GSTP1-114*, and decreased hair mercury concentrations though significant effect modification on the relationship between mercury intake from fish consumption and hair levels were not observed. Previous gene-mercury biomarker studies have assessed the same *GSTP1* SNPs with conflicting results (Custodio et al., 2004; Engström et al., 2008; Gundacker et al., 2009).

Dental professionals provide a convenient sample to assess mercury-gene interactions as they share common exposure routes with the general population (e.g. fish consumption, personal amalgams) along with an additional occupational exposure. While dentists' occupational exposures are typically higher than the average population, a shift toward use of composite resin fillings and safer handling of mercury may result in dentists having urine biomarker levels comparable




Fig. 1. Modification of the relationship between Hg intake from fish consumption and Hair Hg levels by A) *SEPP1 3'UTR* (C>T) genotype and B) *GSS 5'* (C>G) genotype. Note: hair Hg and Hg intake are natural log-transformed, though axes are labeled with untransformed units for ease of interpretation.

to that of the general U.S. population (Eklund, 2010). Here, participating dentists and other related professionals (e.g., dental hygienists, dental assistants) exhibited urine and hair mercury levels similar to that of the general U.S. population according to NHANES (CDC, 2009; McDowell et al., 2004). The best predictors of urine mercury levels were personal dental amalgams and occupational exposure to amalgams with dentists having the greatest exposure and biomarker levels. Study subjects displayed a wide range of methylmercury exposure from fish consumption, enabling the study of both mercury forms.

Immense variability in mercury biomarker measurements and half-lives has been observed in human populations, complicating risk assessment of both elemental and methylmercury (Berglund et al., 2005; Birke et al., 1972; Hursh et al., 1976). Differential metabolism of mercury in the body could affect its half-life and reactivity, as conjugated mercury is less able to induce oxidative stress, bind to thiol groups, and/or inhibit proteins in the body. Functional GSTs, adequate glutathione and selenoproteins are essential for protection against mercury (Ballatori and Clarkson, 1985; Chen et al., 2006; Clarkson and Magos, 2006). Polymorphisms are ubiquitous among the genes encoding these mercury-handling proteins and have the potential to modify mercury toxicokinetics, reflected partially in biomarker levels, via altered enzyme activity or gene expression.

Previous studies assessing the association of genetic polymorphisms with mercury biomarkers focused on seven variants in glutathione synthesizing enzymes and glutathione s-transferases, six of which were genotyped here (see Table 1). Our results follow several trends previously reported, and this study further explores genotype-mercury relationships that were inconsistent in past research. Custodio et al. (2005) found the T allele of *GCLM* rs41303970 to be associated with increased blood, plasma and urine mercury in gold miners occupationally exposed to elemental mercury. Our study did not find associations between urine mercury levels and *GCLM* genotype. Genotype associations may differ based on the level of exposure as the miners had almost seven-fold higher urinary mercury compared to the dental population (miners: median = 4.5 µg/L, maximum = 230 µg/L; dental population: median = 0.7 µg/L, maximum = 9.3 µg/L). Other studies have explored the influence of glutathione-related polymorphisms on urine mercury levels, and a lack of significant outcomes indicate that the polymorphisms studied to date may not influence the variability observed in urinary excretion of elemental mercury. Alternatively, given that the impact of any individual polymorphism on mercury accumulation is expected to be small, studies to date may have insufficient power to detect some associations.

Two previous epidemiological studies have assessed the relationship between glutathione-related polymorphisms and hair mercury levels (Gundacker et al., 2007, 2009), and four explored the modifying effect of polymorphisms on methylmercury accumulation from fish consumption using blood as an exposure biomarker (Custodio et al., 2004; Engström et al., 2008; Gundacker et al., 2007, 2009). Several associative findings linking polymorphisms to hair or blood biomarker levels were inconsistent among studies (Table 1). Gundacker et al. found *GSTM1* and *GSTT1* deletions together (2007) or in combination with other SNPs (2009) to associate with higher hair mercury levels. The present study did not observe significant associations between *GSTM1* or *GSTT1* deletion polymorphisms and hair mercury concentrations when modeling single polymorphisms or combined *GSTT1/GSTM1* genotype (both genes intact vs. at least one deletion).

Conflicting relationships between two *GSTP1* SNPs (105 A>G; 114 C>T) and hair or blood biomarkers were observed among past studies. Given that both SNPs encode amino acid changes in *GSTP1* that decrease enzyme activity (Ali-Osman et al., 1997), these SNPs may alter the ability of *GSTP1* to conjugate glutathione to mercury

or transport mercury-glutathione conjugates. This study suggests that both the *GSTP1*-105 G and *GSTP1*-114 T alleles are linked to decreased hair mercury levels (Table 5). Engström et al. (2008) found that carriers of *GSTP1*-105 G and/or *GSTP1*-114 T accumulated less mercury in erythrocytes per measured polyunsaturated fatty acids (an indicator of fish consumption), suggesting a similar trend as the findings of this study for *GSTP1* variant alleles. In contrast, other studies found associations between *GSTP1*-105 G or *GSTP1*-114 T and higher hair or blood biomarker measurements, after adjustment for fish consumption in many cases and often in combination with other polymorphisms (Custodio et al., 2004; Gundacker et al., 2009). Several factors may influence these incongruent results. While hair and blood, especially erythrocytes, are both biomarkers for methylmercury exposure, 2 cm of scalp hair represent the average exposure over two months with a lag period of approximately one month while blood mercury levels reach a steady-state among consistent fish consumers but fluctuate among infrequent consumers (Berglund et al., 2005). Thus, while similar gene-mercury relationships could be observed in models of hair and blood mercury, they may be influenced by the frequency, duration, and dose of the exposure. Several significant findings in previous studies for *GSTP1*-105 and *GSTP1*-114 involved polymorphism combinations. Our sample size was not large enough to confidently test combinations of all fifteen polymorphisms studied.

Genotype of the *SEPP1* 3'UTR SNP significantly impacted both urine Hg (positive interaction with amalgam) and hair Hg (negative interaction with fish mercury) models. This SNP is found in the 3' untranslated region (UTR), a gene region crucial to the regulation of selenocysteine incorporation. Selenocysteine is an amino acid unique to selenoproteins that can bind mercury-selenium conjugates or methylmercury (Khan and Wang, 2009). *SEPP1* expression depends on genotype and selenium availability and can also be influenced by gender and BMI (Méplan et al., 2007, 2009). The 3'UTR T allele is linked to greater *SEPP1* expression among people supplemented with selenium and higher prevalence of the 60 kDa isoform of *SEPP1* which has more selenocysteine residues and greater mercury binding capacity (Méplan et al., 2007, 2009). Elevated elemental mercury exposure increases *SEPP1* expression and the ability of *SEPP1* to bind mercury as evident in highly exposed miners (Chen et al., 2006). In the dental population, individuals with CT or TT genotype had lower urine mercury levels on average (Table 3). However, the T allele modified the relationship between the source of exposure, dental amalgams, and urine mercury levels. At higher exposure levels, CT/TT individuals excrete more mercury in the urine per amalgam compared to those with the CC genotype (Table 4). The opposite relationship is observed for the hair biomarker whereby each additional T allele is associated with less accumulation of mercury in the hair given methylmercury intake from fish (Fig. 1A). Expression changes of *SEPP1* linked to the T genotype may explain the complex relationship observed between *SEPP1* genotype, mercury biomarker levels, and exposures to mercury that may be complicated further by micronutrient status (e.g. selenium) and chemical speciation of mercury. Higher expression of *SEPP1* from the CT or TT genotypes could affect mercury binding and subsequent distribution to various tissues. Excretion of elemental mercury and methylmercury via urine and hair, respectively, may be differentially modified by this SNP which warrants further investigation and incorporation of selenium biomarker levels into multivariate analysis.

Glutathione synthetase (GSS) is involved in glutathione synthesis, and a SNP upstream of the coding region (5' C>G) may modify the relationship between mercury intake from fish consumption and hair mercury levels. With each variant allele (G), more mercury is accumulated in the hair given intake from fish consumption (Fig. 1B, Table 5). This SNP, which is associated with sporadic amyotrophic lateral sclerosis among individuals with past heavy metal exposure, may influence expression of GSS directly or may be a marker for an

expression-altering SNP (Morahan et al., 2007). Decreased expression of GSS leading to decreased glutathione synthesis could impact the body's ability to eliminate methylmercury as a glutathione conjugate with the higher body burden reflected in hair Hg levels.

We found several nominally significant associations between polymorphisms in glutathione-related enzymes and *SEPP1* and elemental mercury or methylmercury biomarker levels in a population of dental professionals despite several study limitations. This cross-sectional study was dependent upon self-reporting of mercury exposures (fish consumption, number of amalgams handled). Though the voluntary nature of subject participation could lead to selection bias, our subjects did not know their mercury levels or genotype prior to participation in the study, and we obtained a range of mercury biomarker levels similar to that of the general U.S. population (CDC, 2009; McDowell et al., 2004).

In the dually exposed MDA cohort, total Hg in urine was predicted by amalgam exposures and occupation, and hair Hg levels correlated with estimated mercury intake from fish. As such, total Hg in urine and hair were assumed to reflect elemental mercury and methylmercury exposures, respectively. While urine and hair are often used as biomarkers of different mercury species (Berglund et al., 2005), cohorts occupationally exposed to elemental mercury attribute a fraction of hair Hg to the elemental mercury exposure source (Wranová et al., 2008). Likewise, in populations with negligible elemental mercury exposure, urine Hg concentrations correlate with fish consumption (Ohno et al., 2007). While the possibility of such biomarker crossover cannot be eliminated, urine and hair Hg concentrations in the MDA cohort primarily reflect elemental mercury and methylmercury exposures, respectively.

In genotype analyses, the sample size was decreased due to inadequate DNA from a subset of subjects, and not all polymorphisms were genotyped each collection year resulting in sample sizes as small as 204 in some models. Previous gene-mercury biomarker studies found significant impacts of dual polymorphism combinations on mercury biomarker levels. However, due to the sample size and number of loci genotyped, polymorphism combinations could not be tested confidently in this study without encountering errors of multiple testing or insufficient power due to small numbers of cases with some genotype combinations. Without studying polymorphism combinations, we may have missed several significant relationships with mercury biomarkers that would mirror findings in previous studies. Overall, while several significant genotype-mercury biomarker relationships were observed ($p < 0.05$), none remain significant after correction for multiple testing ($p < 0.002$). As such, the nominally significant results reported in the MDA cohort should be considered trends and require further exploration in future cohort studies with greater power.

In conclusion, this study adds to growing knowledge about genetic associations with mercury biomarker levels and inter-individual variability in the toxicokinetics of elemental mercury and methylmercury. Results suggest that five polymorphisms in *GSTP1*, *GSTT1*, *GSS* and *SEPP1* may influence urine and hair mercury levels, though further investigation is warranted. Future research on genetic factors influencing mercury toxicokinetics should assess both major mercury exposure sources (amalgam, fish consumption) in larger cohorts using all three biomarkers (urine, blood, hair) and more polymorphisms in glutathione related enzymes, selenoproteins, and proteins involved in mercury transport to help ascertain sources of biomarker variability and to narrow the gap between the perceived and true health risks accompanying mercury exposure.

Conflict of interest statement

The authors declare that there are no conflicts of interest.

Acknowledgments

We acknowledge the support of Dr. Kevin Sloan and Ms. Andrea Sunderman (Michigan Dental Association). This research was funded by grants from the UM Center for Occupational Health & Safety Engineering's NIOSH Training Grant No. 2T42 OH 008455-04, the Michigan Institute for Clinical and Health Research (MICHR) UL1RR024986, and the UM School of Public Health. JMG was funded through the NIEHS Environmental Toxicology and Epidemiology Training Grant No. T32 007062.

Appendix A. Supplementary data

Supplementary data to this article can be found online at doi:10.1016/j.taap.2011.09.014.

References

- Ali-Osman, F., Akande, O., Antoun, G., Mao, J.-X., Buolamwini, J., 1997. Molecular cloning, characterization, and expression in *Escherichia coli* of full-length cDNAs of three human glutathione S-transferase pi gene variants. *J. Biol. Chem.* 272, 10004–10012.
- ATSDR, 1999. Toxicological Profile for Mercury. U.S. Department of Health and Human Services, Atlanta, GA.
- Ballatori, N., Clarkson, T.W., 1985. Biliary secretion of glutathione and glutathione-metal complexes. *Fundam. Appl. Toxicol.* 5, 816–831.
- Bahnick, D., Sauer, C., Butterworth, B., Kuehl, D., 1994. A national study of mercury contamination of fish. *Chemosphere* 29, 537–546.
- Basu, N., Abare, M., Buchanan, S., Cryderman, D., Nam, D.-H., Sirkin, S., Schmidtt, S., Hu, H., 2010. A combined ecological and epidemiologic investigation of exposure to metals amongst indigenous peoples near the Marlin Mine in Western Guatemala. *Sci. Total. Environ.* 409, 70–77.
- Berglund, M., Lind, B., Björnberg, K.A., Palm, B., Einarsson, O., Vahter, M., 2005. Individual variations of human mercury exposure biomarkers: a cross-sectional assessment. *Environ. Health* 4. doi:10.1186/1476-06X-4-20.
- Birke, G., Johnels, A.G., Plantin, L.O., Sjöstrand, B., Skerfving, S., Westermark, T., 1972. Studies on human exposed to methylmercury through fish consumption. *Arch. Environ. Health* 25, 71–91.
- Centers for Disease Control and Prevention (CDC), 2009. Fourth National Report on Human Exposure to Environmental Chemicals. <http://www.cdc.gov/exposurereport/pdf/FourthReport.pdf> [Accessed 1 November 2010].
- Chen, C., Yu, H., Zhao, J., Li, B., Qu, L., Liu, S., Zhang, P., Chai, Z., 2006. The roles of serum selenium and selenoproteins on mercury toxicity in environmental and occupational exposure. *Environ. Health Perspect.* 114, 297–301.
- Clarkson, T.W., Magos, L., 2006. The toxicology of mercury and its chemical compounds. *Crit. Rev. Toxicol.* 36, 609–662.
- Clarkson, T.W., Vyas, J.B., Ballatori, N., 2007. Mechanisms of mercury disposition in the body. *Am. J. Ind. Med.* 50, 757–764.
- Custodio, H.M., Broberg, K., Wennberg, M., Jansson, J.H., Vessby, B., Hallmans, G., Stegmayr, B., Skerfving, S., 2004. Polymorphisms in glutathione-related genes affect methylmercury retention. *Arch. Environ. Health* 59, 588–595.
- Custodio, H.M., Harari, R., Gerhardsson, L., Skerfving, S., Broberg, K., 2005. Genetic influences on the retention of inorganic mercury. *Arch. Environ. Occup. Health* 60, 17–23.
- Eklund, S.A., 2010. Trends in dental treatment, 1992 to 2007. *J. Am. Dent. Assoc.* 141, 391–399.
- Engström, K.S., Stromberg, U., Lundh, T., Johannsson, I., Vessby, B., Hallmans, G., Skerfving, S., Broberg, K., 2008. Genetic variation in glutathione-related genes and body burden of methylmercury. *Environ. Health Perspect.* 116, 734–739.
- Gundacker, C., Komarnicki, G., Jagiello, P., Gencikova, A., Dahmen, N., Wittmann, K.J., Gencik, M., 2007. Glutathione s-transferase polymorphism, metallothioneins expression, and mercury levels among students in Austria. *Sci. Total. Environ.* 385, 37–47.
- Gundacker, C., Wittmann, K.J., Kukuckova, M., Komarnicki, G., Hikkel, I., Gencik, M., 2009. Genetic background of lead and mercury metabolism in a group of medical students in Austria. *Environ. Res.* 109, 786–796.
- Hursh, J.B., Clarkson, T.W., Cherian, M.G., Vostal, J.J., Vander Mallie, R., 1976. Clearance of mercury (Hg-197, Hg-203) vapor inhaled by human subjects. *Arch. Environ. Health* 31, 302–309.
- Khan, M.A., Wang, F., 2009. Mercury-selenium compounds and their toxicological significance: toward a molecular understanding of the mercury-selenium antagonism. *Environ. Toxicol. Chem.* 28, 1567–1577.
- Lee, B., Hong, Y., Park, H., Ha, M., Koo, B.S., Chang, N., Roh, Y., Kim, B., Kim, Y., Kim, B., Jo, S., Ha, E., 2009. Interaction between *GSTM1/GSTT1* polymorphism and blood mercury on birth weight. *Environ. Health Perspect.* doi:10.1289/ehp.0900731.
- Levine, L., Fahy, J.P., 1945. Evaluation of lead determinations I: the significance of specific gravity. *J. Ind. Hyg. Toxicol.* 27, 217.
- Livak, K.J., Flood, S.J., Marmaro, J., Giusti, W., Deetz, K., 1995. Oligonucleotides with fluorescent dyes at opposite ends provide a quenched probe system useful for detecting PCR product and nucleic acid hybridization. *PCR Methods Appl.* 4, 357–362.

Mason, H.J., Calder, I.M., 1994. The correction of urinary mercury concentrations in untimed random samples. *Occup. Environ. Med.* 51, 287.

McDowell, M.A., Dillon, C.F., Osterloh, J., Bolger, P.M., Pellizzari, E., Fernando, R., et al., 2004. Mercury levels in U.S. children and women of childbearing age: reference range data from NHANES 1999–2000. *Environ. Health Perspect.* 112, 1165–1171.

Méplan, C., Crosley, L.K., Nicol, F., Beckett, G.J., Howie, A.F., Hill, K.E., et al., 2007. Genetic polymorphisms in the human selenoproteins P gene determine the response of selenoproteins markers to selenium supplementation in a gender-specific manner (the SELGEN study). *FASEB* 21, 3063–3074.

Méplan, C., Nicol, F., Burtle, B.T., Crosley, L.K., Arthur, J.R., Mathers, J.C., et al., 2009. Relative abundance of selenoprotein P isoforms in human plasma depends on genotype, Se intake, and cancer status. *Antioxid. Redox. Signal.* 11, 2631–2640.

Mierzykowski, S.E., Carr, K.C., 2001. Total mercury and methylmercury in freshwater mussels *Elliptio complanata* from the Sudbury River watershed, Massachusetts. USFWS. Special Project Report FY98-MEFO-2-EC. Main Field Office, Old Town, ME.

Min, J.L., Lakenberg, N., Bakker-Verweij, M., Suchiman, E., Boomsma, D.I., Slagboom, P.E., Meulenbelt, I., 2006. High microsatellite and SNP genotyping success rates established in a large number of genomic DNA samples extracted from mouth swabs and genotypes. *Twin Res. Hum. Genet.* 9, 501–506.

Morahan, J.M., Yu, B., Trent, R.J., Pamphlett, R., 2007. Genetic susceptibility to environmental toxicants in ALS. *Am. J. Med. Genet. B* 114B, 885–890.

Mordukhovich, I., Wilker, E., Suh, H., Wright, R., Sparrow, D., Vokonas, P.S., Schwartz, J., 2009. Black carbon exposure, oxidative stress genes, and blood pressure in a repeated-measures study. *Environ. Health Perspect.* 117, 1767–1772.

Ohno, T., Sakamoto, M., Kurosawa, T., Dakeishi, M., Iwata, T., Murata, K., 2007. Total mercury levels in hair, toenail, and urine among women free from occupational exposure and their relations to renal tubular function. *Environ. Res.* 103, 191–197.

Paruchuri, Y., Siuniak, A., Johnson, N., Levin, E., Mitchell, K., Goodrich, J.M., et al., 2010. Occupation and environmental mercury exposure among small-scale gold miners in the Talensi-Nabdam District of Ghana's Upper East region. *Sci. Total. Environ.* 408, 6079–6085.

Rooney, J.P.K., 2007. The role of thiols, dithiols, nutritional factors and interacting ligands in the toxicology of mercury. *Toxicology* 234, 145–156.

U.S. Food and Drug Administration (FDA), . Mercury Concentrations in Fish: FDA Monitoring Program (1990–2004). US Department of Health and Human Services <http://www.fda.gov/food/foodsafety/productspecificinformation/seafood/foodbournepathogenscontaminants/methylmercury/ucm115644.htm> (accessed Aug. 3, 2010).

Wranová, K., Čejchanová, M., Spěváčková, V., Korunová, V., Vobecký, M., Spěváček, V., 2008. Mercury and methylmercury in hair of selected groups of Czech population. *Cent. Eur. J. Public Health* 17, 36–40.