

Influenza-associated Hospitalizations by Industry, 2009–10 Influenza Season, United States

Sara E. Luckhaupt, Marie Haring Sweeney, Renee Funk, Geoffrey M. Calvert, Mackenzie Nowell,¹ Tiffany D'Mello, Arthur Reingold, James Meek, Kimberly Yousey-Hindes, Kathryn E. Arnold, Patricia Ryan, Ruth Lynfield, Craig Morin, Joan Baumbach, Shelley Zansky, Nancy M. Bennett, Ann Thomas, William Schaffner, and Timothy Jones

Medscape **ACTIVITY** EDUCATION

Medscape, LLC is pleased to provide online continuing medical education (CME) for this journal article, allowing clinicians the opportunity to earn CME credit.

This activity has been planned and implemented in accordance with the Essential Areas and policies of the Accreditation Council for Continuing Medical Education through the joint sponsorship of Medscape, LLC and Emerging Infectious Diseases. Medscape, LLC is accredited by the ACCME to provide continuing medical education for physicians.

Medscape, LLC designates this Journal-based CME activity for a maximum of 1 *AMA PRA Category 1 Credit(s)*[™]. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

All other clinicians completing this activity will be issued a certificate of participation. To participate in this journal CME activity: (1) review the learning objectives and author disclosures; (2) study the education content; (3) take the post-test with a 70% minimum passing score and complete the evaluation at www.medscape.org/journal/eid; (4) view/print certificate.

Release date: March 15, 2012; Expiration date: March 15, 2013

Learning Objectives

Upon completion of this activity, participants will be able to:

- Distinguish the effect of employment on the risk for hospitalization for influenza
- Assess the risk for hospitalization for influenza across different sectors of industry
- Evaluate the interaction between underlying medical illness and the risk for hospitalization for influenza among workers
- Analyze other factors potentially associated with the risk for hospitalization for influenza among workers.

Editor

Thomas J. Gryczan, MS, Technical Writer/Editor, *Emerging Infectious Diseases*. *Disclosure: Thomas J. Gryczan, MS, has disclosed no relevant financial relationships.*

CME Author

Charles P. Vega, MD, Health Sciences Clinical Professor; Residency Director, Department of Family Medicine, University of California, Irvine. *Disclosure: Charles P. Vega, MD, has disclosed no relevant financial relationships.*

Authors

Disclosures: Sara E. Luckhaupt, MD, MPH; Mackenzie Nowell, MPH; Tiffany D'Mello, MPH; Arthur Reingold, MD; James Meek, MPH; Kimberly Yousey-Hindes; Renee Funk, DVM; Patricia Ryan, MS; Ruth Lynfield, MD; Craig Morin, MPH; Joan Baumbach, MD, MPH; Shelley Zansky, PhD; Nancy M. Bennett, MD, MS; Ann Thomas, MD, MPH; and Timothy Jones, MD, have disclosed no relevant financial relationships. Marie Haring Sweeney, PhD, MPH, has disclosed the following relevant financial relationships: owns stock, stock options, or bonds from P&G, Apple, Lucent, J&J. Geoffrey M. Calvert, MD, MPH, has disclosed the following relevant financial relationships: owns stock, stock options, or bonds from Bristol-Myers Squibb, Abbott, Amgen, Baxter, Gen-Probe, Johnson & Johnson, Pfizer, Gilead, St. Jude Medical, Stryker, Teva, Novartis. Kathryn E. Arnold, MD, has disclosed the following relevant financial relationships: owns stock, stock options, or bonds from Steris Corp, Kimberley Clark Corp. William Schaffner, MD, has disclosed the following relevant financial relationships: served as an advisor or consultant for Merck, sanofi-Pasteur, Pfizer, GSK, Dynavax.

Author affiliations: Centers for Disease Control and Prevention, Cincinnati, Ohio, USA (S.E. Luckhaupt, M. H. Sweeney, G.M. Calvert); Centers for Disease Control and Prevention, Atlanta, Georgia, USA (R. Funk, M. Nowell, T. D'Mello); University of California, Berkeley, California, USA (A. Reingold); Yale University, New Haven, Connecticut, USA (J. Meek, K. Yousey-Hindes); Georgia Division of Public Health, Atlanta (K. E. Arnold); Maryland Department of Health and Mental Hygiene, Baltimore, Maryland, USA (P. Ryan), Minnesota Department of Health, Minneapolis, Minnesota, USA (R. Lynfield, C. Morin); New Mexico Department of Health, Santa Fe, New Mexico, USA (J. Baumbach); New York State Department of Health, Albany, New York, USA (S. Zansky); University of Rochester, Rochester, New York, USA (N. M. Bennett); Oregon Public Health Division, Portland, Oregon, USA (A. Thomas); Vanderbilt University, Nashville, Tennessee, USA (W. Schaffner); and Tennessee Department of Health, Nashville (T. Jones)

DOI: <http://dx.doi.org/10.3201/eid1804.110337>

¹Current affiliation: Florida Department of Health, Orlando, Florida, USA.

In response to pandemic (H1N1) 2009, data were collected on work status and industry of employment of 3,365 adults hospitalized with laboratory-confirmed influenza during the 2009–10 influenza season in the United States. The proportion of workers hospitalized for influenza was lower than their proportion in the general population, reflecting underlying protective characteristics of workers compared with nonworkers. The most commonly represented sectors were transportation and warehousing; administrative and support and waste management and remediation services; health care; and accommodation and food service.

Although the Occupational Safety and Health Administration, Department of Labor, and the National Institute for Occupational Safety and Health (NIOSH), Centers for Disease Control and Prevention (CDC), have issued guidance to assist workplaces in responding to influenza pandemics (1,2), more information is needed about which specific groups of workers are at highest risk for acquiring or having complications from pandemic and seasonal influenza. Specifically, this information is needed for recognizing and responding to increased risks for infection among key occupational groups (e.g., health care workers, school teachers, retail and food service workers, and others with substantial exposure to the general public); informing persons who develop guidance for key policy questions, including the prioritization of groups to receive vaccine, school closing policies, and appropriate personal protective equipment use; and providing data that might trigger more in-depth case studies of clusters of disease occurring among specific workers.

During the influenza (H1N1) 2009 pandemic, NIOSH explored multiple sources of data on the occupations of affected persons. The occupational distribution of all confirmed (H1N1) 2009 pandemic influenza case-patients from 4 states during the early phase of the pandemic (April–July 2009) has been reported by Suarathana et al. (3) but as the pandemic progressed and case counts rapidly increased, it became impossible to collect occupational information on all case-patients.

Thus, during the fall wave of the pandemic, NIOSH worked with the CDC Emerging Infections Program (EIP) to collect data on the industry and occupation of the subset of adults hospitalized with laboratory-confirmed influenza. Hospitalized case-patients, many of whom have underlying medical risk factors for severe disease, are not representative of all persons who acquire influenza. Thus, studying them provides little insight into the risk of acquiring influenza. However, examining the distribution of industry of employment of these persons provides some clues about specific groups of workers that might be most commonly affected by severe influenza.

Methods

EIP Data for Hospitalized Influenza Case-Patients

The EIP is coordinated and funded by CDC. It consists of a network of 10 state health departments (California, Colorado, Connecticut, Georgia, Maryland, Minnesota, New Mexico, New York, Oregon, and Tennessee) and their collaborators in local health departments, academic institutions, other federal agencies, and public health and clinical laboratories. The network comprises a catchment area of ≈ 16.8 million persons ≥ 18 years of age. The population studied by the EIP is roughly representative of the US population on the basis of demographic characteristics such as age, sex, race, urban residence, and population density and percentage at or below the poverty level. During the 2009–10 influenza season (September 1, 2009–April 30, 2010), the 10 EIP sites performed active population-based surveillance for laboratory-confirmed influenza-related hospitalizations (see www.cdc.gov/ncpcid/deiss/eip/index.html for information about EIP). This period coincided with circulation and dominance of the pandemic (H1N1) 2009 virus strain. Institutional Review Board approval for EIP adult influenza-associated hospitalization surveillance activities during the 2009–10 influenza season was obtained from all sites, including CDC.

As described elsewhere (4), cases were defined as persons ≥ 18 years of age hospitalized for community-onset, laboratory-positive influenza infection. Case-patients were residents of the defined EIP catchment areas and were admitted to a surveillance-area hospital during the influenza season and within 14 days of receiving a positive influenza test result. Laboratory confirmation of influenza was obtained by virus culture, immunofluorescence antibody staining, reverse transcription PCR, or a commercially available rapid diagnostic test. Written documentation of a positive influenza test result in the medical chart was acceptable as evidence of laboratory confirmation. Persons who had positive results for influenza > 3 days after hospital admission were considered to have nosocomial influenza and were excluded from this study. Staff at each EIP site identified cases by contacting hospital laboratories, medical records departments, and infection control practitioners, and by reviewing databases of state-reportable conditions.

Once a person who met the surveillance case definition was identified, his or her hospital medical chart was abstracted and a standardized data collection instrument was completed. The instrument included a question about employment in selected health care occupations and an open text field to record other occupational information.

NIOSH received a deidentified dataset of 4,511 hospitalized cases of influenza from the 2009–10 influenza season from the CDC Influenza EIP program, as of May

18, 2010. One hundred forty-eight (3.3%) case-patients were excluded from analyses because of incomplete data collection. An additional 11 (0.2%) case-patients were excluded because they were hospitalized outside the EIP catchment area, although they resided in a catchment area. We classified case-patients by work status (worker, nonworker, or unknown). Nonworkers included students, homemakers, retired persons, disabled persons, and nonworkers not elsewhere classified. Nonworkers not elsewhere classified included occupation text entries of none, not employed, unemployed, incarcerated, and homeless. All nursing home residents, regardless of occupational information recorded, were grouped with disabled persons because of their similar high prevalence of underlying medical conditions. If the text field for occupation was blank or indicated that the work status of the case-patient was unknown, we classified the work status as unknown. Trained NIOSH coders used responses to the health care worker question and occupational text entries to assign 2-digit codes for industry sector from the North American Industrial Classification System (5) and 2-digit codes for occupational group from the Standard Occupation Classification system (6) to the workers.

The EIP data collection instrument also included information about several underlying conditions associated with increased risk for influenza (asthma, cystic fibrosis, other chronic lung disease, chronic cardiovascular disease, renal disease, chronic metabolic disease [including diabetes], hemoglobinopathy, neuromuscular disorder, diagnosis of cancer [excluding nonmelanoma skin cancer] in the past 12 months, immunosuppressive condition, seizure disorder, Guillain-Barré syndrome, lymphoma or leukemia, cognitive dysfunction, pregnancy, and obesity). The EIP program does not collect any data on tobacco use, socioeconomic factors (e.g., income), or access to primary care among hospitalized influenza case-patients.

National Health Interview Survey Reference Data for Employed US Population

Reference data for the employed US population was obtained from the 2010 National Health Interview Survey (NHIS) public use dataset (www.cdc.gov/nchs/nhis/quest_data_related_1997_forward.htm). The NHIS is a cross-sectional in-person household survey conducted annually by the National Center for Health Statistics, CDC. Data are collected on the civilian noninstitutionalized population of the United States, and thus exclude persons in long-term care facilities (e.g., nursing homes), correctional facilities, active-duty Armed Forces personnel (although civilian family members are included), and US nationals living in foreign countries. The survey uses a multistage clustered sample design with oversampling of black, Hispanic, and Asian persons and produces nationally representative data

on health insurance coverage, health care access and use, health status, health behavior, and other health-related topics.

Data Analyses

The proportions of adults hospitalized for influenza (EIP cases) by employment status were compared with expected proportions in the US population according to the 2010 NHIS, by age group. Among employed adults hospitalized for influenza, the proportions employed in each industry sector were compared with proportions of the US population employed in each industry sector. Ratios >1.0 indicated overrepresentation of an industry sector in the EIP dataset compared with what would be expected if workers from all industry sectors had the same risk for hospitalization because of influenza, which would lead to equal distributions of industry sectors between the 2 datasets. Confidence intervals were calculated by using the χ^2 statistic to approximate the Poisson distribution.

We also used data from the 2010 NHIS to estimate the proportion of US adults employed in each sector who reported ≥ 1 underlying medical conditions, were current smokers, had relatively low annual earnings ($< \$35,000$), and had a usual place to go for health care. We included health conditions reported in the NHIS that most closely matched the underlying health conditions for which data were collected in the EIP influenza study (i.e., asthma, emphysema, chronic bronchitis, cardiovascular disease [hypertension, coronary heart disease, angina, history of myocardial infarction, other heart condition, and/or history of stroke], renal failure, diabetes, diagnosis of cancer [excluding nonmelanoma skin cancer] in the past 12 months, epilepsy, pregnancy, and obesity [body mass index ≥ 30]). These estimates provide some group-level background information about underlying characteristics of workers employed in each industry sector that might affect their risks of hospitalization because of influenza (4,7–12).

Data analyses were performed by using SAS version 9.2 for Windows (SAS Institute, Cary, NC, USA). To account for the complex sampling design of the NHIS, all analyses of NHIS data were completed by using SAS survey procedures appropriate for complex samples and sampling information included in the public use dataset. Estimates based on NHIS data with relative SE $>30\%$ are not presented because of low reliability/precision.

Results

Data were available for 4,352 case-patients who had laboratory-confirmed influenza and who were hospitalized during September 1, 2009–April 24, 2010. Of these case-patients, 3,365 (77.3%) had adequate information recorded to classify them according to work status: 1,283 workers, 96 students, 86 homemakers, 472 retired persons, 535 nursing

home residents and disabled persons, and 893 nonworkers not elsewhere classified.

Overall, workers represent a much lower proportion of EIP hospitalized influenza case-patients compared with their proportions in the general US population according to NHIS data for every age group (EIP:NHIS ratio range 0.61–0.66) (Table 1). A total of 1,070 (83.4%) current workers were assigned 2-digit North American Industrial Classification System industry codes. Industry sectors with overrepresentation among hospitalized influenza case-patients (EIP data) compared with the 2010 NHIS reference population data were transportation and warehousing (ratio 1.53, 95% CI 1.18–1.94), administrative and support and waste management and remediation services (ratio 1.51, 95% CI 1.18–1.91), health care (ratio 1.47, 95% CI 1.26–1.70), and accommodation and food services (ratio 1.35, 95% CI 1.10–1.65) (Table 2).

In general, industry sectors with the highest prevalence of underlying chronic medical conditions (per NHIS, e.g., social assistance, public administration) were not overrepresented in the EIP database (Table 2). For every industry sector, the proportion of workers with underlying medical conditions among hospitalized influenza (EIP) case-patients was higher than the proportion of workers in the general population (NHIS) with underlying medical conditions (Table 2). These ratios varied by industry sector, but neither the industry sector with the highest proportion of workers with underlying medical conditions among hospitalized influenza case-patients (education services) nor the industry sector with the highest ratio of case-patients with underlying conditions compared with workers in the general population with underlying conditions (arts, entertainment, and recreation) were overrepresented in the EIP database (Table 2).

According to the 2010 NHIS reference population data, some industry sectors overrepresented in the EIP database had a higher prevalence of demographic characteristics that might place them at increased risk

for influenza-associated hospitalization. For example, the accommodation and food services industry sector had the highest prevalence of current smokers (30.9%) and workers with relatively low earnings (<\$35,000 per year) (92.3%) (Table 2). With regards to health care access, the industry sectors with the lowest proportion of workers who report having a usual place to go for health care (other than an emergency department) include accommodation and food services (63.9%) and administrative and support and waste management and remediation services (67.7%) (Table 2).

Discussion

Pandemic (H1N1) 2009 virus, which emerged in April 2009, resulted in substantial illness and deaths among working-aged adults. As reported, cumulative rates of laboratory-confirmed, influenza-associated hospitalizations in EIP sites were 2.4/10,000 population for all persons 18–49 years of age during August 30, 2009–March 27, 2010, which was $\approx 6\times$ higher than the influenza-associated hospitalization rate for this age group in 2008–09, when seasonal influenza (H1N1) virus was the predominant strain (13).

We found that workers made up a much lower proportion of EIP hospitalized influenza case-patients than the general US population. This finding was not unexpected because of favorable underlying characteristics of workers compared with nonworkers, such as younger age and lower prevalence of diagnosed underlying medical conditions, which make workers less likely to be hospitalized for influenza. Furthermore, among workers, certain industry sectors are overrepresented in the EIP dataset than what would be expected if workers from all industry sectors had the same risk for hospitalization for influenza, suggesting that the risk for severe influenza varied among different groups of workers during the 2009–10 influenza season.

To date, most evaluations of work-related risk for influenza have focused on health care workers (14–17).

Table 1. Employment status among US adults hospitalized with influenza September 1, 2009–April 24, 2010 by age group, compared with data from the 2010 NHIS*

Status	No. (%) cases†	Estimated US populations in thousands (weighted %) from NHIS†	Ratio (95% CI)
Age 18–49 y			
Employed	825 (45.2)	94,862 (71.3)	0.63 (0.59–0.68)
Not employed	999 (54.8)	38,170 (28.7)	1.91 (1.79–2.03)
Employment status unknown	586 (24.3)	36 (0)	NA
Age 50–64 y			
Employed	409 (41.2)	36,045 (62.4)	0.66 (0.60–0.73)
Not employed	583 (58.8)	21,675 (37.6)	1.56 (1.44–1.70)
Employment status unknown	300 (23.2)	24 (0)	NA
Age ≥ 65 y			
Employed	49 (8.9)	5,697 (14.7)	0.61 (0.45–0.80)
Not employed	500 (91.1)	32,980 (85.3)	1.07 (0.98–1.17)
Employment status unknown	101 (15.5)	15 (0)	NA

*NHIS, National Health Interview Survey; NA, not applicable.

†Percentage among case-patients/respondents with known work status and among all case-patients/respondents with unknown work status.

RESEARCH

There is some evidence that household exposures are more predictive of influenza infection among these workers than occupational exposures (16), but there is also evidence that occupational acquisition occurs (14,17). We found that persons working in the health care industry were overrepresented among hospitalized persons with influenza compared with what would be expected if workers from all industry sectors had the same risk for hospitalization

because of influenza. However, health care workers were not the only worker group overrepresented.

Although the ratios were only modestly increased, these results suggest that groups of workers other than those employed in health care may also be at increased risk for influenza severe enough to result in hospitalization. Overrepresentation of an industry sector in the EIP dataset may be related to demographic and underlying health

Table 2. Hospitalized influenza case-patients by industry sector per EIP (September 1, 2009–April 24, 2010), ratios compared with distribution of employed US adults per NHIS, and characteristics of employed US adults by industry sector per NHIS*

Industry sector of employment (NAICS code)	% Employed hospitalized influenza case-patients, EIP	Weighted % employed adults, NHIS†	Ratio (95% CI)‡	% Case-patients with underlying condition, EIP§	% Adults, NHIS†			
					With underlying condition¶	Who smoke	Annual earnings <\$35,00	With usual place for health care
Transportation and warehousing (48, 49)	6.26	4.10	1.53 (1.18–1.94)	76.12	54.07	20.79	48.23	78.36
Administrative and support and waste management and remediation (56)	6.64	4.38	1.51 (1.18–1.91)	73.24	49.82	25.34	80.74	67.69
Health care (62, except for 624)	16.17	11.01	1.47 (1.26–1.70)	81.40	52.46	16.24	59.12	87.72
Accommodation and food (72)	9.07	6.70	1.35 (1.10–1.65)	73.20	39.32	30.88	92.27	63.92
Other (81)	6.26	5.15	1.22 (0.94–1.54)	75.00	44.88	17.37	78.74	77.23
Social assistance (624)	3.18	2.70	1.18 (0.82–1.64)	82.35	60.66	14.32	81.83	89.66
Information (51)	2.90	2.54	1.14 (0.78–1.62)	87.10	51.19	15.54	44.20	84.80
Retail trade (44, 45)	12.43	11.03	1.13 (0.94–1.34)	85.61	46.35	22.64	80.93	75.84
Finance and insurance (52)	4.95	4.41	1.12 (0.84–1.47)	69.81	48.59	14.60	41.36	87.91
Education (61)	9.63	10.25	0.94 (0.77–1.14)	89.32	50.00	8.40	53.79	91.72
Professional, scientific, and technical (54)	6.26	6.84	0.92 (0.71–1.16)	71.64	41.06	13.35	28.65	85.08
Arts, entertainment, and recreation (71)	1.78	2.04	0.87 (0.52–1.36)	78.95	39.32	21.40	75.67	80.40
Construction (23)	5.05	6.61	0.76 (0.57–1.00)	57.41	46.50	29.09	63.51	70.01
Real estate and rental and leasing (53)	1.21	1.92	0.63 (0.34–1.08)	76.92	45.90	23.33	63.24	78.29
Public administration (92)	3.18	5.44	0.58 (0.40–0.82)	82.35	57.93	14.64	34.99	92.16
Manufacturing (31–33)	4.30	9.56	0.45 (0.33–0.60)	71.74	52.02	21.75	49.79	81.93
Utilities (22)	–	1.00	–	–	53.67	18.20	24.09	88.59
Agriculture, forestry, fishing, and hunting (11)	–	1.18	–	–	44.75	19.36	82.97	71.40
Wholesale trade (42)	–	2.58	–	–	49.76	23.71	48.68	82.70
Mining (21)	–	0.50	–	–	44.93	27.03	35.78	78.99
All employed	38.10	59.50	0.64 (0.59–0.70)	77.05	48.53	19.17	60.39	81.09
All nonemployed	61.90	40.50	1.53 (1.42–1.65)	87.56	65.20	19.62	96.77	84.52

*EIP, Emerging Infections Program; NHIS, National Health Interview Survey; NAICS, North American Industry Classification System; –, <5 cases.

†Weighted estimates based on 2010 NHIS public use dataset (www.cdc.gov/nchs/nhis/quest_data_related_1997_forward.htm).

‡Proportion among EIP cases:proportion of employed adults, per NHIS.

§Include asthma, cystic fibrosis, other chronic lung disease, chronic cardiovascular disease, renal disease, chronic metabolic disease (including diabetes), hemoglobinopathy, neuromuscular disorder, cancer (excluding nonmelanoma skin cancer) in past 12 mo, immunosuppressive condition, seizure disorder, Guillain-Barré syndrome, lymphoma or leukemia, cognitive dysfunction, pregnancy, and obesity.

¶Include asthma, emphysema, chronic bronchitis, cardiovascular disease (hypertension, coronary heart disease, angina, myocardial infarction, other heart condition, or stroke), renal failure, diabetes, cancer (excluding nonmelanoma skin cancer) in past 12 mo, epilepsy, pregnancy, and obesity (body mass index ≥30).

characteristics of the sector's work force that put them at increased risk for acquiring influenza and for being hospitalized with influenza, but it may also partially reflect occupational risk factors for influenza (e.g., exposure to ill members of the public).

Because the EIP data only include cases, it is difficult to assess the potential reasons for overrepresentation of certain industry sectors, but general population estimates of potential contributing factors based on 2010 NHIS data provide some clues. We examined 4 factors for which data or expert consensus suggests associations with hospitalization (or severe outcomes in general) caused by influenza: underlying medical conditions (4,7,8), current smoking behavior (8,9), low income (10–12), and timely access to primary care (8,10), as measured by reporting a usual place to go for health care.

For example, the most highly overrepresented groups of workers among EIP cases, transportation and warehousing and administrative and support and waste management and remediation services, also had the highest prevalence of some unfavorable demographic characteristics, which might place them at increased risk for influenza-associated hospitalization. Workers in the accommodation and food services sector were also overrepresented among EIP cases. It seems logical that these workers may be at increased occupational risk for acquiring influenza because of their high level of interaction with the general public, but NHIS data also suggest some demographic factors that might increase their risk for hospitalization because of severe influenza (e.g., low earnings, smoking, lack of access to medical care).

On the other hand, variation in underlying health status, socioeconomic status, and access to health care by industry group does not appear to explain all of our findings regarding overrepresentation of groups among hospitalized influenza case-patients compared with the general working population. Health care workers are overrepresented among EIP cases despite being relatively healthy and having relatively high earnings and access to health care. Construction workers are underrepresented among EIP cases despite having relatively low earnings and access to health care. These workers in construction might have a relatively low risk for acquiring influenza because of low interaction with the public.

We also found some industry sectors in which we would expect a relatively high level of interaction with the public (e.g., public administration, education) that were not overrepresented among EIP cases. Even if these workers have an increased risk for acquiring influenza from the public, they might have a low risk for progressing to severe influenza requiring hospitalization because of their relatively high earnings and access to health care.

This study has several limitations in addition to the major limitation of relying on a secondary data source (NHIS) for information on the characteristics of workers by industry sector. No useful information on work status was available for 22.7% of EIP hospitalized influenza case-patients. This fact likely reflects the inconsistency of occupational data available in typical hospital records. There is the potential for misclassification of work status and, among workers, misclassification of industry sector because of inconsistency in narrative data recorded for occupation. Although we called the variable we collected occupation, there were more entries that reflected codable industry sectors than reflected codable occupational groups. Thus, we only reported results by industry. Furthermore, in most cases, the available information only enabled industry to be coded at a broad, nonspecific level. For example, it was impossible to distinguish whether many of the health care workers worked in inpatient or outpatient settings.

Our study examined systematically collected influenza surveillance data according to occupational variables. Benefits of using data from the EIP program include laboratory confirmation of influenza and representation of a large population from geographically diverse areas in the United States. Although we were able to identify specific groups of workers that were most heavily affected by severe influenza during the 2009–10 influenza season, more research is needed to understand the reasons for the increased incidence of severe influenza among specific groups of workers. Concurrently, any interventions that focus on these groups of workers should be evaluated for effectiveness and efficiency.

Acknowledgments

We thank state health departments in California, Colorado, Connecticut, Georgia, Maryland, Minnesota, New Mexico, New York, Oregon, and Tennessee; collaborators in local health departments; academic institutions; other federal agencies; public health and clinical laboratories; infection preventionists; and healthcare providers for contributing data to the EIP network; and Laurie Kamimoto, Liping Pan, Pam Schumacher, Matthew Hirst, Elizabeth Smith, Jia Li, SangWoo Tak, and David Weissman for contributing to the study and the manuscript.

Dr Luckhaupt is a medical officer at CDC in Cincinnati, Ohio. Her research interests include surveillance and epidemiology of various chronic and acute health outcomes among various groups of workers.

References

1. Occupational Safety and Health Administration, US Department of Labor. Guidance on preparing workplaces for an influenza pandemic. OSHA 3327–05R, 2009 [cited 2010 Jul 8]. <http://www.osha.gov/Publications/OSHA3327pandemic.pdf>

2. Centers for Disease Control and Prevention. Guidance for businesses and employers to plan and respond to the 2009–2010 influenza season. Updated February 22, 2010 [cited 2010 Jul 8]. <http://www.cdc.gov/h1n1flu/business/guidance/#top>
3. Suarathana E, McFadden JD, Laney AS, Kreiss K, Anderson HA, Hunt DC, et al. Occupational distribution of persons with confirmed 2009 H1N1 influenza. *J Occup Environ Med*. 2010;52:1212–6. <http://dx.doi.org/10.1097/JOM.0b013e3181fd32e4>
4. Dao CN, Kamimoto L, Nowell M, Reingold A, Gershman K, Meek J, et al. Adult hospitalizations for laboratory-positive influenza during the 2005–2006 through 2007–2008 seasons in the United States. *J Infect Dis*. 2010;202:881–8. <http://dx.doi.org/10.1086/655904>
5. Office of Management and Budget. North American industry classification system (NAICS). United States 2007 [cited 2012 Jan 6]. <http://www.census.gov/naics>
6. Office of Management and Budget. Standard occupational classification manual (SOC). United States 2000 [cited 2012 Jan 6]. http://stats.bls.gov/soc/soc_home.htm
7. Jain S, Kamimoto L, Bramley AM, Schmitz AM, Benoit SR, Louie J, et al. Hospitalized patients with 2009 H1N1 influenza in the United States, April–June 2009. *N Engl J Med*. 2009;361:1935–44. <http://dx.doi.org/10.1056/NEJMoa0906695>
8. Gilca R, De Serres G, Boulianne N, Ouhoumane N, Papenburg J, Douville-Fradet M, et al. Risk factors for hospitalization and severe outcomes of 2009 pandemic H1N1 influenza in Quebec, Canada. *Influenza Other Respi Viruses*. 2011;5:247–55. <http://dx.doi.org/10.1111/j.1750-2659.2011.00204.x>
9. Huttunen R, Heikkinen T, Syrjänen J. Smoking and the outcome of infection. *J Intern Med*. 2011;269:258–69. <http://dx.doi.org/10.1111/j.1365-2796.2010.02332.x>
10. Bouye K, Truman BI, Hutchins S, Richard R, Brown C, Guillory JA, et al. Pandemic influenza preparedness and response among public-housing residents, single-parent families, and low-income populations. *Am J Public Health*. 2009;99:S287–93. <http://dx.doi.org/10.2105/AJPH.2009.165134>
11. Thompson DL, Jungk J, Hancock E, Smelser C, Landen M, Nichols M, et al. Risk factors for 2009 pandemic influenza A (H1N1)-related hospitalization and death among racial/ethnic groups in New Mexico. *Am J Public Health*. 2011;101:1776–84.
12. Yousey-Hindes KM, Hadler JL. Neighborhood socioeconomic status and influenza hospitalizations among children: New Haven County, Connecticut, 2003–2010. *Am J Public Health*. 2011;101:1785–9.
13. Centers for Disease Control and Prevention. Update: influenza—United States, August 30, 2009–March 27, 2010, and composition of the 2010–11 influenza vaccine. *MMWR Morb Mortal Wkly Rep*. 2010;59:423–30.
14. Centers for Disease Control and Prevention. Novel influenza A (H1N1) virus infections among healthcare personnel—United States, April–May 2009. *MMWR Morb Mortal Wkly Rep*. 2009;58:641–5.
15. Santos CD, Bristow RB, Vorenkamp JV. Which healthcare workers were most affected during the spring 2009 H1N1 pandemic? *Disaster Med Public Health Prep*. 2010;4:47–54.
16. Williams CJ, Schweiger B, Diner G, Haaman F, Krause G, Nienhaus A, et al. Seasonal influenza risk in hospital healthcare workers is more strongly associated with household than occupational exposures: results from a prospective cohort study in Berlin, Germany, 2006/07. *BMC Infect Dis*. 2010;10:8. <http://dx.doi.org/10.1186/1471-2334-10-8>
17. Wise ME, De Perio M, Halpin J, Jhung M, Magill S, Black SR, et al. Transmission of pandemic (H1N1) 2009 influenza to healthcare personnel in the United States. *Clin Infect Dis*. 2011;52:S198–204. <http://dx.doi.org/10.1093/cid/ciq038>

Address for correspondence: Sara E. Luckhaupt, Centers for Disease Control and Prevention, 4676 Columbia Pkwy, Mailstop R17, Cincinnati, OH 45226-1998, USA; email: pks8@cdc.gov

All material published in *Emerging Infectious Diseases* is in the public domain and may be used and reprinted without special permission; proper citation, however, is required.

Get the content you want delivered to your inbox.



Table of Contents
Podcasts
Ahead of Print Articles
Medscape CME™
Specialized Content

Online subscription: www.cdc.gov/ncidod/eid/subscrib.htm

EMERGING INFECTIOUS DISEASES®



April 2012

Vector-borne Infections



The Charles F. Buxton Foundation. Archives of the Buxton-Perry Air Center at Buffalo State College, Buffalo, New York
Gift of the Charles F. Buxton Foundation, 2006

EMERGING INFECTIOUS DISEASES®

EDITOR-IN-CHIEF

D. Peter Drotman

Managing Senior Editor

Polyxeni Potter, Atlanta, Georgia, USA

Senior Associate Editor

Brian W.J. Mahy, Bury St. Edmunds, Suffolk, UK

Associate Editors

Paul Arguin, Atlanta, Georgia, USA
 Charles Ben Beard, Ft. Collins, Colorado, USA
 Ermias Belay, Atlanta, Georgia, USA
 David Bell, Atlanta, Georgia, USA
 Corrie Brown, Athens, Georgia, USA
 Charles H. Calisher, Ft. Collins, Colorado, USA
 Michel Drancourt, Marseille, France
 Paul V. Effler, Perth, Australia
 David Freedman, Birmingham, Alabama, USA
 Peter Gerner-Smidt, Atlanta, Georgia, USA
 Stephen Hadler, Atlanta, Georgia, USA
 Nina Marano, Atlanta, Georgia, USA
 Martin I. Meltzer, Atlanta, Georgia, USA
 David Morens, Bethesda, Maryland, USA
 J. Glenn Morris, Gainesville, Florida, USA
 Patrice Nordmann, Paris, France
 Tanja Popovic, Atlanta, Georgia, USA
 Didier Raoult, Marseille, France
 Pierre Rollin, Atlanta, Georgia, USA
 Ronald M. Rosenberg, Fort Collins, Colorado, USA
 Dixie E. Snider, Atlanta, Georgia, USA
 Frank Sorvillo, Los Angeles, California, USA
 David Walker, Galveston, Texas, USA
 J. Todd Weber, Atlanta, Georgia, USA
 Henrik C. Wegener, Copenhagen, Denmark

Founding Editor

Joseph E. McDade, Rome, Georgia, USA

Copy Editors Claudia Chesley, Karen Foster, Thomas Gryczan, Carol Snarey, P. Lynne Stockton

Production Carrie Huntington, Ann Jordan, Shannon O'Connor, Reginald Tucker

Editorial Assistant Christina Dzikowski

Social Media/Communications Sarah Logan Gregory

Emerging Infectious Diseases is published monthly by the Centers for Disease Control and Prevention, 1600 Clifton Road, Mailstop D61, Atlanta, GA 30333, USA. Telephone 404-639-1960, fax 404-639-1954, email eideditor@cdc.gov.

The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the Centers for Disease Control and Prevention or the institutions with which the authors are affiliated.

All material published in Emerging Infectious Diseases is in the public domain and may be used and reprinted without special permission; proper citation, however, is required.

Use of trade names is for identification only and does not imply endorsement by the Public Health Service or by the U.S. Department of Health and Human Services.

EDITORIAL BOARD

Dennis Alexander, Addlestone Surrey, United Kingdom
 Timothy Barrett, Atlanta, Georgia, USA
 Barry J. Beaty, Ft. Collins, Colorado, USA
 Martin J. Blaser, New York, New York, USA
 Sharon Bloom, Atlanta, GA, USA
 Christopher Braden, Atlanta, Georgia, USA
 Mary Brandt, Atlanta, Georgia, USA
 Arturo Casadevall, New York, New York, USA
 Kenneth C. Castro, Atlanta, Georgia, USA
 Louisa Chapman, Atlanta, Georgia, USA
 Thomas Cleary, Houston, Texas, USA
 Vincent Deubel, Shanghai, China
 Ed Eitzen, Washington, DC, USA
 Daniel Feikin, Baltimore, Maryland, USA
 Anthony Fiore, Atlanta, Georgia, USA
 Kathleen Gensheimer, Cambridge, Massachusetts, USA
 Duane J. Gubler, Singapore
 Richard L. Guerrant, Charlottesville, Virginia, USA
 Scott Halstead, Arlington, Virginia, USA
 David L. Heymann, London, UK
 Charles King, Cleveland, Ohio, USA
 Keith Klugman, Atlanta, Georgia, USA
 Takeshi Kurata, Tokyo, Japan
 S.K. Lam, Kuala Lumpur, Malaysia
 Stuart Levy, Boston, Massachusetts, USA
 John S. MacKenzie, Perth, Australia
 Marian McDonald, Atlanta, Georgia, USA
 John E. McGowan, Jr., Atlanta, Georgia, USA
 Tom Marrie, Halifax, Nova Scotia, Canada
 Philip P. Mortimer, London, United Kingdom
 Fred A. Murphy, Galveston, Texas, USA
 Barbara E. Murray, Houston, Texas, USA
 P. Keith Murray, Geelong, Australia
 Stephen M. Ostroff, Harrisburg, Pennsylvania, USA
 David H. Persing, Seattle, Washington, USA
 Richard Platt, Boston, Massachusetts, USA
 Gabriel Rabinovich, Buenos Aires, Argentina
 Mario Raviglione, Geneva, Switzerland
 David Relman, Palo Alto, California, USA
 Connie Schmaljohn, Frederick, Maryland, USA
 Tom Schwan, Hamilton, Montana, USA
 Ira Schwartz, Valhalla, New York, USA
 Tom Shinnick, Atlanta, Georgia, USA
 Bonnie Smoak, Bethesda, Maryland, USA
 Rosemary Soave, New York, New York, USA
 P. Frederick Sparling, Chapel Hill, North Carolina, USA
 Robert Swanepoel, Pretoria, South Africa
 Phillip Tarr, St. Louis, Missouri, USA
 Timothy Tucker, Cape Town, South Africa
 Elaine Tuomanen, Memphis, Tennessee, USA
 John Ward, Atlanta, Georgia, USA
 Mary E. Wilson, Cambridge, Massachusetts, USA

∞ Emerging Infectious Diseases is printed on acid-free paper that meets the requirements of ANSI/NISO 239.48-1992 (Permanence of Paper)

EMERGING INFECTIOUS DISEASES

April 2012



On the Cover

Charles E. Burchfield
(1893–1967)
Camouflage design (1918)
Watercolor and graphite on
paper mounted on black paper
(27.3 cm x 32.1 cm)

The Charles E. Burchfield
Foundation Archives at the
Burchfield Penney Art Center at
Buffalo State College, Buffalo,
New York
Gift of the Charles E. Burchfield
Foundation, 2006

About the Cover p. 706

Geographic Distribution of Hantaviruses Associated with Neotomine and Sigmodontine Rodents, Mexico 571

M.L. Milazzo et al.
El Moro Canyon virus and Limestone Canyon
virus are widely distributed and may cause
hantavirus pulmonary syndrome.

Shiga Toxin–producing *Escherichia coli* serotype O78:H– in Family, Finland, 2009 577

T. Lienemann et al.
STEC strains carrying *stx_{1c}* and *hlyA* genes can
invade the human bloodstream.

Identification of Intermediate in Evolutionary Model of Enterohemorrhagic *Escherichia coli* O157 582

C. Jenke et al.
Single-nucleotide polymorphism typing found
missing link between human strains in strain
from deer.

Emergence of Unusual G6P[6] Rotaviruses in Children, Burkina Faso, 2009–2010..... 589

J. Nordgren et al.
High incidence highlights the need for long-term
surveillance of rotavirus strains.

Comparison of *Escherichia coli* ST131 Pulsotypes by Epidemiologic Traits, 1967–2009..... 598

J.R. Johnson et al.
Certain high-prevalence pulsed-field gel
electrophoresis types exhibited distinctive
temporal patterns and epidemiologic
associations.

Research

Medscape EDUCATION ACTIVITY



Determinants for Autopsy after Unexplained Deaths Possibly Resulting from Infectious Causes, United States 549

L. Liu et al.

Autopsy findings, clinical history, and diagnostic
tools can aid surveillance and investigation of
infectious diseases.



p. 612

Medscape EDUCATION ACTIVITY



Influenza-associated Hospitalizations by Industry, 2009–10 Influenza Season, United States 556

S.E. Luckhaupt et al.

Certain industries were overrepresented among
employed adults with this disease.

p. 617

Risk Factors for Chronic Q Fever, the Netherlands 563

L.M. Kampschreur et al.

Previous cardiac valvular surgery, vascular
prosthesis, aortic aneurysm, renal insufficiency,
and older age increased risk.



EMERGING INFECTIOUS DISEASES

April 2012

Policy Review

Lessons Learned during Dengue Outbreaks in the United States, 2001–2011..... 608

A.A. Adalja et al.

Public health authorities should involve the clinical, laboratory community, and local communities in vector control and case reporting.

Historical Reviews

Malaria in Highlands of Ecuador since 1900..... 615

L.L. Pinault and F. F. Hunter

Eliminated after 1950, malaria may be reemerging in a new region.

Dengue and US Military Operations from Spanish–American War through Today..... 623

R.V. Gibbons et al.

Dengue may remain problematic for military personnel until an effective vaccine is licensed.

Dispatches

631 *Bartonella* spp. in Rats and Zoonoses, Los Angeles, California, USA

V.A.K.B. Gundi et al.

634 Detection of *Plasmodium* spp. in Human Feces

M. Jirků et al.

637 Extraintestinal Infections Caused by *Salmonella enterica* Subspecies II–IV

S.L. Abbott et al.

640 Subclinical Infections with Crimean-Congo Hemorrhagic Fever Virus, Turkey

H. Bodur et al.

643 Crimean-Congo Hemorrhagic Fever, Kazakhstan, 2009–2010

B. Knust et al.



p. 626

p. 664



646 Vector Blood Meals and Chagas Disease Transmission Potential, United States

L. Stevens et al.

650 *emm59* Group A *Streptococcus* Strains, United States

N. Fittipaldi et al.

653 Characterization of *Mycobacterium orygis* as *M. tuberculosis* Complex Subspecies

J. van Ingen et al.

656 Cosavirus Infection in Persons with and without Gastroenteritis, Brazil

A. Stöcker et al.

660 Drug Susceptibility of *Mycobacterium tuberculosis* Beijing Genotype and Association with MDR TB

J.E.M. de Steenwinkel et al.

664 Highly Divergent Novel Lyssavirus in an African Civet

D.A. Marston et al.

668 *Coccidioides posadasii* Infection in Bats, Brazil

R. de Aguiar Cordeiro et al.

671 Surveillance for West Nile Virus, Dengue, and Chikungunya Infections, Italy, 2010

F. Gobbi et al.

674 De Novo Daptomycin-Nonsusceptible Enterococcal Infections

T. Kelesidis et al.

677 Dengue in Patients with Central Nervous System Manifestations, Brazil

F. Araújo et al.

EMERGING INFECTIOUS DISEASES

April 2012

- 680 Human Parvovirus 4 Infection, Cameroon

M. Lavoie et al.


- 684 Neuroinvasive Disease and West Nile Virus Infection, North Dakota, USA, 1999–2008

P.J. Carson et al.

Another Dimension

- 687 Leaving the Hospital

A. Silver



p. 669

Letters

- 688 West Nile Virus Lineage 2 from Blood Donor, Greece

- 689 Tuberculosis Screening before Anti-Hepatitis C Virus Therapy in Prisons


- 691 Deficient Reporting in Avian Influenza Surveillance, Mali

- 693 Myxozoan Parasite in Brain of Critically Endangered Frog

- 695 Mayaro Virus Infection in Swiss Traveler Returning from Amazon Basin, Northern Peru

- 697 Meningoencephalitis Complicating Relapsing Fever in Traveler Returning from Senegal

p. 694



- 698 Orthopoxvirus Infection in Buffaloes, Brazil

- 700 MSSA ST398, New York and New Jersey, USA

- 702 *Rickettsia monacensis* and Mediterranean Spotted Fever-like Illness, Italy

- 704 *Leishmania* Resistance to Miltefosine Associated with Genetic Marker

- 706 Prolonged KI Polyomavirus Infection in Immunodeficient Child

- 708 African Swine Fever Virus Caucasus Isolate in European Wild Boars of All Ages

About the Cover

- 709 Military Magic or Nature's Fool

Online Report

Multidisciplinary and Evidence-based Method for Prioritizing Diseases of Food-producing Animals and Zoonoses

http://wwwnc.cdc.gov/eid/article/18/4/11-1151_article.htm

The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the Centers for Disease Control and Prevention or the institutions with which the authors are affiliated.



The EID journal is now using **Quick Response codes** (QR codes) to make its content more interactive, engaging, and accessible. QR codes can be interpreted by numerous smartphone apps. Free downloads of readers are available online. These codes will allow readers to access CME, articles, podcasts, and other relevant content, easily from smartphones.