



Journal of Toxicology and Environmental Health, Part B: Critical Reviews

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uteb20>

Respiratory Health Effects of Large Animal Farming Environments

Sara May^a, Debra J. Romberger^{a b} & Jill A. Poole^a

^a Pulmonary, Critical Care, Sleep & Allergy Division, Department of Internal Medicine, University of Nebraska Medical Center, Omaha, Nebraska, USA

^b Veterans Administration Nebraska Western Iowa Healthcare System, Omaha, Nebraska, USA
Version of record first published: 30 Nov 2012.

To cite this article: Sara May, Debra J. Romberger & Jill A. Poole (2012): Respiratory Health Effects of Large Animal Farming Environments, Journal of Toxicology and Environmental Health, Part B: Critical Reviews, 15:8, 524-541

To link to this article: <http://dx.doi.org/10.1080/10937404.2012.744288>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

RESPIRATORY HEALTH EFFECTS OF LARGE ANIMAL FARMING ENVIRONMENTS

Sara May¹, Debra J. Romberger^{1,2}, Jill A. Poole¹

¹Pulmonary, Critical Care, Sleep & Allergy Division, Department of Internal Medicine, University of Nebraska Medical Center, Omaha, Nebraska, USA

²Veterans Administration Nebraska Western Iowa Healthcare System, Omaha, Nebraska, USA

With increases in large animal-feeding operations to meet consumer demand, adverse upper and lower respiratory health effects in exposed agriculture workers are a concern. The aim of this study was to review large animal confinement feeding operational exposures associated with respiratory disease with a focus on recent advances in the knowledge of causative factors and cellular and immunological mechanisms. A PubMed search was conducted with the keywords airway, farm, swine, dairy, horse, cattle inflammation, organic dust, endotoxin, and peptidoglycan, among items were published between 1980 and now. Articles were selected based on their relevance to environmental exposure and reference to airway diseases. Airway diseases included rhinitis, sinusitis, mucus membrane inflammation syndrome, asthma, chronic bronchitis, chronic obstructive pulmonary disease, hypersensitivity pneumonitis, and organic dust toxic syndrome. There is lower prevalence of immunoglobulin (Ig) E-mediated asthma and atopy in farmers and their children, but organic dust worsens existing asthma. Multiple etiologic factors are linked to disease, including allergens, organic dusts, endotoxins, peptidoglycans, and gases. Large animal confinement feeding operations contain a wide diversity of microbes with increasing focus on gram-positive bacteria and archaeobacteria as opposed to gram-negative bacteria in mediating disease. Toll-like receptors (TLR) and nucleotide oligomerization domain (NOD)-like innate immune pathways respond to these exposures. Finally, a chronic inflammatory adaptation, tolerance-like response in chronically exposed workers occurs. Large animal confinement farming exposures produce a wide spectrum of upper and lower respiratory tract diseases due to the complex diversity of organic dust, particulates, microbial cell wall components, and gases and resultant activation of various innate immune receptor signaling pathways.

Chronic airway diseases that develop from exposure to large animal-feeding operations include a spectrum of upper- and lower-respiratory-tract disorders: rhinitis, mucus membrane inflammation syndrome, sinusitis, asthma, asthma-like syndrome, chronic bronchitis, chronic obstructive pulmonary disease (COPD), hypersensitivity pneumonitis, and organic dust toxic syndrome (ODTS) (Eduard et al., 2004, 2009; Girard and Cormier, 2010; Kogevinas et al., 1999; Monso et al., 2004;

Omland, 2002; Reynolds et al., 1996; Von Essen and Donham, 1999). These diseases commonly occur following exposure to large animal-feeding operation farming environments, particularly swine confinement facilities and commercial cattle feedlots. In North America, changes in farming practices have occurred to meet consumer demand for meat globally. This includes an increase in large animal-feeding operations, which may potentially impact respiratory disease development.

This study was supported by grants from the National Institute of Environmental Health Sciences (R01: ES019325 to JAP) and National Institute of Occupational Safety Health (R01OH008539 to DJR and P01OH010162 to JAP and DJR). The authors thank Lisa Chudomelka for her expertise and help with organizing and editing this article.

Address correspondence to Jill A. Poole, MD, Pulmonary, Critical Care, Sleep & Allergy Division, Department of Internal Medicine, University of Nebraska Medical Center, 985300 The Nebraska Medical Center, Omaha, NE 68198-5300, USA. E-mail: japoole@unmc.edu

The majority of hogs are currently raised in confinement with confinement facilities housing an average of 4,646 hogs in the United States in 2004, which is an elevation from an average farm size of 945 hogs in 1992 (Key and McBride, 2008; <http://webarchives.cdlib.org/sw1vh5dg3r/http://ers.usda.gov/AmberWaves/April08/Features/USHogFarms.htm>). Similarly, the farm size in Canada also increased, and it was reported for the first time in 2010 that the average number of hogs exceeded 1,600 head (Key and McBride, 2008; <http://webarchives.cdlib.org/sw1vh5dg3r/http://ers.usda.gov/AmberWaves/April08/Features/USHogFarms.htm>). Changes in farming strategies have also occurred in Europe. News sources have reported the formation of factory farms in Poland housing approximately 10,000 swine (<http://news.bbc.co.uk/2/hi/science/nature/4035081.stm>). Similarly, almost all slaughtered cattle for beef consumption are raised in commercial feedlots. Forty percent of all cattle bred for beef are placed on feedlots that hold 32,000 or more head of cattle while gaining weight for slaughtering (U.S. Department of Agriculture, 2009, www.ers.usda.gov/briefing/cattle/background.htm).

These changes from small, family-owned farms that are still prevalent in Europe and parts of the United States and Canada, toward large-scale farming practices, has potential impact on respiratory diseases. Kirkhorn and Garry (2000) estimated that 50% of swine farmers/workers are at risk of developing respiratory-tract symptoms. Cough with or without sputum production, chest tightness, and wheeze are symptoms that are more commonly reported by confinement workers than the general population (Von Essen et al., 2010; Zejda et al., 1993). Chronic or repetitive exposure to these organic dust environments is also associated with lung function decline over time (Donham et al., 1995; Reynolds et al., 1996; Von Essen et al., 2010; Zejda et al., 1993). Even though farmers use tobacco significantly less than the general population, this population has a substantial increase in morbidity and mortality from respiratory disease (American Thoracic Society, 1998). An overview of the respiratory disease

consequences associated with farming practices was last comprehensively reviewed by the American Thoracic Society (1998). The aim of this report is to review the current knowledge of respiratory diseases associated with large animal confinement feeding operational exposures with focus on recent advances in the knowledge of causative factors and cellular and immunological mechanisms underlying disease manifestations.

AIRWAY DISEASE MANIFESTATIONS

Upper Airway Respiratory Diseases

Upper respiratory diseases along the spectrum of airway diseases associated with large animal confinement farming include rhinitis, sinusitis, and mucous membrane inflammation syndrome, with allergic and irritant rhinitis symptoms most frequently reported (Kirkhorn and Garry, 2000; Poole et al., 2007; Reynolds et al., 1996; Slager et al., 2010; Von Essen and Donham, 1999; Von Essen and Romberger, 2003). Within swine veterinarians, 60% reported rhinitis symptoms (Poole et al., 2007a), and among farmers in the large U.S. Agricultural Health Study, 67% of farmers reported rhinitis, with 39% having three or more episodes of rhinitis in the past year (Slager et al., 2010). Nasal symptoms including congestion, rhinorrhea, sneezing, and/or pruritus are commonly associated with acute and chronic inflammation of the nasal mucous membrane. Nasal lavage fluid demonstrates increases in levels of neutrophils, interleukin (IL)-8/CXCL8, and IL-6 from farmers following swine barn exposure (Burch et al., 2010; Dosman et al., 2000; Hiel et al., 2009; Palmberg et al., 2002; Von Essen and Romberger, 2003). Importantly, proper use of the N-95 respirator results in significant decreases in the proinflammatory nasal response in farmers; however, N-95 respirators are not routinely worn at the facility (Dosman et al., 2000).

Mucus membrane inflammation syndrome is a syndrome characterized by nasal, eye, and throat complaints found in animal confinement workers and associated with increases in nasal

lavage IL-1 α , IL-1 β , and IL-6 levels (Von Essen and Donham, 1999; Von Essen and Romberger, 2003). It is most commonly reported if workers are exposed to dusts and gases within swine confinement facilities (Kirkhorn and Garry, 2000; Von Essen and Donham, 1999). Nasal symptoms such as congestion, rhinorrhea, and pruritus occur in 50% of swine confinement workers, and 25% have sinusitis (Kirkhorn and Garry, 2000; Von Essen and Romberger, 2003).

Asthma

Asthma, defined as a chronic inflammatory disease of the airways characterized by variable and recurring symptoms, reversible air-flow obstruction, airway hyperresponsiveness, and bronchospasm, is associated with large animal farming environmental exposures. It is now well recognized that children raised on farms have less allergic, IgE-mediated asthma (Braun-Fahrlander, 2003; Eduard et al., 2004; Ege et al., 2011; Ernst and Cormier, 2000; Klintberg et al., 2001; Merchant et al., 2005; Rennie et al., 2012; Riedler et al., 2000). It was postulated that exposure to endotoxin or other bacterial components abundantly present in various farming environments leads to decreased IgE-mediated disease development (Braun-Fahrlander et al., 1999; Braun-Fahrlander, 2003; Chen et al., 2007; Cleave et al., 2010; Ernst and Cormier, 2000; Klintberg et al., 2001; Lawson et al., 2011a; 2011b; Pekkanen et al., 2001; Portengen et al., 2002; Radon et al., 2004; Riedler et al., 2000), which is also consistent with the so-called hygiene hypothesis. Indeed, data from large European childhood studies including Prevention of Allergy–Risk Factors for Sensitization in Children Related to Farming and Anthroposophic Lifestyle (PARSIFAL) and the multidisciplinary Study to Identify the Genetic and Environmental Causes of Asthma in the European Community (GABRIEL) demonstrated an increase in exposure to microorganisms in farm children with a corresponding inverse relation with the prevalence of asthma and atopy compared to reference groups (Ege et al., 2011). The PARSIFAL study sampled dust from mattresses

of farm children and used single-strand conformation polymorphism (SSCP) to identify bacteria in school children of both rural and suburban Germany. The GABRIEL study was conducted in elementary school children from Austria, southern Germany, and Switzerland and sampled bacteria in dust from bedroom floors using culture techniques. Both studies found a lower prevalence of asthma (PARSIFAL study: odds ratio [OR] 0.49, confidence interval [CI] 0.35–0.69; GABRIEL study: OR 0.76, CI 0.65–0.89) and atopy (PARSIFAL study: OR 0.34, CI 0.18–0.34; GABRIEL study: OR 0.51, CI 0.46–0.57) in farm children, as well as elevated exposure to the amount and diversity of microorganisms in the dust (Ege et al., 2011).

In Austria, Riedler et al. (2000) conducted a cross-sectional survey of allergic diseases and environmental factors of more than 2000 children aged 8–10 yr. Data showed lower prevalence of hay fever (18.8% vs. 10.3%, $p = .0002$), asthma (1.1% vs. 3.9%, $p = .017$), and positive skin prick test (18.8% vs. 32.7%, $p = .001$) in children living on a farm. The risk reduction was only seen when regular exposure to farm animals was evaluated (Riedler et al., 2000). In addition, there was decreased incidence of asthma and atopy in rural Canadian children raised on a farm, especially girls for current wheeze (OR 0.7) and asthma (OR 0.59) (Ernst and Cormier, 2000). However, an increased rate of respiratory symptoms was reported in children who cleaned large animal pens (Farthing et al., 2009).

Despite these observations of lower prevalence of atopy and IgE-mediated asthma in children raised on farms, there are also reports of increased incidence of asthma, particularly non-IgE-mediated asthma, in agriculturally exposed children and adults (Lawson et al., 2011a, 2011b). In a large cohort study in rural Iowa, in the United States, the asthma rate was increased in children of swine farmers with reported prevalence rates of asthma approaching greater than 50% in children living on farms where swine were fed antibiotic-supplemented food (Merchant et al., 2005). However, these farm children displayed less evidence for IgE-mediated disease and atopy. Moreover,

Merchant et al. (2005) reported that two-thirds of the children had symptoms consistent with severe asthma, yet did not have a physician diagnosis of asthma, suggesting that asthma may be underdiagnosed in this population. It was speculated that while endotoxin levels are higher in agricultural environments (Mayeux, 1997) and may be protective of atopy development via modifying the immune response, high endotoxin might also exacerbate already existing asthma (Cleave et al., 2010; Lawson et al., 2011a, 2011b; Reed and Milton, 2001). It was also proposed that children with asthma are more sensitive to the proinflammatory effects of endotoxin marked by nonspecific airway hyperresponsiveness (Lawson et al., 2011a).

The European Community Respiratory Health Survey reported that the highest risk of developing occupational asthma was the occupation of farmer (OR 2.6, CI 1.3 to 5.4), followed by agricultural workers (OR 1.8, CI 1.0 to 3.2) (Eduard et al., 2009; Kogevinas et al., 1999). In adult farmers, asthma symptoms are more likely regarded as work-exacerbated asthma (WEA), as opposed to occupational asthma (OA) (Eduard et al., 2009). Subjects with moderate or severe asthma and/or subjects not receiving optimal treatment of their asthma may develop WEA when exposed to potential irritants such as dusts, fumes, and sprays (Tarlo and Liss, 2003; Tarlo et al., 2009). OA is new-onset asthma induced by an environmental aspect of the workplace and 90% of the time is IgE-mediated to components, which include animal dander, storage mites, and cockroach (Tarlo and Liss, 2003; Tarlo et al., 2009). A small portion of cases is irritant-induced OA, which is asthma induced by high concentrations of inhaled irritants in confinement facilities. Irritant-induced OA needs to be considered if symptoms of asthma occur within 24 h of the exposure. Finally, gender may play a role in OA, because investigators reported four new cases of OA with symptoms of wheezing and cough in only females without previous atopy or asthma only a couple of weeks after starting a full-time position in swine production facilities (Von Essen et al., 2010; Dosman et al., 2004). However, 2 yr later, the same

authors published three more cases of newly diagnosed OA in swine confinement workers. These three cases were all males, but similar to the females, none had known previous atopy or asthma (Dosman et al., 2006).

There are also reports of an asthma-like syndrome that is described by cough, chest tightness, dyspnea, and wheezing. However, unlike asthma, the asthma-like syndrome does not demonstrate signs of airway obstruction on pulmonary function testing (PFT), and lacks signs of persistent airway inflammation and eosinophil influx, yet methacholine challenge reveals airway hyperresponsiveness in symptomatic workers (Cleave et al., 2010; Von Essen et al., 2010). Asthma-like symptoms are most pronounced upon returning to work after a weekend or vacation holiday, and are reported in up to 50% of swine confinement workers and grain elevator operators (Kirkhorn and Garry, 2000; Von Essen and Romberger, 2003). Future studies are warranted to better define and characterize the spectrum of occupational asthma disorders resulting from large animal farming practices.

Chronic Bronchitis and Chronic Obstructive Pulmonary Disease (COPD)

Farmers and agriculture workers are at an increased risk of respiratory morbidity and mortality from chronic bronchitis and COPD (Eduard et al., 2009). Chronic bronchitis, defined by symptoms of a cough productive of sputum for 3 months of a year for at least 2 consecutive years, affects animal production workers. Eduard and colleagues (2009) reviewed multiple studies from European farmers and found that Danish swine farmers had the highest prevalence of chronic bronchitis at 32%, compared to 28% in farmers that had swine and cattle. Farmers who did not raise any livestock displayed the lowest prevalence at 18.6% (Omland, 2002). In addition, livestock farmers and dairy farmers demonstrated significantly increased risk of developing COPD (livestock farmers, OR 1.4, CI: 1.1 to 1.7; dairy farmers, O.R.1.3, CI: 1.0 to 1.7) (Eduard et al., 2009). Moreover, raising more than one type of

livestock enhanced the risk of farmers developing chronic bronchitis and COPD as compared to crop farmers (Eduard et al., 2009). The livestock farmers also showed the lowest forced expiratory volume in 1 s (FEV₁), consistent with the pattern of lung function decline (Eduard et al., 2009).

Organic Dust Toxic Syndrome

Organic dust toxic syndrome (ODTS) is defined as a febrile illness characterized by malaise, myalgias, chest tightness, headache, and nausea following exposure to large amounts of organic dust. This clinical syndrome was first described in the United States approximately 40 years ago in silo unloading (May et al., 1986; Von Essen and Romberger, 2003; Von Essen et al., 1990). Although farmers were more likely to describe cough or chest tightness while handling grain and grain sorghum (Von Essen et al., 1990), ODTS has since been described following exposure to a variety of organic dust environments (Von Essen and Romberger, 2003). Onset of symptoms usually occurs 4–8 h after exposure and may last for several days. Clinical evaluation demonstrates peripheral blood neutrophilia, but otherwise normal chest radiograph imaging, arterial blood gas measurements, and pulmonary function testing (Von Essen and Romberger, 2003). Although this is typically a self-limiting disease, there is an increased risk of developing bronchitis. The reported prevalence of organic dust toxic syndrome may be as high as 34% in swine confinement workers (Donham et al., 1995; Vogelzang, et al., 1999; Von Essen and Romberger, 2003).

Hypersensitivity Pneumonitis

Although hypersensitivity pneumonitis shares similar clinical symptoms with ODTS, specific diagnostic guidelines exist. These guidelines include evidence for interstitial lung disease by clinical history and physical examination, restrictive pattern found on pulmonary function testing, and chest radiographic imaging findings consistent with interstitial lung

disease (Girard and Cormier, 2010; Kirkhorn and Garry, 2000). Hypersensitivity pneumonitis is induced by a specific antigenic exposure that elicits an antibody response. Hypersensitivity pneumonitis is an important respiratory disease in farmers, as the first cases of hypersensitivity pneumonitis were referred to as “farmer’s lung” because the etiologic agent is commonly fungi (and bacteria) associated with moldy grain and hay. Because hypersensitivity pneumonitis is more associated with grain farming and has recently been reviewed in detail by Girard and Cormier (2010), refer to that review article for more details.

In interim summary, large animal confinement farming is associated with upper and lower airway diseases (Table 1). Rhinitis is a common feature reported by exposed workers with evidence of increased inflammatory markers of neutrophils and cytokines/chemokines by nasal lavage fluid analysis. Overall, studies indicate a decrease in the prevalence of IgE-mediated/allergic asthma, but increased prevalence of non-IgE-mediated asthma/asthma symptoms. Chronic bronchitis and COPD are common in the adult workers and lead to increased respiratory morbidity and mortality. Given the large spectrum of potential respiratory diseases affecting exposed agriculture workers, a thorough clinical investigation is warranted to appropriately phenotype the respiratory disease in order to facilitate appropriate therapy.

CAUSATIVE FACTORS IN LARGE ANIMAL FARMING ENVIRONMENTS

Allergens

There are a number of environmental factors that might lead to adverse respiratory health outcomes in workers exposed to large animal feeding operations, including allergens, toxic gases, organic dusts, and microbial cell wall components. Allergens are antigens defined as macromolecular agents capable of inducing IgE immune responses and provoking allergic reactions in sensitized subjects. In relation to farming, macromolecules that

TABLE 1. Respiratory Diseases Associated with Large Animal Farming

Disease	Symptoms	Diagnosis	Subjects Studied	Pathophysiology	Source
Rhinitis	Congestion, rhinorrhea, sneezing, pruritus	Defined by the presence of one or more symptoms	Swine veterinarians Agricultural workers - Livestock breeding - Livestock handling - Dairy production	Neutrophil influx IL-8/CXCL8, IL-6 levels	(Slager et al., 2010) (Poole et al., 2007) (Dosman et al., 2000)
Mucous membrane syndrome	Combination of nasal, eye and throat complaints with congestion, rhinorrhea and pruritus	No defined criteria, based on reported symptoms in confinement workers	Swine confinement workers	Increased levels of IL-1 α , IL-1 β , IL-6	(Romberger et al., 2001) (Kirkhorn et al., 2000) (Von Essen et al., 1999)
Asthma	Variable and reoccurring: wheeze, cough, chest tightness, shortness of breath	History of intermittent symptoms with either 1) reversible airflow obstruction by spirometry or peak flow monitoring, or 2) positive bronchoprovocation test	Children of farmers Farmers Agricultural workers	IgE-mediated eosinophilic influx Occupational Workplace exacerbated	(Lawson et al., 2011) (Riedler et al., 2000) (Merchant et al., 2005) (Eduard et al., 2009) (Kogevinas et al., 1999) (Dosman et al., 2004) (Tarlo, 2003)
Asthma-like syndrome	Cough, chest tightness, dyspnea, wheezing	Symptoms similar to asthma Positive bronchoprovocation if symptomatic No obstruction noted on PFT No eosinophil influx	Swine confinement workers Grain elevator operators	Normal pulmonary function testing Airway hyper-responsiveness to methacholine challenge Increased Neutrophils	(Kirkhorn et al., 2000) (Romberger et al., 2001) (Von Essen et al., 2010)
Chronic bronchitis COPD	Productive cough for at least 3 mo/yr for at least 2 consecutive years	Symptoms of chronic cough, sputum production, dyspnea or wheezing Known risk factor exposure FEV ₁ <80 % predicted, FEV ₁ /FVC <70% predicted, decreased DLCO	Farmers - Swine - Multiple livestock - Dairy	Absence of eosinophilic infiltrates Organic dust: multiple agents - Muramic acid - Endotoxins - Storage mites - Ammonia	(Eduard et al., 2009) (Omland et al., 2002) (Zhiping et al., 1996)
Organic dust toxic syndrome	Fever, malaise, myalgia, chest tightness, headache, nausea	Flu-like illness Occurs 4–8 h after exposure No sensitization Resolution of symptoms within days	Farmers - Grain - Swine Confinement Workers	Symptoms start 4–8 h after exposure Neutrophilia Neutrophils on BALF Normal imaging, Normal blood gases	(Romberger et al., 2001) (Donham et al., 1990) (Vogelzang et al., 1999) (Von Essen et al., 1990) (May, 1986)
Hyper-sensitivity pneumonitis	Fever, malaise, nausea, chest tightness, headache	Acute exposure - Flu-like illness - Occurs 4–8 h after exposure - CXR with infiltrates - Hypoxia Sensitization occurs Chronic exposure - Exposure history - Interstitial disease	Farmers: - Moldy grain - Moldy vegetables - Cattle feed	Specific antigenic exposure with antibody response Lymphocytes influx Granuloma formation Production of TNF, IFN- γ , IL-12, IL-18	(Girard et al., 2010) (Kirkhorn et al., 2000) (Zhiping et al., 1996)

were shown to be antigenic include animal dander, pollens, insect fragments, storage mites, and fungal molds (Kirkhorn and Garry, 2000).

Organic Dust

Organic dust is a complex mixture of an abundant variety of substances that includes dust particles of various sizes and microbial cell wall components. Forty percent of these dust particles are in the respirable range with a median diameter of 4 μm or less, and may be deposited at the level of the terminal bronchioles and alveoli (Kilburn, 1984; Kirkhorn and Garry, 2000). In swine confinement buildings, the proportion of dust within the respirable range may be as high as 60% of the total dust (Kirkhorn and Garry, 2000). It was reported that dust levels are highest in finishing buildings, which are buildings that house mature hogs prior to going to market (Kirkhorn and Garry, 2000). Dust concentration in confinement buildings was shown to induce adverse pulmonary effects at concentrations above 2.4–2.5 mg/m^3 for swine and 0.16 mg/m^3 for poultry. At these concentrations, workers demonstrate a 5% decrease in postshift FEV_1 measurements (Donham et al., 1995; Reynolds et al., 1996). Moreover, the size of the respirable fraction may be important in disease development in large animal feeding operations. A recent study examined inflammation produced by different particulate matter (PM) size from dairy farms in California (Vogel et al., 2012). The dust collected from 5 different farms was divided into 4 different size groups and showed that PM with a cutoff of 4.2 μm induced the highest proinflammatory cytokine release from macrophages as compared to smaller (cutoff of 2.1 μm) and larger (cutoff of 10.2 μm) PM (Vogel et al., 2012). In addition, PM from the farms induced a greater proinflammatory mediator response than the same size PM collected from urban sites (Vogel et al., 2012). Similarly, other studies found that coarse PM (2.5–10 μm) induced enhanced inflammatory responses from alveolar macrophages in vitro more than fine and ultrafine PM collected from urban areas, and

further evaluation found microbial agents in the coarse PM were the major component (Becker et al., 2002, 2003; Ghio et al., 2012). However, Alexis and colleagues (2006) found that coarse particulate, even with neutralizing the microbial components, induced macrophage inflammation more than fine and ultrafine PM, suggesting the size of PM is an important factor. Future studies need to investigate the interplay between microbial components, dust size fractions, and inflammatory potential related to dusts from animal confinement operations.

Endotoxin

Organic dusts are rich in endotoxins from the cell wall of gram-negative bacteria (Mayeux, 1997). Several studies associate endotoxin levels in swine confinement structures, livestock farming, and animal feed industry with adverse respiratory health outcomes (Donham et al., 1995; Dosman et al., 2006; Reynolds et al., 1996; Schwartz et al., 1995; Vogelzang et al., 1998). In humans, inhalation and intravenous administration of lipopolysaccharide (LPS), a large molecule found in the outer membrane of gram-negative bacteria consisting of a lipid and a polysaccharide commonly referred to as endotoxin, precipitate multiple clinical inflammatory effects including fever, shivering, arthralgia, influenza-like syndrome, dry cough, dyspnea, chest tightness, and leukocytosis (American Thoracic Society, 1998; Mayeux, 1997). LPS administration also results in an acute, dose-dependent decrease in lung function and in diffusion capacity, and there is resultant airway obstruction (American Thoracic Society, 1998). Concentrations of endotoxin at 1000–2000 ng/m^3 may produce ODTs and at concentrations of 100–200 ng/m^3 may induce bronchoconstriction (American Thoracic Society, 1998; Kirkhorn and Garry, 2000). However, there is not a universal association with levels of endotoxin within swine confinement facilities and respiratory health consequences because studies reported high exposure without symptoms or low exposure with a possible dose-response relationship (Rask-Andersen et al., 1989). Moreover, Sundblad

and colleagues (2009) found that inhalation endotoxin challenges with endotoxin only did not reproduce proinflammatory responses similar to those of swine barn exposure challenges, even though the pure endotoxin concentration was 200-fold higher than the endotoxin concentration measured within the swine barns.

Peptidoglycans

Because human studies have not shown a universal association with endotoxin and respiratory disease outcomes, and laboratory studies demonstrated that organic dusts from large animal feeding operations scrubbed of endotoxin retain significant inflammatory ability (Poole et al., 2008; Romberger et al., 2002), recent studies searched for potentially important, non-endotoxin agents. Culture-dependent and culture-independent techniques demonstrated a high burden of gram-positive bacteria (10^8 16S rRNA gene copies per cubic meter of air), rather than gram-negative bacteria, in organic dust samples from swine confinement buildings (Nehme et al., 2009). Applied molecular approaches found that the majority (93.8% of sequences related to gram-positive anaerobic bacteria) of the bacteria in organic dust from swine facilities were anaerobic gram-positive bacteria with the most likely source being manure (Nehme et al., 2009). Larsson and colleagues (1999) found that gram-positive bacteria elicit inflammation in alveolar and epithelial cells (Larsson et al., 1999). Moreover, a higher concentration of total bacteria was found through these techniques as compared with standard culture-dependent identification techniques (Nehme et al., 2008). Actinomycetes and molds were also reported (Letourneau et al., 2010a, 2010b). Peptidoglycan is derived predominately from gram-positive bacteria as it is 85% of the cell wall, but it is also found in gram-negative bacteria, making up approximately 5% of the cell wall. Muramic acid, the basic backbone of peptidoglycan, is detected by mass spectrometry techniques (Szponar and Larsson, 2001). Levels of muramic acid have been associated with airway inflammation in European swine farmers. High concentrations

of muramic acid were reported in swine confinement facilities (15 ng/mg) and dairy barns (5 ng/mg) in the United States (Poole et al., 2010). These concentrations were statistically higher than comparative environments, such as grain elevators or domestic houses, but their association with airway disease manifestations in U.S. agriculture workers is not known. Poole and colleagues (2011b) demonstrated that exposure to peptidoglycan alone elicits proinflammatory cellular mediator release and airway inflammatory consequences in rodents similar to that observed with swine facility organic dust extracts.

Animal Confinement Gases

Large animal feeding operations release high levels of gases due to by-products of animal waste, primarily manure. Hydrogen sulfide (H_2S) is found in both swine and dairy confinement building and, although uncommonly, may lead to death from asphyxiation and/or pulmonary edema while cleaning manure pits in buildings with under-building manure pits (CDC, 1993). H_2S at lower levels has an odor of "rotten egg smell," but at levels that produce pulmonary edema or unconsciousness, the gas is odorless due to olfactory paralysis (Kirkhorn and Garry, 2000). Studies linked ammonia exposure of greater than 2 h/d for 6 yr to sinusitis and chronic obstructive pulmonary conditions (Von Essen and Donham, 1999). The threshold limit value for ammonia is 25 ppm and this level is rarely exceeded in swine confinement buildings, but studies documented decrease in FEV_1 at levels above 7.5 ppm in swine confinement workers (Von Essen and Romberger, 2003).

In interim summary, recent advances into understanding the causative factors eliciting respiratory disease within large animal confinement exposures have identified multiple important components as opposed to a single agent. These components included allergens, toxic gases, particulates, organic dust, and microbial cell wall products. These environments are now recognized for the wide diversity of microbes, particularly focused on the influence

of gram-negative endotoxins and gram-positive peptidoglycans. Future research to investigate the association of particle size with microbial components human respiratory disease consequences from large animal farming exposure environments is warranted.

INNATE IMMUNITY

To help determine biologic relevance of these microbial components known as pathogen associated molecular patterns (PAMP) such as endotoxin and peptidoglycan, studies targeted specific pattern recognition receptors (PRR) to determine functional significance. Of the many PRR, Toll-like receptor (TLR) 2 recognizes gram-positive bacteria through peptidoglycans, lipoteichoic acid, and lipoproteins (Akira et al., 2006) and the TLR4 complex recognizes gram-negative bacteria endotoxins (Akira et al., 2006). In general, activation of TLR results in signaling through common intracellular signaling molecules (i.e., MyD88) and transcription factors (i.e., nuclear factor [NF]- κ B, MAP kinases) to elicit production of proinflammatory mediators including IL-1, IL-6, IL-8, and tumor necrosis factor (TNF)- α (Senthilselvan et al., 2008, 2009).

Toll-Like Receptor 4

The TLR4 complex includes TLR4, CD14, and LPS binding protein (LBP). Polymorphisms of the TLR4 gene (299 and 399) were shown to alter the signaling pathway such that changes in pulmonary function are reduced, but IL-6 and IL-8 levels are not downregulated, following exposure to LPS in human studies (Senthilselvan et al., 2009). Further, humans with the variant in TLR4 (299/399) demonstrated a decrease in cross-shift FEV₁ following a high endotoxin swine barn exposure challenge as compared to wild-type humans ($-8.48 \pm 1.52\%$ vs. $-11.46 \pm 1.79\%$, $p = .0001$), but no marked difference after a low endotoxin swine barn exposure challenge (Senthilselvan et al., 2009). In addition, CD14 is a high-affinity receptor for endotoxin and transfers endotoxin to TLR4. Polymorphisms of CD14 (CD14-159TT) were

associated with elevated levels of soluble CD14 (LeVan, 2005), and LeVan et al. (2005) found that these polymorphisms in nonsmoking farmers (CD14/-1619GG and CD14/159TT) were associated with diminished FEV₁ and decreased self-reporting of wheeze. Finally, in animal studies, mice deficient in TLR4 signaling demonstrated a reduction in neutrophil influx, but not cytokine production, following a 1-d exposure to swine barn air (Charavaryamath et al., 2008). However, these TLR4-deficient animals were not protected from developing acute airway hyperresponsiveness following swine barn air exposure.

Toll-Like Receptor 2

Investigations directed at determining the functional importance of gram-positive bacteria in organic dust environments focused on the TLR2 signaling pathway in cell culture and animal studies. TLR2 gene and protein expression is increased on airway epithelial cells following incubation with swine facility organic dust extract in a concentration-dependent manner (Bailey et al., 2008), and organic dust-induced airway epithelial cell cytokine release is dampened with anti-TLR2 antibody blockade (von Scheele et al., 2010). Further, TLR2-deficient mice displayed an altered airway response following organic dust exposure marked by a reduction in dust-induced airway inflammatory cell influx (approximate 55% reduction) and cytokine/chemokine production (Poole et al., 2011b). However, similar to observations in TLR4-deficient animals, TLR2 mice are not protected from the enhanced organic dust-induced airway hyperresponsiveness. In human studies, there appears to be a role for TLR2 polymorphisms. Eder and colleagues (2004) found that children of farmers with polymorphism in TLR2 gene (TLR2/-16934) have less atopy (14% vs. 27%, $p = .023$), IgE-mediated disease (3% vs. 14%, $p = .01$) and asthma (3% vs. 13%, $p = .012$) as compared to farm children without the polymorphism. However, this same polymorphism was not protective against atopy in children who did not live on a farm.

Other PRR

Nucleotide oligomerization domains (NOD)-like molecules are intracellular molecules that sense and respond to bacterial components. Less has been described about the role of NOD in exposed large animal farmers. However, NOD2, which recognizes muramyl dipeptide, is upregulated in antigen presenting cells following organic dust exposure, and mice deficient in NOD2 display an increased, as opposed to decreased, airway inflammatory response following swine confinement facility organic dust exposures (Poole et al., 2011a). In addition, gene polymorphisms in NOD1, which recognizes γ -D-Glu-mDAP from gram-negative bacteria and are also known as Caspase Recruitment Domain (CARD) 4, have been associated with respiratory disease in farm children. Children of farmers with the T allele in CARD4/-21596 display decreased atopy (5.8% vs. 19.4%, $p < .01$) and IgE-mediated diseases (hay fever 1.7% vs. 13%, $p < .01$; asthma 1.7% vs. 7.6%, $p < .05$) compared to reference group (Eder et al., 2006).

Intracellular Signaling Pathways

Although investigations focused on upstream innate immune receptor events found roles for TLRs NODs, there have been several studies determining relevant and potentially important intracellular signaling molecules. Protein kinase C (PKC) is an important catalytic enzyme that mediates proinflammatory cytokine release from human bronchial epithelial cells following swine confinement feeding operation and cattle feed lot organic dust exposure (Romberger et al., 2002; Wyatt et al., 2007). Further, activation of specific PKC isoforms in a sequential manner was described (Wyatt et al., 2010). In human bronchial epithelial cells, PKC α is rapidly activated by 1 h with dust exposure, which subsequently leads to activation of PKC ϵ . Inhibition of PKC α resulted in reduced dust-induced IL-6 and IL-8 release, whereas inhibition of PKC ϵ leads to less dust-induced IL-8 release. Adhesion of lymphocytes to epithelial cells, epithelial cell migration, and

wound healing in the setting of dust exposure is dependent on PKC (Mathisen et al., 2004; Slager et al., 2007). NF- κ B is a known signaling pathway that is activated by TLR4 receptor recognition and hence induces inflammation with exposure to gram-negative bacteria. It was shown that NF- κ B has enhanced binding and reporter activity when macrophages are treated with organic PM from dairy farms (Vogel et al., 2012).

In interim summary, several innate immune receptors/sensors including TLR2, TLR4, NOD1, and NOD2 are implicated in recognizing and mediating large animal confinement organic dust exposure induced inflammatory consequences (Table 2). These studies also underscore the complexity of the environmental exposure because targeting one receptor alone does not explain the entirety of disease inflammatory manifestations. Therefore, further studies to understand the host response to this diverse exposure environment is needed with a focus suggested on common signaling pathways or multiple receptor targeted approaches.

ACUTE AND CHRONIC INFLAMMATORY-ADAPTATION RESPONSE

Acute Inflammatory Response

The acute inflammatory response following swine confinement facility organic dust exposure has been well characterized. Collectively, swine confinement organic dust exposure in humans and rodents shows an acute inflammatory response marked by neutrophil influx and TNF- α , IL-6, and IL-8/CXCL8 release in both the upper and lower respiratory tract (Larsson et al., 1994, 1997, 2002; Palmberg et al., 2002; Poole et al., 2009; Sundblad et al., 2009; Von Essen and Romberger, 2003). Specifically, healthy volunteer subjects with no prior agriculture exposure demonstrated a 100-fold increase in neutrophils and a 3- to 4-fold rise in lymphocytes and macrophages in bronchoalveolar lavage fluid (BALF) following a swine barn exposure challenge (Sundblad et al., 2009). Exhaled nitric oxide was numerically

TABLE 2. Immune Signaling Pathways Implicated in Mediating Large-Animal Confinement Exposure-Induced Respiratory Diseases

Implicated pathways	Associated response and/or disease
IgE-mediated/allergen-induced	Rhinitis Asthma
Toll-like receptor pathways - TLR4 - CD14 - TLR2	Innate immune response to microbial cell wall products from gram-negative (TLR4) and gram-positive (TLR2) bacteria Lung-function decline Lower-respiratory-tract symptoms Absence of TLR2 or TLR4 associated with decreased inflammation following organic dust challenges in mice Human genetic variants: Cross-shift changes in FEV ₁ (TLR4-299-399 variant) Decreased FEV ₁ (CD14/-1619GG and CD14/159TT) Decreased asthma (TLR2/-16934)
Intracellular nucleotide oligomerization domain (NOD) sensors - NOD1 - NOD2	Absence of NOD2 associated with increased inflammation following organic dust challenges in mice. Human genetic variants: Decreased asthma (NOD1, T allele in CARD4/-21596)
Activation of PKC α , PKC ϵ	Mediates IL-6 and IL-8 release in airway epithelial cells Important in wound healing, epithelial cell migration, and adhesion in dust-exposed epithelial cells
Activation of NF- κ B	Enhanced binding and activity following dust exposures Important transcription factor in TLR signaling

elevated in controls compared to farmers when exposed to dust ($11.7 \text{ ppb} \pm 0.6 \text{ SEM}$ vs. 10.2 ± 1.6 ; $p = .023$) and exhaled nitric oxide measures almost doubled from baseline readings in healthy controls following a swine barn exposure ($p = .002$) (Sundblad et al., 2009). Finally, bronchial hyperresponsiveness to methacholine challenge occurs in healthy volunteer subjects following swine barn exposure challenges (Ek et al., 2005; Larsson et al., 2002; Sundblad et al., 2002).

Chronic Inflammatory Adaptation Response

Less is known regarding the chronic inflammatory response to large animal farming environmental exposures observed in agriculture workers. Schwartz et al. (1995) showed that there is basement membrane thickening with increase in fibronectin and hyaluronic acid in swine farmers. It is now appreciated that following repetitive or chronic exposure to these organic dust environments, the airway inflammatory response is less pronounced as compared to naive subjects, but yet exposed agriculture workers experience lung function decline and chronic airways disease. This

phenomenon has been termed the chronic inflammation adaptation response (Palmberg et al., 2002; Pedersen et al., 1996; Sundblad et al., 2009; Von Essen and Romberger, 2003).

In support of the chronic inflammatory adaptation response, studies demonstrated that bronchial responsiveness to methacholine challenge was significantly higher in naive, healthy controls as compared to swine farmers following an organic dust exposure challenge. Farmers displayed less exposure-induced symptoms of chills, runny nose, cough, and chest tightness (Palmberg et al., 2002; Sundblad et al., 2009; Von Essen and Romberger, 2003). Moreover, baseline BALF sampling from swine confinement workers demonstrated increased numbers of neutrophils, macrophages, and levels of TNF- α , IL-6, and IL-8 as compared to healthy controls, but the magnitude of the inflammatory response following an environmental challenge was less pronounced in these farmers as compared to naive individuals (Sundblad et al., 2009). The mechanisms to explain the chronic inflammation adaptation response are not entirely clear. L-Selectin may play an important role because shedding of this molecule into its soluble form is associated with a decrease in

inflammation, and swine farmers have elevated serum L-selectin, which suggests its role in the adaptation or tolerant response (Israel-Assayag and Cormier, 2002).

Cell culture modeling revealed that repetitive exposure to swine confinement organic dust extracts impairs the development of human macrophages marked by downregulation of (1) innate immune cell surface markers such as MHC class II and costimulatory molecule, CD80 and CD86 expression, (2) impaired phagocytic and bactericidal ability, and (3) diminished cytokine/chemokine responsiveness, which was also found to be independent of endotoxin (Poole et al., 2007b, 2008). Animal models have been developed that resemble the human chronic inflammatory adaptation response. For example, rodents demonstrate an increase in airway hyperresponsiveness to a one-time exposure of swine barn air and/or dust extracts, which was not observed following repetitive exposures. Further, acute exposure-induced cytokine/chemokine significantly diminishes with repetitive exposures; however, repetitive dust exposures induce histopathologic changes in the lung parenchyma, evidenced by an influx of neutrophils, T cells, B cells, and macrophages dependent on length of exposure (Charavaryamath et al., 2008; Poole et al., 2009). Th17 skewing is proposed because acute organic dust exposure increases soluble IL-17 and IL-17A-expressing lymphocytes in bronchoalveolar lavage fluid (BALF) of healthy volunteers (Ivanov et al., 2005) and repetitive organic dust exposure promotes a Th1/Th17 lung microenvironment in rodent lungs (Poole et al., 2012).

Overall, the acute inflammatory response following large animal confinement organic dust exposure is well characterized and is marked by influx of neutrophils and lymphocytes and proinflammatory cytokine/chemokine release. With repetitive or chronic exposures, there is a dampening of the acute inflammatory response, consistent with an adaptation-like phenomenon. This chronic response appears to be, in part, mediated by T-cell responses,

particularly a polarized Th1/Th17 state. Future investigations to further understand the chronic inflammatory-adaptation response and to investigate therapies to reduce disease burden in chronically exposed workers are warranted.

CONCLUSION

Since the late 1990s, there has been knowledge of multiple respiratory diseases associated with working in animal confinement environments including rhinitis, chronic bronchitis, COPD, asthma, asthma-like, ODS, and hypersensitivity pneumonia. Although the development of IgE-mediated respiratory disease is decreased in farming children, reports show that non-IgE-mediated respiratory diseases are prevalent in children and adults exposed to modern large animal feeding operations, and that large animal farming is a strong risk factor toward the development of occupational asthma diseases. Recent studies highlight that confinement workers have an increased prevalence of rhinitis (Poole et al., 2007; Slager et al., 2010) and non-IgE mediated occupational asthma (Dosman et al., 2004, 2006). Importantly, Merchant and colleagues (2005) demonstrated that children exposed to swine confinement environments in the United States had an increased prevalence of non-IgE-mediated asthma symptoms, approaching 50% of the study population. The agents responsible for respiratory diseases are not completely understood, but it is increasingly appreciated that these environments are complex and include endotoxins, peptidoglycans, and respirable dust particles. Advances into understanding specific causative factors determined that PM size is important, with coarse PM inducing enhanced inflammation (Vogel et al., 2012). Endotoxin, signaling through TLR4 (Charavaryamath et al., 2008), remains an important causative agent, but recent studies show important roles for gram-positive bacteria (Nehme et al., 2009; Poole et al., 2008, 2010) and for targeting the TLR2 pathway. Finally, advances in understanding the chronic inflammatory adaptation response in humans and animal modeling should provide the tools

to implement strategies to ultimately prevent and/or reduce respiratory disease burden in exposed workers.

REFERENCES

- American Thoracic Society. 1998. Respiratory health hazards in agriculture. *Am. J. Respir. Crit. Care Med.* 158: S1–S76.
- Akira, S., Uematsu, S., and Takeuchi, O. 2006. Pathogen recognition and innate immunity. *Cell* 124: 783–801.
- Alexis N. E., Lay, J. C., Zeman, K., Bennett, W. E., Peden, D. B., Soukup, J. M., Devlin, R. B., and Becker, S. 2006. Biological material on inhaled coarse fraction particulate matter activates airway phagocytes in vivo in healthy volunteers. *J. Allergy Clin. Immunol.* 117: 1396–1403.
- Bailey, K. L., Poole, J. A., Mathisen, T. L., Wyatt, T. A., Von Essen, S. G., and Romberger, D. J. 2008. Toll-like receptor 2 is upregulated by hog confinement dust in an IL-6-dependent manner in the airway epithelium. *Am. J. Physiol. Lung Cell Mol. Physiol.* 294: L1049–L1054.
- Becker, S., Fenton, M. J., and Soukup, J. M. 2002. Involvement of microbial components and toll-like receptors 2 and 4 in cytokine responses to air pollution particles. *Am. J. Respir. Cell Mol. Biol.* 27: 611–618.
- Becker, S., Soukup, J. M., Sioutas, C., and Cassee, F. R. 2003. Response of human alveolar macrophages to ultrafine, fine, and coarse urban air pollution particles. *Exp. Lung Res.* 29: 29–44.
- Braun-Fahrlander, C. 2003. Environmental exposure to endotoxin and other microbial products and the decreased risk of childhood atopy: Evaluating developments since April 2002. *Curr. Opin. Allergy Clin. Immunol.* 3: 325–329.
- Braun-Fahrlander, C., Gassner, M., Grize, L., Neu, U., Sennhauser, F. H., Varonier, H. S., Vuille, J. C., and Wuthrich, B. 1999. Prevalence of hay fever and allergic sensitization in farmer's children and their peers living in the same rural community. SCARPOL team. Swiss study on childhood allergy and respiratory symptoms with respect to air pollution. *Clin. Exp. Allergy* 29: 28–34.
- Burch, J. B., Svendsen, E., Siegel, P. D., Wagner, S. E., Von Essen, S., Keefe, T., Mehaffy, J., Martinez, A. S., Bradford, M., Baker, L., Cranmer, B., Saito, R., Tessari, J., Linda, P., Andersen, C., Christensen, O., Koehncke, N., and Reynolds, S. J. 2010. Endotoxin exposure and inflammation markers among agricultural workers in Colorado and Nebraska. *J. Toxicol. Environ. Health A* 73: 5–22.
- Centers for Disease Control. 1993. Fatalities attributed to entering manure waste pits—Minnesota, 1992. *Morbidity and Mortality Weekly Rep.* 42(17): 325–329. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/00020468.htm> (accessed April 21, 2012).
- Charavaryamath, C., Juneau, V., Suri, S. S., Janardhan, K. S., Townsend, H., and Singh, B. 2008. Role of toll-like receptor 4 in lung inflammation following exposure to swine barn air. *Exp. Lung Res.* 34: 19–35.
- Chen, Y., Rennie, D., Cormier, Y., McDuffie, H., Pahwa, P., and Dosman, J. 2007. Reduced risk of atopic sensitization among farmers: The Humboldt study. *Int. Arch. Allergy Immunol.* 144: 338–342.
- Cleave, J., Willson, P. J., Town, J., and Gordon, J. R. 2010. Fractionation of swine barn dust and assessment of its impact on the respiratory tract following repeated airway exposure. *J. Toxicol. Environ. Health A* 73: 1090–1101.
- Cormier, Y., Duchaine, C., Israel-Assayag, E., Bedard, G., Laviolette, M., and Dosman, J. 1997. Effects of repeated swine building exposures on normal naive subjects. *Eur. Respir. J.* 10: 1516–1522.
- Donham, K. J., Reynolds, S. J., Whitten, P., Merchant, J. A., Burmeister, L., and Popendorf, W. J. 1995. Respiratory dysfunction in swine production facility workers: Dose-response relationships of environmental exposures and pulmonary function. *Am. J. Ind. Med.* 27: 405–418.
- Dosman, J. A., Lawson, J. A., Kirychuk, S. P., Cormier, Y., Biem, J., and Koehncke, N. 2006. Three new cases of apparent

- occupational asthma in swine confinement facility employees. *Eur. Respir. J.* 28: 1281–1282.
- Dosman J. A., Lawson, J. A., Kirychuk, S. P., Cormier, Y., Biem, J., and Koehncke, N. 2004. Occupational asthma in newly employed workers in intensive swine confinement facilities. *Eur. Respir. J.* 24: 698–702.
- Dosman, J. A., Senthilselvan, A., Kirychuk, S. P., Lemay, S., Barber, E. M., Willson, P., Cormier, Y., and Hurst, T. S. 2000. Positive human health effects of wearing a respirator in a swine barn. *Chest* 118: 852–860.
- Eder, W., Klimecki, W., Yu, L., von Mutius, E., Riedler, J., Braun-Fahrlander, C., Nowak, D., Martinez, F. D., and ALEX Study Team. 2004. Toll-like receptor 2 as a major gene for asthma in children of European farmers. *J. Allergy Clin. Immunol.* 113: 482–488.
- Eder, W., Klimecki, W., Yu, L., von Mutius, E., Riedler, J., Braun-Fahrlander, C., Nowak, D., Holst, O., Martinez, F. D., and ALEX Team. 2006. Association between exposure to farming, allergies and genetic variation in CARD4/NOD1. *Allergy* 61: 1117–1124.
- Eduard, W., Douwes, J., Omenaas, E., and Heederik, D. 2004. Do farming exposures cause or prevent asthma? Results from a study of adult Norwegian farmers. *Thorax* 59: 381–386.
- Eduard, W., Pearce, N., and Douwes, J. 2009. Chronic bronchitis, COPD, and lung function in farmers: The role of biological agents. *Chest* 136: 716–725.
- Ege, M. J., Mayer, M., Normand, A. C., Genuneit, J., Cookson, W. O., Braun-Fahrlander, C., Heederik, D., Piarroux, R., von Mutius, E., and GABRIELA Transregio 22 Study Group. 2011. Exposure to environmental microorganisms and childhood asthma. *N. Engl. J. Med.* 364: 701–709.
- Ek, A., Palmberg, L., Sundblad, B. M., and Larsson, K. 2005. Salmeterol does not alter increased bronchial responsiveness caused by organic dust exposure. *Chest* 128: 3038–3047.
- Ernst, P., and Cormier, Y. 2000. Relative scarcity of asthma and atopy among rural adolescents raised on a farm. *Am. J. Respir. Crit. Care Med.* 161: 1563–1566.
- Farthing, P., Rennie, D., Pahwa, P., Janzen, B., and Dosman, J. 2009. The association between farming activities and respiratory health in rural school age children. *J. Agromed.* 14: 256–262.
- Ghio, A. J., Carraway, M. S., and Madden, M. C. 2012. Composition of air pollution particles and oxidative stress in cells, tissues, and living systems. *J. Toxicol. Environ. Health B* 15: 1–21.
- Girard, M., and Cormier, Y. 2010. Hypersensitivity pneumonitis. *Curr. Opin. Allergy Clin. Immunol.* 10: 99–103.
- Hiel, D., von Scheele, I., Sundblad, B. M., Larsson, K., and Palmberg, L. 2009. Evaluation of respiratory effects related to high-pressure cleaning in a piggery with and without robot pre-cleaning. *Scand. J. Work Environ. Health* 35: 376–383.
- Israel-Assayag, E., and Cormier, Y. 2002. Adaptation to organic dust exposure: A potential role of L-selectin shedding? *Eur. Respir. J.* 19: 833–837.
- Ivanov, S., Palmberg, L., Venge, P., Larsson, K., and Linden, A. 2005. Interleukin-17A mRNA and protein expression within cells from the human bronchoalveolar space after exposure to organic dust. *Respir. Res.* 6: 44.
- Key, N., and McBride, W. D. 2008. Amber waves. The economics of food, farming, natural resources, and rural America. Technology, larger farm size increased productivity on U.S. hog farms. Available at <http://webarchives.cdlib.org/sw1vh5dg3r/http://ers.usda.gov/AmberWaves/April08/Features/USHogFarms.htm> (accessed May 10, 2012).
- Kilburn, K. H. 1984. Particles causing lung disease. *Environ. Health Perspect.* 55: 97–109.
- Kirkhorn, S. R., and Garry, V. F. 2000. Agricultural lung diseases. *Environ. Health Perspect.* 108: 705–712.
- Klinberg, B., Berglund, N., Lilja, G., Wickman, M., and van Hage-Hamsten, M. 2001. Fewer allergic respiratory disorders among farmers' children in a closed birth cohort from Sweden. *Eur. Respir. J.* 17: 1151–1157.

- Kogevinas, M., Anto, J. M., Sunyer, J., Tobias, A., Kromhout, H., and Burney, P. 1999. Occupational asthma in europe and other industrialised areas: A population-based study. european community respiratory health survey study group. *Lancet* 353: 1750–1754.
- Larsson, B. M., Larsson, K., Malmberg, P., and Palmberg, L. 2002. Airways inflammation after exposure in a swine confinement building during cleaning procedure. *Am. J. Ind. Med.* 41: 250–258.
- Larsson B. M., Larsson, K., Malmberg, P., and Palmberg, L. 1999. Gram positive bacteria induce IL-6 and IL-8 production in human alveolar macrophages and epithelial cells. *Inflammation* 23: 217–230.
- Larsson, B. M., Palmberg, L., Malmberg, P. O., and Larsson, K. 1997. Effect of exposure to swine dust on levels of IL-8 in airway lavage fluid. *Thorax* 52: 638–642.
- Larsson, K. A., Eklund, A. G., Hansson, L. O., Isaksson, B. M., and Malmberg, P. O. 1994. Swine dust causes intense airways inflammation in healthy subjects. *Am. J. Respir. Crit. Care Med.* 150: 973–977.
- Lawson, J. A., Dosman, J. A., Rennie, D. C., Beach, J., Newman, S. C., and Senthilselvan, A. 2011. The association between endotoxin and lung function among children and adolescents living in a rural area. *Can. Respir. J.* 18: e89–e94.
- Lawson, J. A., Dosman, J. A., Rennie, D. C., Beach, J., Newman, S. C., and Senthilselvan, A. 2011. Relationship of endotoxin and tobacco smoke exposure to wheeze and diurnal peak expiratory flow variability in children and adolescents. *Respirology* 16: 332–339.
- Letourneau, V., Nehme, B., Meriaux, A., Masse, D., Cormier, Y., and Duchaine, C. 2010a. Human pathogens and tetracycline-resistant bacteria in bioaerosols of swine confinement buildings and in nasal flora of hog producers. *Int. J. Hyg. Environ. Health* 213: 444–449.
- Letourneau, V., Nehme, B., Meriaux, A., Masse, D., and Duchaine, C. 2010b. Impact of production systems on swine confinement buildings bioaerosols. *J. Occup. Environ. Hyg.* 7: 94–102.
- LeVan, T. D., Von Essen, S., Romberger, D. J., Lambert, G. P., Martinez, F. D., Vasquez, M. M., and Merchant, J. A. 2005. Polymorphisms in the CD14 gene associated with pulmonary function in farmers. *Am. J. Respir. Crit. Care Med.* 171: 773–779.
- Mathisen, T., Von Essen, S. G., Wyatt, T. A., and Romberger, D. J. 2004. Hog barn dust extract augments lymphocyte adhesion to human airway epithelial cells. *J. Appl. Physiol.* 96: 1738–1744.
- May, J. J., Stallones, L., Darrow, D., and Pratt, D. S. 1986. Organic dust toxicity (pulmonary mycotoxicosis) associated with silo unloading. *Thorax* 41: 919–923.
- Mayeux, P. R. 1997. Pathobiology of lipopolysaccharide. *J. Toxicol. Environ. Health* 51: 415–435.
- Merchant, J. A., Naleway, A. L., Svendsen, E. R., Kelly, K. M., Burmeister, L. F., Stromquist, A. M., Taylor, C. D., Thorne, P. S., Reynolds, S. J., Sanderson, W. T., Chrischilles, E. A. 2005. Asthma and farm exposures in a cohort of rural Iowa children. *Environ. Health Perspect.* 113: 350–356.
- Monso, E., Riu, E., Radon, K., Magarolas, R., Danuser, B., Iversen, M., Morera, J., and Nowak, D. 2004. Chronic obstructive pulmonary disease in never-smoking animal farmers working inside confinement buildings. *Am. J. Ind. Med.* 46: 357–362.
- Nehme, B., Gilbert, Y., Letourneau, V., Forster, R. J., Veillette, M., Villemur, R., and Duchaine, C. 2009. Culture-independent characterization of archaeal biodiversity in swine confinement building bioaerosols. *Appl. Environ. Microbiol.* 75: 5445–5450.
- Nehme, B., Letourneau, V., Forster, R. J., Veillette, M., and Duchaine, C. 2008. Culture-independent approach of the bacterial bioaerosol diversity in the standard swine confinement buildings, and assessment of the seasonal effect. *Environ. Microbiol.* 10: 665–675.
- O'Connor, A. M., Auvermann, B., Bickett-Weddle, D., Kirkhorn, S., Sargeant, J. M., Ramirez, A., and Von Essen, S. G. 2010.

- The association between proximity to animal feeding operations and community health: A systematic review. *PLoS One* 5: e9530.
- Omland, O. 2002. Exposure and respiratory health in farming in temperate zones—A review of the literature. *Ann. Agric. Environ. Med.* 9: 119–136.
- Palmberg, L., Larsson, B. M., Malmberg, P., and Larsson, K. 2002. Airway responses of healthy farmers and nonfarmers to exposure in a swine confinement building. *Scand. Work Environ. Health* 28: 256–263.
- Pedersen, B., Iversen, M., Bundgaard Larsen, B., and Dahl, R. 1996. Pig farmers have signs of bronchial inflammation and increased numbers of lymphocytes and neutrophils in BAL fluid. *Eur. Respir. J.* 9: 524–530.
- Pekkanen, J., Xu, B., and Jarvelin, M. R. 2001. Gestational age and occurrence of atopy at age 31—a prospective birth cohort study in Finland. *Clin. Exp. Allergy* 31: 95–102.
- Poole, J. A., Alexis, N. E., Parks, C., MacInnes, A. K., Gentry-Nielsen, M. J., Fey, P. D., Larsson, L., Allen-Gipson, D., Von Essen, S. G., and Romberger, D. J. 2008. Repetitive organic dust exposure in vitro impairs macrophage differentiation and function. *J. Allergy Clin. Immunol.* 122: 375–382.
- Poole, J. A., Dooley, G. P., Saito, R., Burrell, A. M., Bailey, K. L., Romberger, D. J., Mehaffy, J., and Reynolds, S. J. 2010. Muramic acid, endotoxin, 3-hydroxy fatty acids, and ergosterol content explain monocyte and epithelial cell inflammatory responses to agricultural dusts. *J. Toxicol. Environ. Health A* 73: 684–700.
- Poole, J. A., Kielian, T., Wyatt, T. A., Gleason, A. M., Stone, J., Palm, K., West, W. W., and Romberger, D. J. 2011a. Organic dust augments nucleotide-binding oligomerization domain expression via an NF- κ B pathway to negatively regulate inflammatory responses. *Am. J. Physiol. Lung Cell Mol. Physiol.* 301: L296–L306.
- Poole, J. A., LeVan, T. D., Slager, R. E., Qiu, F., Severa, L., Yelinek, J., Carlson, M. L., Bush, J., Bolin, N., Wyatt, T. A., Romberger, D., and Von Essen, S. G. 2007a. Bronchodilator responsiveness in swine veterinarians. *J. Agromed.* 12: 49–54.
- Poole, J. A., Wyatt, T. A., Kielian, T., Oldenburg, P., Gleason, A. M., Bauer, A., Golden, G., West, W. W., Sisson, J. H., and Romberger, D. J. 2011b. Toll-like receptor 2 regulates organic dust-induced airway inflammation. *Am. J. Respir. Cell Mol. Biol.* 45: 711–719.
- Poole, J. A., Wyatt, T. A., Oldenburg, P. J., Elliott, M. K., West, W. W., Sisson, J. H., Von Essen, S. G., and Romberger, D. J. 2009. Intranasal organic dust exposure-induced airway adaptation response marked by persistent lung inflammation and pathology in mice. *Am. J. Physiol. Lung Cell Mol. Physiol.* 296: L1085–L1095.
- Poole, J. A., Wyatt, T. A., Von Essen, S. G., Hervert, J., Parks, C., Mathisen, T., and Romberger, D. J. 2007b. Repeat organic dust exposure-induced monocyte inflammation is associated with protein kinase C activity. *J. Allergy Clin. Immunol.* 120: 366–373.
- Poole, J. A., Gleason, A. M., Bauer, C., West, W. W., Alexis, N., Reynolds, S. J., Romberger, D. J., and Kielian, T. 2012. $\alpha\beta$ T cells and a mixed Th1/Th17 response are important in organic dust-induced airway disease. *Ann. Allergy Asthma Immunol.* 109:266–273.
- Portengen, L., Sigsgaard, T., Omland, O., Hjort, C., Heederik, D., and Doekes, G. 2002. Low prevalence of atopy in young Danish farmers and farming students born and raised on a farm. *Clin. Exp. Allergy* 32: 247–253.
- Radon, K., Ehrenstein, V., Pram, G., and Nowak, D. 2004. Childhood visits to animal buildings and atopic diseases in adulthood: An age-dependent relationship. *Am. J. Ind. Med.* 46: 349–356.
- Rask-Andersen, A., Malmberg, P., and Lundholm, M. 1989. Endotoxin levels in farming: Absence of symptoms despite high exposure levels. *Br. J. Ind. Med.* 46: 412–416.
- Reed, C. E., and Milton, D. K. 2001. Endotoxin-stimulated innate immunity: A contributing factor for asthma. *J. Allergy Clin. Immunol.* 108: 157–166.
- Rennie, D. C., Lawson, J. A., Senthilselvan, A., Willson, P. J., and Dosman, J. A. 2012.

- Domestic endotoxin exposure and asthma in children: Epidemiological studies. *Front. Biosci.* 4: 56–73.
- Reynolds, S. J., Donham, K. J., Whitten, P., Merchant, J. A., Burmeister, F., and Pendorf, W. J. 1996. Longitudinal evaluation of dose-response relationships for environmental exposures and pulmonary function in swine production workers. *Am. J. Ind. Med.* 29: 33–40.
- Riedler, J., Eder, W., Oberfeld, G., and Schreuer, M. 2000. Austrian children living on a farm have less hay fever, asthma and allergic sensitization. *Clin. Exp. Allergy* 30: 194–200.
- Romberger, D. J., Bodlak, V., Von Essen, S. G., Mathisen, T., and Wyatt, T. A. 2002. Hog barn dust extract stimulates IL-8 and IL-6 release in human bronchial epithelial cells via PKC activation. *J. Appl. Physiol.* 93: 289–296.
- Schwartz, D. A., Donham, K. J., Olenchock, S. A., Pendorf, W. J., Van Fossen, D. S., Burmeister, L. F., and Merchant, J. A. 1995. Determinants of longitudinal changes in spirometric function among swine confinement operators and farmers. *Am. J. Respir. Crit. Care Med.* 151: 47–53.
- Senthilselvan, A., Dosman, J. A., Chenard, L., Burch, L. H., Predicala, B. Z., Sorowski, R., Schneberger, D., Hurst, T., Kirychuk, S., Gerds, V., Cormier, Y., Rennie, D. C., and Schwartz, D. A. 2009. Toll-like receptor 4 variants reduce airway response in human subjects at high endotoxin levels in a swine facility. *J. Allergy Clin. Immunol.* 123: 1034–1040.
- Senthilselvan, A., Rennie, D., Chenard, L., Burch, L. H., Babiuk, L., Schwartz, D. A., and Dosman, J. A. 2008. Association of polymorphisms of toll-like receptor 4 with a reduced prevalence of hay fever and atopy. *Ann Allergy Asthma Immunol* 100: 463–468.
- Slager, R. E., Allen-Gipson, D. S., Sammut, A., Heires, A., DeVasure, J., Von Essen, S., Romberger, D. J., and Wyatt, T. A. 2007. Hog barn dust slows airway epithelial cell migration in vitro through a PKC alpha-dependent mechanism. *Am. J. Physiol. Lung Cell Mol. Physiol.* 293: L1469–L1474.
- Slager, R. E., Simpson, S. L., Levan, T. D., Poole, J. A., Sandler, D. P., and JHoppin, J. A. 2010. Rhinitis associated with pesticide use among private pesticide applicators in the agricultural health study. *J. Toxicol. Environ. Health A* 73: 1382–1393.
- Sundblad, B. M., Larsson, B. M., Palmberg, L., and Larsson, K. 2002. Exhaled nitric oxide and bronchial responsiveness in healthy subjects exposed to organic dust. *Eur. Respir. J.* 20: 426–431.
- Sundblad, B. M., von Scheele, I., Palmberg, L., Olsson, M., and Larsson, K. 2009. Repeated exposure to organic material alters inflammatory and physiological airway responses. *Eur. Respir. J.* 34: 80–88.
- Szponar, B., and Larsson, L. 2001. Use of mass spectrometry for characterising microbial communities in bioaerosols. *Ann. Agric. Environ. Med.* 8: 111–117.
- Tarlo, S. M., and Liss, G. M. 2003. Occupational asthma: An approach to diagnosis and management. *Can. Med. Assoc. J.* 168: 867–871.
- Tarlo, S. M., Malo, J. L., and Third Jack Pepys Workshop on Asthma in the Workplace Participants. 2009. An official ATS proceedings: Asthma in the workplace: The third Jack Pepys Workshop on Asthma in the Workplace: Answered and unanswered questions. *Proc. Am. Thorac. Soc* 6: 339–349.
- U.S. Department of Agriculture. 2009. The economics of food, farming, natural resources, and rural America. Economic Research Service. Available at www.ers.usda.gov/briefing/cattle/background.htm
- Vogelzang, P. F., van der Gulden, J. W., Folgering, H., Kolk, J. J., Heederik, D., Preller, L., Tielen, M. J., and van Schayck, C. P. 1998. Endotoxin exposure as a major determinant of lung function decline in pig farmers. *Am. J. Respir. Crit. Care Med.* 157: 15–18.
- Vogelzang, P. F., van der Gulden, J. W., Tielen, M. J., Folgering, H., and van Schayck, C. P. 1999. Health-based selection for asthma, but not for chronic bronchitis, in pig farmers: An evidence-based hypothesis. *Eur. Respir. J.* 13: 187–189.

- Von Essen, S., and Donham, K. 1999. Illness and injury in animal confinement workers. *Occup. Med.* 14: 337–350.
- Von Essen, S., Moore, G., Gibbs, S., and Larson, K. L. 2010. Respiratory issues in beef and pork production: Recommendations from an expert panel. *J. Agromed.* 15: 216–225.
- Von Essen, S., and Romberger, D. 2003. The respiratory inflammatory response to the swine confinement building environment: The adaptation to respiratory exposures in the chronically exposed worker. *J. Agric. Safety Health* 9: 185–196.
- Von Essen, S. G., Thompson, A. B., Robbins, R. A., Jones, K. K., Dobry, C. A., and Rennard, S. I. 1990. Lower respiratory tract inflammation in grain farmers. *Am. J. Ind. Med.* 1: 75–76.
- Von Scheele, I., Larsson, K., and Palmberg, L. 2010. Budesonide enhances toll-like receptor 2 expression in activated bronchial epithelial cells. *Inhal. Toxicol.* 22: 493–499.
- Wyatt, T. A., Slager, R. E., Devasure, J., Auvermann, B. W., Mulhern, M. L., Von Essen, S., Mathisen, T., Floreani, A. A., and Romberger, D. J. 2007. Feedlot dust stimulation of interleukin-6 and -8 requires protein kinase c epsilon in human bronchial epithelial cells. *Am. J. Physiol. Lung Cell Mol. Physiol.* 293: L1163–L1170.
- Wyatt, T. A., Slager, R. E., Heires, A. J., DeVasure, J. M., Von Essen, S. G., and Romberger, D. J. 2010. Sequential activation of protein kinase C isoforms by organic dust is mediated by tumor necrosis factor. *Am. J. Respir. Cell Mol. Biol.* 42: 706–715.
- Zejda, J. E., Hurst, T. S., Rhodes, C. S., Barber, E. M., McDuffie, H. H., and Dosman, J. A. 1993. Respiratory health of swine producers. Focus on young workers. *Chest* 103: 702–709.
- Zhiping, W., Malmberg, P., Larsson, B. M., Larsson, K., Larsson, L., and Saraf, A. 1996. Exposure to bacteria in swine-house dust and acute inflammatory reactions in humans. *Am. J. Respir. Crit. Care Med.* 154: 1261–1266.