



[Oncology Letters](#)
[International Journal of Oncology](#)
[Molecular and Clinical Oncology](#)
[Experimental and Therapeutic Medicine](#)
[International Journal of Molecular Medicine](#)
[Biomedical Reports](#)
[Oncology Reports](#)
[Molecular Medicine Reports](#)
[World Academy of Sciences Journal](#)
[International Journal of Functional Nutrition](#)
[International Journal of Epigenetics](#)
[Medicine International](#)

International Journal of Oncology


[Journal Home](#)
[Current Issue](#)
[Forthcoming Issue](#)
[Special Issues](#)
[Most Read](#)
[Most Cited \(Dimensions\)](#)
[Most Cited \(CrossRef\)](#)
[Social Media](#)
[Archive](#)

Apoptosis induced by tungsten carbide-cobalt nanoparticles in JB6 cells involves ROS generation through both extrinsic and intrinsic apoptosis pathways

Authors: Jinshun Zhao, Linda Bowman, Ruth Magaye, Stephen S. Leonard, Vincent Castranova, Min Ding

[View Affiliations](#)

Published online on: February 15, 2013 <https://doi.org/10.3892/ijo.2013.1828>

Pages: 1349-1359

Metrics: Total Views: **7428** (Spandidos Publications: **7428** | PMC Statistics: **0**)

Total PDF Downloads: **1373** (Spandidos Publications: **1373** | PMC Statistics: **0**)

Cited By (CrossRef): 31 citations [View Articles](#)



34 Total citations
9 Recent citations

4.92 Field Citation Ratio
1.28 Relative Citation Ratio



Referenced in 1 policy sources

27 readers on Mendeley

[See more details](#)

Abstract

In this study, apoptosis and related signaling induced by WC-Co nanoparticles were investigated in JB6 cells and rat lung macrophages. Electron spin resonance (ESR) and fluorescent staining indicated that both WC-Co nanoparticles and fine particles stimulated reactive oxygen species (ROS) generation. Catalase exhibited an inhibitory



April 2013

Volume 42 Issue 4

Print ISSN: 1019-6439

Online ISSN: 1791-2423

[Sign up for eToc alerts](#)

[Recommend to Library](#)

Article Options

Viewing Options

[View Abstract](#)

[Full Text](#)

[Download PDF](#)

Citations

[Cite This Article](#)

[Download Citation](#)

[Create Citation Alert](#)

[Cited By](#)

Related Articles

on Spandidos Publications

Similar Articles

on Google Scholar

on Pub Med

Information

[Online Submission](#)

[Information for Authors](#)

[Language Editing](#)

[Information for Reviewers](#)

[Editorial Policies](#)

[Editorial Academy](#)

[Join Editorial Academy](#)

[Aims and Scope](#)

[Abstracting and Indexing](#)

[Bibliographic Information](#)

[Information for](#)

Reprints and
permissions
Contact the Editor

General
Information

About Spandidos

Conferences

Job Opportunities

Contact

Terms and
Conditions

cytotoxicity and apoptotic induction than fine particles. Western blot analysis showed activation of proapoptotic factors including Fas, Fas-associated protein with death domain (FADD), caspase 3, 8 and 9, BID and BAX. In addition, both cytochrome c and apoptosis-inducing factor (AIF) were upregulated and released from mitochondria to the cytoplasm. Our findings demonstrate that, on a mass basis, WC-Co nanoparticles exhibit higher cytotoxicity and apoptotic induction than fine particles. Apoptosis induced by WC-Co nanoparticles and fine particles involves both extrinsic and intrinsic apoptosis pathways.

Introduction

Nano-scale particle research has recently become a very important field in materials science. Nanoparticles (1 to 100 nm) usually have physical properties different from those of large particles (1–10 μ m). It has been found that nanoparticles exhibit a variety of novel properties, depending on particle size, including magnetic, optical, and other physical properties as well as surface reactivity [1]. Hard metals consisting of tungsten carbide (WC) and metallic cobalt (Co) particles are important industrial materials [2]. WC-Co nanoparticle composites have the potential to replace standard materials for tools and wear parts because of their increased hardness and toughness. By means of particle size reduction, the fracture toughness and wear resistance of WC-Co can be increased significantly. Research evidence indicated that WC-Co fine particle mixture was carcinogenic in exposed workers [2]. Based on the fact that nanoparticles of metallic nickel and titanium dioxide (TiO₂) cause more pronounced toxicity than fine particles [3–7], the pathogenic effects of WC-Co particles may well vary with their size.

Among the possible mechanisms of metal toxicity, apoptosis plays an important role in carcinogenesis [8]. While a number of known or suspected human carcinogenic metallic compounds have been shown to induce apoptosis, the relevance of these observations for the carcinogenic process is, however, still not completely understood [9]. Apoptosis induced by WC-Co fine particles has been reported in previous *in vitro* studies [2,9]; however, the signal pathways induced by WC-Co particles still remain elusive.

Apoptosis is a highly regulated process that is involved in physiological as well as pathological conditions [10–13]. Deregulation of apoptosis has been implicated in carcinogenesis, tumor progression and resistance of tumor cells to radio- and chemotherapy [14,15]. Malfunction of apoptosis plays an important role in many disease processes. An inefficient elimination of mutated cells may favor carcinogenesis or tumorigenesis; failure to clear inflammatory cells may prolong the inflammation because of the release of damaging histotoxins [9]. However, excessive apoptosis has been shown to contribute to pulmonary fibrosis in mice [16]. Furthermore, enhanced apoptosis may indirectly trigger compensatory cell proliferation to ensure tissue homeostasis and promote the fixation of mutagenic events or multiplication of mutated cells. Studies have demonstrated that apoptosis is also involved in pulmonary disorders, such as acute lung injury, diffuse alveolar damage, idiopathic pulmonary fibrosis, and other lung disorders caused by bleomycin, silica, endotoxins, and the deposition of immune complexes [16–19]. Inhibition of apoptosis by gene deletion strategies or by caspase inhibitors abrogate the pathologic effects of these agents [16–20], supporting the potential role of apoptosis in the inflammatory, pulmonary fibrosis, and immunopathologic disorders. Therefore, the apoptotic properties may be important for elucidating the mechanisms of adverse health effects induced by WC-Co particles.

Accordingly, the objectives of this study are to compare the difference in cytotoxicity and apoptosis induced by WC-Co nano- and fine particles, and to elucidate the mechanisms of cell death induced by WC-Co particles. This study provides insight into the role of apoptosis in the possible pathogenicity and carcinogenicity induced by WC-Co nanoparticles.

Materials and methods

Materials

WC-Co nanoparticles (99.9% pure, molecularly mixed at the ratio of 85:15

Journal Metrics

Impact Factor: **5.65**

Ranked Oncology
(total number of cites)

CiteScore: **8.2**

CiteScore Rank: **Q1:**
#52/340 Oncology

Source Normalized
Impact per Paper (SNIP):
1.045

SCImago Journal Rank
(SJR): **1.405**

Tweets by @IJOncol

International Journal of
Oncology Retweeted

 **Spandidos Publicatic**
@SpandidosP

Replying to @SpandidosP

The latest Impact Factor for International Journal of Oncology, according to Journal Citation Reports 2021 (Clarivate), is 5.650 (cf. 3.899 last year; an increase of 45%, with the journal having an IF >5 for the first time in its existence).


Jul 1, 2021

International Journal of
Oncology Retweeted

 **Spandidos Publicatic**
@SpandidosP

Replying to @SpandidosP

The overall trend in Impact Factors for Spandidos Publications Journals featured in JCR over the course of the last decade is shown in the attached graphic.



Jul 1, 2021

International Journal of
Oncology Retweeted

 **Spandidos Publicatic**
@SpandidosP

Replying to @SpandidosP

The new Impact Factors for the Spandidos Publications journals that are indexed in JCR are shown in tabulated form in the attached graphic.

minimal essential medium (EMEM) was obtained from Lonza (Walkersville, MD). Fetal bovine serum (FBS), trypsin, penicillin/streptomycin and L-glutamine were purchased from Life Technologies Inc. (Gaithersburg, MD). YO-PRO-1 [YP, 1 mM solution in dimethyl sulfoxide (DMSO)] and propidium iodide (PI, 1.0 mg/ml in water) were purchased from Invitrogen (Carlsbad, CA). Hoechst 33342, 2', 7'-dichlorodihydrofluorescein diacetate (H2DCFDA) and dihydroethidium (DHE) were purchased from Molecular Probes (Eugene, OR). Anti-h/m caspase-8 antibody was obtained from R&D Systems (Minneapolis, MN). BID and cleaved caspase 3 antibodies were purchased from Cell Signaling Technology (Beverly, MA). All other antibodies were obtained from Santa Cruz Biotechnology Co. (Santa Cruz, CA). Cell proliferation kit I (MTT assay kit) was obtained from Roche Applied Science (Penzberg, Germany). Mitochondria Staining Kit and catalase were purchased from Sigma-Aldrich (Saint Louis, MO).

Preparation of WC-Co particles

Stock solutions of WC-Co nano- or fine particles were prepared by sonification on ice using a Branson Sonifier 450 (Branson Ultrasonics Corp., Danbury, CT) in sterile PBS (10 mg/ml) for 30 sec, then kept on ice for 15 sec and sonicated again for a total of 3 min at a power of 400 W. Before use, these particles were diluted to a designed concentration in fresh culture medium. All samples were prepared under sterile conditions.

Surface area and size distribution measurements

Surface area of WC-Co particles was measured using the Gemini 2360 Surface Area Analyzer (Micromeritics, Norcross, GA) with a flowing gas technique according to the manufacturer's instructions. The size distribution of WC-Co particles was detected using scanning electron microscopy (SEM). Briefly, WC-Co particles were prepared by sonification. Then, the samples were diluted in double-distilled water and air dried onto a carbon planchet. Images were collected on a scanning electron microscope (Hitachi S-4800, Japan) according to the manufacturer's instructions. Optimas 6.5 image analysis software (Media Cybernetics, Bethesda, MD) was used to measure the diameter of WC-Co particles.

Cell culture

Mouse epidermal JB6 cells were maintained in 5% FBS EMEM containing 2 mM L-glutamine and 1% penicillin-streptomycin (10,000 U/ml penicillin and 10 mg/ml streptomycin) at standard culture conditions (37°C, 80% humidified air and 5% CO₂). For all treatments, cells were grown to 80% confluence.

Determination of free radical formation

All ESR measurements were conducted using a Varian E9 ESR spectrometer and a flat cell assembly. Hyperfine couplings were measured (to 0.1 G) directly from magnetic field separation using potassium tetraperoxochromate (K₃CrO₈) and 1,1-diphenyl-2-picrylhydrazyl as reference standards. An EPR DAP 2.0 program was used for data acquisition and analyses. Reactants were mixed in test tubes in a total final volume of 450 µl. The reaction mixture was then transferred to a flat cell for ESR measurement. The concentrations given in the figure legends are final concentrations and measurements were made at room temperature and under ambient air, except those specifically indicated otherwise. The receiver gain, time constant, sweep time, sweep width, modulation frequency, modulation amplitude and microwave power were set constant to allow relative intensity comparisons of spectra. All spectra shown are the accumulation of five scans. Hyperfine couplings constants were determined using the WinSim program of the NIEHS public EPR software tools, available over the internet (<http://epr.niehs.nih.gov>). The relative radical concentration was estimated by measuring the peak-to-peak height (mm) of the observed spectra.

H2DCFDA and DHE are used for staining general ROS or oxygen radicals (•O₂⁻) produced in intact cells, respectively. Hoechst 33342 is a nucleic acid stain. JB6 cells were seeded into a 24-well plate. Cells were grown 24 h and then starved in 0.1% FBS

International Journal of Molecular Medicine	4,351	3,686
Oncology Letters	2,967	2,513
Molecular Medicine Reports	2,952	2,190
Experimental and Therapeutic Medicine	2,447	1,760

Jul 1, 2021

International Journal of Oncology Retweeted

 **Spandidos Publications**
@SpandidosP

Following the release of Journal Citation Reports 2021 (published by Clarivate), we are pleased to announce some significant increases in the Impact Factors for several of our Journals. More information to come in the following thread.

Jul 1, 2021

International Journal of Oncology Retweeted

 **Spandidos Publications**
@SpandidosP

We would like to thank all our authors, reviewers, Editorial Boards and Academies, and office staff in Athens and London for their contributions to these achievements.

Jun 28, 2021

[Embed](#) [View on Twitter](#)

fresh 0.1% FBS EMEM. The images were captured with a fluorescence microscope (Axiovert 100M, Zeiss, Germany).

Cytotoxicity assay

Cytotoxicity of WC-Co particles to JB6 cells was assessed by an MTT assay kit following the manufacturer's instructions. Briefly, cells were plated in 100 ml EMEM at a density of 10^4 cells/well in a 96-well plate. The cells were grown for 24 h and then, treated with various concentrations of WC-Co particles. After 24-h incubation, 10 ml MTT labeling reagent was added in each well and the plates were further incubated for 4 h. Afterward, 100 ml solubilization solution was added to each well and the plate was incubated overnight at 37°C. The optical density (OD) of the wells was measured at a wavelength of 575 nm with reference of 690 nm using an ELISA plate reader. Results were calculated using the OD measured without cells.

Detection of apoptosis

YP staining was used to detect cell apoptosis in JB6 cells. Briefly, cells were seeded onto a 24-well plate overnight. Then, cells were treated with/without various concentrations of WC-Co nano- or fine particles for 24 h. Before microscopy, YP dye was added into the cultures (10 μ g/ml) and cells were further incubated for 1 h. Then, cells were washed two times with EMEM medium. Apoptotic cells were monitored using a fluorescent microscope (Axiovert 100M). Percentage of cells exhibiting apoptosis was calculated.

Sub-cellular fractionation and western blot analysis

Briefly, cells were plated onto a 100×20 mm cell culture dish. The cultures were grown for 24 h and then starved in 0.1% FBS EMEM overnight. Cells were treated with/without WC-Co nano- or fine particles. After treatment, the cells were extracted with 1X SDS sample buffer supplemented with protease inhibitor cocktail (Sigma-Aldrich). Protein concentrations were determined using the bicinchoninic acid method (Pierce, Rockford, IL). Equal amounts of proteins were separated by 4–12% Tris glycine gels. Immunoblots for expression of Fas, FADD, caspase 3, caspase 8, caspase 9, BID, cleaved BID, BAX, Bcl-2, cytochrome c, AIF, lamin A/C, β -actin and β -tubulin were detected. Experiments were performed three or more times, and equal loading of protein was ensured by measuring β -actin and β -tubulin expression.

To prepare the subcellular fractionation, after treatment cells were washed twice with cold PBS. Then, cells were resuspended in 50 μ l 1X Cytosol Extraction Buffer Mix and incubated on ice for 10 min. Cells were homogenized by passing through 22-gauge needles 30 times. The lysate was centrifuged at $700 \times g$ for 10 min at 4°C to remove unbroken cells and nuclei. The supernatant was then re-centrifuged at $10,000 \times g$ for 30 min at 4°C. The resulting supernatant was the cytosolic fraction while the pellet contained mitochondria. The cytosolic fraction was diluted using 50 μ l of 2X SDS sample buffer. The mitochondrial pellet was resuspended in 20 μ l Mitochondria Extraction Buffer Mix plus 20 μ l of 2X SDS sample buffer. These two different fractions were used for western blot analysis.

Detection of mitochondrial membrane permeability

JB6 cells were seeded onto a 24-well plate overnight. Cells were treated with/without WC-Co nano- or fine particles supplemented with or without catalase (10,000 U/ml) for 24 h. Changes in mitochondrial membrane permeability were evaluated using a mitochondrial staining kit (JC1 staining) according to the manufacturer's instructions. Briefly, a staining mixture was prepared by mixing the staining solution with an equal volume of the EMEM medium. Cells were incubated in the staining mixture (0.4 ml/well) for 30 min at 37°C in a humidified atmosphere containing 5% CO₂. Thereafter, cells were washed two times in medium, followed by addition of fresh medium. Mitochondrial membrane permeability was monitored on a fluorescence microscope (Axiovert 100M).

Isolation of rat lung macrophages

Brown Norway rats were obtained from Charles River Laboratories (Wilmington, MA). The animals were housed in AAALA-accredited facility, specific-pathogen-free,

acclimated for at least 5 days before use, and were housed in ventilated cages that were provided with HEPA-filtered air, with α -Dri virgin cellulose chips and hardwood β -chips used as bedding. The rats were maintained on ProLab 3500 diet and tap water, both of which were provided *ad libitum*. Rats were anesthetized using brevitil sodium (40 mg/kg), and exposed to 100 μ l/rat of 1% WC-Co nanoparticles suspended in sterile saline (100 μ l of saline for control rat) by intranasal droplet application 4 times (totally 4 g/rat) on day 0, 7, 14 and 21, respectively. At 24 h after the last treatment, rats were sacrificed by intraperitoneal injection of 0.2 g/kg pentobarbital, and then the descending aorta was cut. The trachea was exposed, and a cannula was inserted. Bronchoalveolar lavage fluid (BALF) was collected using saline. After centrifugation, cells in BALF were seeded into a 6-well plate in 5% FBS EMEM containing 2 mM L-glutamine and 1% penicillin-streptomycin (10,000 U/ml penicillin and 10 mg/ml streptomycin) at standard culture conditions (37°C, 80% humidified air and 5% CO₂).

After 4-h incubation, the unattached cells were washed out by using 5% FBS EMEM for 3 times. Attached cells (rat lung macrophages) were used for mitochondrial membrane permeability or apoptotic detections. Mitochondrial membrane permeability of rat lung macrophages was evaluated using a mitochondrial staining kit according to the manufacturer's instructions as stated above.

Apoptotic detection of rat lung macrophages

Dual staining using YP and PI were used to distinguish between apoptosis and necrosis (or late apoptosis) as described previously [21,22] with some modifications. Rat lung macrophages were obtained as stated above. YP and PI were added into the cultures with a final concentration of 10 μ g/ml and 1 μ M, respectively. Thereafter, rat lung macrophages were incubated in 5% FBS EMEM medium containing YP and PI dyes at 37°C in a humidified atmosphere containing 5% CO₂ for 1 h. Then, cells were washed two times in medium, followed by addition of fresh medium. Apoptotic cells were monitored using a fluorescence microscope (Axiovert 100M). YP stained cells were detected with a blue excitation filter. PI stained cells were measured using a green excitation filter.

Statistical analysis

Data are presented as means \pm standard errors (SE) of the number of experiments/samples. Data were analyzed using one-way analysis of variance (ANOVA). Significance was set at $p \leq 0.05$.

Results

Surface area and size distribution of WC-Co particles


Gemini 2360 Surface Area Analyzer and scanning electron microscopy were used to measure the surface area and size distribution of WC-Co particles, respectively. The average surface area of WC-Co nano and fine particles were published previously [23]. The average surface area of WC-Co nanoparticles was 2.73 m²/g compared to 0.16 m²/g for fine particles. The average size distribution of WC-Co nano- and fine particles is 95.53 nm and 39.00 μ m, respectively.

Free radical generation induced by WC-Co particles in JB6 cells

The ESR spin trapping technique and fluorescent staining were used to detect free radical generation during incubation of WC-Co nano- or fine particles with JB6 cells (Fig. 1). Using the ESR spin trapping technique, we observed the formation of free radicals in JB6 cells exposed to WC-Co nano- or fine particles. After 5 min exposure of JB6 cells to WC-Co particles, DMPO radical adducts were recorded. The ESR signal from WC-Co nanoparticles was much stronger than fine particles (Fig. 1A). Addition of catalase, an H₂O₂ scavenger decreased the radical generation, while addition of H₂O₂ dramatically increased the ROS adduct signal (Fig. 1A). These results suggest that ROS generated during exposure of WC-Co particles to JB6 cells was formed via a metal-dependent Fenton reaction. H2DCFDA, a general ROS sensitive dye, and DHE, an \bullet O₂⁻ specific dye, were used to monitor ROS generation induced by WC-Co particles in intact

nanoparticles induced stronger ROS generation, indicated by DHE staining, compared to fine particles (Fig. 1B).


Figure 1



Effects of WC-Co particles on cell viability and apoptotic induction

To determine whether there is a difference in the cytotoxicity induced by different sizes of WC-Co particles, various concentrations [5–100 μ g/cm 2] of WC-Co nano- or fine particles were used to study the effects on cell viability in JB6 cells using the MTT assay (Fig. 2). After 24-h treatment, both WC-Co nano- and fine particles caused a dose-dependent cytotoxicity in JB6 cells (Fig. 2A). Cytotoxicity induced by WC-Co nanoparticles was significantly higher than fine particles between the concentration of 40 and 100 μ g/cm 2 . To detect apoptosis induced by WC-Co particles, YP staining was used. Results indicated that both WC-Co nano- and fine particles induced JB6 cell apoptosis (Fig. 2B). The percentage of apoptotic cells was significantly higher in cells treated with nanoparticles than fine particles between the concentration of 40 and 100 μ g/cm 2 . At the concentration of 100 μ g/cm 2 , there was an 8-fold increase in apoptosis induced by nanoparticles compared to fine particles (Fig. 2C).


Figure 2



Effects of WC-Co particles on Fas, FADD, caspase 3, 8 and 9, lamin A/C, BID, BAX and Bcl-2

To investigate the involvement of extrinsic signals in the apoptotic process induced by WC-Co particles, expression of Fas and FADD was examined. JB6 cells were treated with WC-Co nano- or fine particles (100 μ g/cm 2) for 1, 3, 6 and 8 h, respectively, and then the protein expression patterns were detected by western blot analysis (Fig. 3). Results demonstrated that WC-Co nano- and fine particles, especially nanoparticles, stimulated Fas and FADD expression.


Figure 3



are two types of apoptotic caspases: initiator caspases and effector caspases. Initiator caspases (e.g. caspase 8) cleave inactive pro-forms of effector caspases, thereby activating them. Effector caspases (e.g. caspase 3) in turn cleave other protein substrates resulting in the apoptotic process. Our results indicated that caspase 3 and 8 were significantly cleaved and a slight activation of caspase 9 was observed. BAX and BID, two proapoptotic members of Bcl-2 family, were upregulated by WC-Co particles. Bcl-2, an anti-apoptotic factor, was also upregulated. The cellular morphology associated with the apoptotic process has been well characterized by membrane blebbing, formation of apoptotic bodies, and chromosome condensation. These apoptotic changes are the result of the cleavage of cellular proteins, such as lamin and actin [25,26]. In this study, cleavage of lamin A and β -actin were observed, especially in cells treated with WC-Co nanoparticles. The cleavage of lamin A and β -actin were detected as early as 1 h post-exposure of WC-Co nanoparticles. Interestingly, WC-Co nanoparticles induced lamin C upregulation without cleavage.

Effects of WC-Co particles on cytochrome c and AIF

The intrinsic pathway of apoptosis is initiated in the mitochondria. Its activation involves release of cytochrome *c* and other proapoptotic factors from the mitochondrial intermembrane space [27]. AIF is a proapoptotic factor, a mitochondrial protein [28]. It is normally confined to the mitochondrial intermembrane space [29]. After release from mitochondria into the cytoplasm, both cytochrome *c* and AIF can stimulate cell apoptosis by initiating the intrinsic pathway. To test the effects of WC-Co particles on the intrinsic pathway of apoptosis, JB6 cells were treated with nano- or fine particles [100 $\mu\text{g}/\text{cm}^2$] for 1, 3, 6 and 8 h, respectively. Western blots revealed that both nano- and fine particles induced cytochrome *c* and AIF upregulation as well as release from mitochondria to the cytoplasm after 1 h treatment, with nano-WC-Co exhibiting greater potency (Fig. 4A).




Effects of WC-Co particles on mitochondrial membrane permeability

Mitochondrial membrane permeability change is a hallmark for apoptosis [30]. JB6 cells were treated with or without various concentrations of WC-Co particles for 24 h. Mitochondrial membrane permeability was evaluated using a JC-1 staining kit according to the manufacturer's instructions. The results indicated that both WC-Co nano- and fine particles induced a significant change in the mitochondrial membrane permeability compared to control, with nano WC-Co exhibiting greater potency. Valinomycin (1 μM /ml culture medium) and H₂O₂ (500 μM) for 1 h were used as positive controls and Hoechst 33342 was used for nucleic localization (Fig. 4B).

Effect of catalase on WC-Co particle-induced mitochondrial membrane permeability alteration


According to the results obtained above as well as our previous studies [23], ROS may play an important role in WC-Co particle-induced mitochondrial membrane permeability alteration. To verify this hypothesis, the inhibitory effect of catalase on WC-Co particle-induced mitochondrial membrane permeability alteration was examined by using a mitochondrial staining kit (Fig. 5). The results showed that

membrane permeability damage induced by H₂O₂ (a free radical-apoptotic reagent), but not by valinomycin (a non-free radical-apoptotic reagent).



Apoptosis and mitochondrial membrane permeability alteration induced by WC-Co nanoparticles in rat lung macrophages

Apoptosis and mitochondrial membrane permeability alteration induced by WC-Co nanoparticles in rat lung macrophages were investigated as described in Materials and methods. Results indicated that lung macrophages from WC-Co nanoparticle-treated rats exhibited more apoptosis and mitochondrial membrane permeability damage compared to control (Fig. 6).



Discussion

With the increased use of nanoparticles in modern industries, inhaled nanoparticles are increasingly being recognized as a potential health threat [31]. Hard metal, a mixture of WC-Co, is an important industrial material. Occupational exposure of workers to the WC-Co particle mixture has been shown to be carcinogenic, and Co itself can also induce occupational asthma [32]. WC-Co nanoparticles were introduced in industry [33] to improve hardness and toughness compared to standard WC-Co materials. The increasing use of WC-Co nanoparticles may lead to an increased human exposure and adverse health effects [34]. The International Agency for Research on Cancer (IARC) has classified Co and its compounds as possibly carcinogenic to humans [35]. However, the molecular mechanisms involved in hard metal-induced carcinogenesis are unclear. In addition, the health effects of occupational exposure to WC-Co nanoparticles are unknown. The molecular events mediating cellular responses to WC-Co particles also remain to be elucidated.

It is well known that the toxicity of particles to the lung in both occupational and environmental settings is not only related to exposure but also to the particle size. We have previously shown that nanoparticles of TiO₂ and metallic nickel produce higher cytotoxicity than their conventional fine particles, respectively [6,7]. Accordingly, WC-Co nanoparticles may be more toxic than WC-Co fine particles.

In this study, we observed that both WC-Co nano- and fine particles induced ROS generation in a dose-dependent manner. This result is in agreement with our previous study which indicated that compared to fine particle WC-Co nanoparticles generated a higher level of free radicals and induced greater oxidative stress, as evidenced by a decrease of GSH levels [23]. Further studies indicated that catalase inhibits WC-Co particle-induced mitochondrial membrane permeability damage in JB6 cells. These results demonstrate that hydrogen peroxide may play an important role in the pathogenicity induced by WC-Co particles.

Using MTT and apoptotic assays, we observed that both WC-Co nano- and fine particles

demonstrated TiO₂ nanoparticles to be more inflammatory than fine TiO₂ particles [4]. However, when normalized to surface area, they found that the dose-response curves for the nano- and fine particles were similar, suggesting that the pulmonary toxicity might be mediated by surface effects. In the present study, surface area of WC-Co nanoparticles was 17-fold greater than fine particles. However, WC-Co nanoparticles exhibited potency for cytotoxicity and apoptosis induction which was somewhat less than 17-fold greater than fine particles. Therefore, in agreement with our previous study with TiO₂ particles [7], specific surface area measured by gas absorption, tends to over correct for the greater cytotoxicity probably because it over estimates the surface area of agglomerates.

Apoptosis and necrosis are two forms of cell death that have been defined on the basis of distinguishable, morphological criteria [36]. Apoptosis is a programmed cell death which is now widely recognized as being of critical importance in health and disease. Although studies have demonstrated that WC-Co fine particles induce cell apoptosis [2,9], the molecular pathways have not been well investigated. Our results suggest that, after 24-h treatment, both WC-Co nano- and fine particles induce JB6 apoptosis in a dose response manner within a certain dose range. Further studies revealed that treatment with WC-Co nanoparticles by intranasal droplet application induced both apoptosis and necrosis (or late apoptosis) in rat lung macrophages. Tozawa *et al* reported with a single intratracheal administration, WC-Co dusts induced marked fibrotic changes in rat pulmonary tissues after 6 months [37]. Combined with our results, apoptosis or necrosis in rat lung macrophages induced by WC-Co nanoparticles may play an important role in the formation of the experimental pneumoconiosis [37].

In mammals, signaling cascades culminating in apoptotic cell death can be divided into intrinsic or extrinsic pathways. The extrinsic pathway is initiated upon activation of death receptors. The intrinsic pathway can be initiated by cellular stresses, such as cytochrome *c* release from mitochondria into the cytoplasm. In this study, WC-Co particles induced Fas, FADD upregulation and caspase 8 activation. These results imply that in the apoptotic process induced by WC-Co particles, the extrinsic signal pathway is initiated. To investigate the involvement of intrinsic pathways in the apoptotic process induced by WC-Co particles, AIF and cytochrome *c* release from mitochondria to the cytoplasm were examined. Our results show that WC-Co particles induced upregulation and release of AIF and cytochrome *c* from mitochondria to the cytoplasm. In addition, the mitochondrial permeability assay showed that WC-Co particles induced significant changes in the mitochondrial membrane permeability, which was in parallel with western blot results. These results indicate that the apoptotic process induced by WC-Co particles involves both intrinsic and extrinsic pathways.

Lamins are nuclear membrane structural components, which are important for maintaining normal cell functions. Proteolysis of lamins has been observed in different cells undergoing apoptosis [38]. Degradation of lamina proteins can be triggered by both the extrinsic and the intrinsic pathways [39]. Our results show that lamin A, but not C, was cleaved in cells treated with WC-Co nanoparticles, suggesting involvement of lamin A in the apoptotic process induced by WC-Co nanoparticles. The actin cytoskeleton is capable of responding to complex signaling cascades. The major cytoskeletal filament, actin, can be degraded during the execution phase of apoptosis [40]. Reports indicate that disruption of actin filament integrity promptly induces apoptosis in adherent epithelial cells [41]. In addition, the dynamic state of actin is important in the regulation of ion channels [42]. In the present study, WC-Co particles induced β-actin cleavage after 1-h treatment in JB6 cells. These results, combined with our previous study with metallic nickel particles [7] and a report by White *et al* [41], suggest that the actin cytoskeletal network may act as a target for apoptosis and an early signaling component toward apoptotic commitment in the apoptotic process induced by metallic particles.

The Bcl-2 proteins are a family of proteins involved in the response to apoptosis. Some of these proteins, such as BAX and BID, are proapoptotic, while others, such as Bcl-2, are anti-apoptotic. In the present study, WC-Co particles induced both proapoptotic

conditions for the selection of cells that have become resistant to apoptosis, which may also be important in the WC-Co-induced carcinogenic process. Therefore, further research is needed to elucidate the role of activation of Bcl-2 in the carcinogenicity of WC-Co particles.

Mitochondrial dysfunction has been shown to participate in the induction of apoptosis. Changes in the mitochondrial membrane permeability are considered early events in apoptosis [43]. Many proapoptotic proteins, including cytochrome *c*, AIF, heat shock proteins, Smac/Diablo, and endonuclease G, can be released from the mitochondria into the cytoplasm following the formation of a pore in the mitochondrial membrane [44]. AIF is involved in initiating a caspase-independent pathway of apoptosis by causing DNA fragmentation and chromatin condensation [45]. However, cytochrome *c* induces apoptosis in a caspase-dependent pathway [46]. In this study, WC-Co particles, especially nanoparticles, induced a significant increase in the mitochondrial membrane permeability in JB6 cells after 24-h treatment and release of cytochrome *c* and AIF from mitochondria into cytoplasm. In addition, lung macrophages from rats exposed to WC-Co nanoparticles by intranasal droplet application (4 g/rat) for 21 days exhibited significant mitochondrial membrane permeability alteration in rat lung macrophages. These results indicate that the intrinsic mitochondrial pathway is involved in WC-Co particle-induced apoptosis.

Caspases, a family of aspartic acid-specific proteases, are the major effectors of apoptosis. The execution of apoptosis comprises both caspase-dependent and caspase-independent processes. In the present study, WC-Co particles significantly activated caspase 3 and 8. Caspase 9 was slightly activated. Taken together, our results suggest that apoptosis induced by WC-Co particles in JB6 cells involves both caspase-dependent and caspase-independent pathways.

In conclusion, the major findings of the present study show that both WC-Co nano- and fine particles induced JB6 cell and rat lung macrophage apoptosis. WC-Co nanoparticles elicited higher cytotoxicity and apoptotic induction than fine particles. WC-Co particles stimulated ROS generation in JB6 cells and rat lung macrophages. Catalase exhibited partly inhibitory effects on WC-Co particle-induced ROS generation and mitochondrial membrane permeability damage. To our knowledge, this is the first study showing that WC-Co particles activated both extrinsic and intrinsic apoptotic pathways in JB6 cells, which include upregulation of Fas, FADD, caspase 3, 8 and 9, BAX and BID, as well as release of AIF and cytochrome *c* from mitochondria into the cytoplasm. Lamin A and β -actin were cleaved. Furthermore, an increase of mitochondrial membrane permeability was observed in both WC-Co particle-treated JB6 cells and rat lung macrophages. On a mass basis, WC-Co nanoparticles exhibited higher cytotoxicity and apoptotic induction than fine particles. Particle size may play a critical role in the toxicity of this hard metal. Specific surface area tends to overcorrect for the greater toxicity when comparing the cytotoxicity of WC-Co nano- with fine particles. Apoptosis induced by WC-Co nano- and fine particles in JB6 cells may be mediated through ROS generation.

Acknowledgements

This study was partly supported by the National Nature Science Foundation of China (Grant no. 81273111), the Foundations of Innovative Research Team of Educational Commission of Zhejiang Province (T200907), the Nature Science Foundation of Ningbo city (Grant no. 2012A610185), the Ningbo Scientific Project (SZX11073), the Scientific Innovation Team Project of Ningbo (no. 2011B82014), Innovative Research Team of Ningbo (2009B21002) and K.C. Wong Magna Fund in Ningbo University.

References

- 1 Dagani R: Nanostructured materials promise to advance range of technologies. Chem Engineer News. 23:18–24. 1992. [View Article](#) : [Google Scholar](#)

- Lombaert N, Lison D, Van Hummelen P and Kirsch-Volders M: In vitro expression of hard metal dust (WC-Co) – responsive genes in human peripheral blood mononucleated cells. *Toxicol Appl Pharmacol.* 227:299–312. 2008. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
- 3 Driscoll KE, Maurer JK, Lindenschmidt RC, Romberger D, Rennard SI and Crosby L: Respiratory tract responses to dust: relationships between dust burden, lung injury, alveolar macrophage fibronectin release, and the development of pulmonary fibrosis. *Toxicol Appl Pharmacol.* 106:88–101. 1990. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 4 Oberdorster G, Ferin J and Lehnert BE: Correlation between particle size, in vivo particle persistence, and lung injury. *Environ Health Perspect.* 102(Suppl 5): 173–179. 1994. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 5 Oberdorster G: Pulmonary effects of inhaled ultrafine particles. *Int Arch Occup Environ Health.* 74:1–8. 2001. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 6 Zhao J, Bowman L, Zhang X, et al: Titanium dioxide (TiO₂) nanoparticles induce JB6 cell apoptosis through activation of the caspase-8/Bid and mitochondrial pathways. *J Toxicol Environ Health A.* 72:1141–1149. 2009. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 7 Zhao J, Bowman L, Zhang X, et al: Metallic nickel nano- and fine particles induce JB6 cell apoptosis through a caspase-8/AIF mediated cytochrome c-independent pathway. *J Nanobiotechnol.* 7:22009. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 8 Pulido MD and Parrish AR: Metal-induced apoptosis: mechanisms. *Mutat Res.* 533:227–241. 2003. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 9 Lombaert N, De Boeck M, Decordier I, Cundari E, Lison D and Kirsch-Volders M: Evaluation of the apoptogenic potential of hard metal dust (WC-Co), tungsten carbide and metallic cobalt. *Toxicol Lett.* 154:23–34. 2004. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 10 Raff MC: Social controls on cell survival and cell death. *Nature.* 356:397–400. 1992. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 11 Steller H: Mechanisms and genes of cellular suicide. *Science.* 267:1445–1449. 1995. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 12 White E: Life, death, and the pursuit of apoptosis. *Genes Dev.* 10:1–15. 1996. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 13 Wyllie AH, Kerr JF and Currie AR: Cell death: the significance of apoptosis. *Int Rev Cytol.* 68:251–306. 1980. [View Article](#) : [Google Scholar](#)
 - 14 Lu X, Lee M, Tran T and Block T: High level expression of apoptosis inhibitor in hepatoma cell line expressing Hepatitis B virus. *Int J Med Sci.* 2:30–35. 2005. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 15 Ramp U, Krieg T, Caliskan E, et al: XIAP expression is an independent prognostic marker in clear-cell renal carcinomas. *Hum Pathol.* 35:1022–1028. 2004. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 16 Kuwano K, Miyazaki H, Hagimoto N, et al: The involvement of Fas-Fas ligand pathway in fibrosing lung diseases. *Am J Respir Cell Mol Biol.* 20:53–60. 1999. [View](#)

- Bardales RH, Xie SS, Schaefer RF and Hsu SM: Apoptosis is a major pathway responsible for the resolution of type II pneumocytes in acute lung injury. *Am J Pathol.* 149:845–852. 1996.[PubMed/NCBI](#)
- 18 Levine AJ: p53, the cellular gatekeeper for growth and division. *Cell.* 88:323–331. 1997. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 19 Matute-Bello G, Winn RK, Jonas M, Chi EY, Martin TR and Liles WC: Fas (CD95) induces alveolar epithelial cell apoptosis in vivo: implications for acute pulmonary inflammation. *Am J Pathol.* 158:153–161. 2001. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 20 McCabe MJ Jr: Mechanisms and consequences of silica-induced apoptosis. *Toxicol Sci.* 76:1–2. 2003. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 21 Gawlitta D, Oomens CW, Baaijens FP and Bouten CV: Evaluation of a continuous quantification method of apoptosis and necrosis in tissue cultures. *Cytotechnology.* 46:139–150. 2004. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 22 Boffa DJ, Waka J, Thomas D, et al: Measurement of apoptosis of intact human islets by confocal optical sectioning and stereologic analysis of YO-PRO-1-stained islets. *Transplantation.* 79:842–845. 2005. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 23 Ding M, Kisin ER, Zhao J, et al: Size-dependent effects of tungsten carbide-cobalt particles on oxygen radical production and activation of cell signaling pathways in murine epidermal cells. *Toxicol Appl Pharmacol.* 241:260–268. 2009. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 24 Grossmann J, Mohr S, Lapentina EG, Fiocchi C and Levine AD: Sequential and rapid activation of select caspases during apoptosis of normal intestinal epithelial cells. *Am J Physiol.* 274:G1117–G1124. 1998.[PubMed/NCBI](#)
 - 25 Okinaga T, Kasai H, Tsujisawa T and Nishihara T: Role of caspases in cleavage of lamin A/C and PARP during apoptosis in macrophages infected with a periodontopathic bacterium. *J Med Microbiol.* 56:1399–1404. 2007. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 26 Mashima T, Naito M and Tsuruo T: Caspase-mediated cleavage of cytoskeletal actin plays a positive role in the process of morphological apoptosis. *Oncogene.* 18:2423–2430. 1999. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 27 Caroppi P, Sinibaldi F, Fiorucci L and Santucci R: Apoptosis and human diseases: mitochondrion damage and lethal role of released cytochrome C as proapoptotic protein. *Curr Med Chem.* 16:4058–4065. 2009. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 28 Susin SA, Lorenzo HK, Zamzami N, et al: Molecular characterization of mitochondrial apoptosis-inducing factor. *Nature.* 397:441–446. 1999. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 29 Susin SA, Dugas E, Ravagnan L, et al: Two distinct pathways leading to nuclear apoptosis. *J Exp Med.* 192:571–580. 2000. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 30 Piacenza L, Irigoin F, Alvarez MN, et al: Mitochondrial super-oxide radicals mediate programmed cell death in *Trypanosoma cruzi*: cytoprotective action of mitochondrial iron superoxide dismutase overexpression. *Biochem J.* 403:323–334. 2007.

- Long TC, Tajuba J, Sama P, et al: Nanosize titanium dioxide stimulates reactive oxygen species in brain microglia and damages neurons in vitro. *Environ Health Perspect.* 115:1631–1637. 2007. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
- 32 Lison D: Human toxicity of cobalt-containing dust and experimental studies on the mechanism of interstitial lung disease (hard metal disease). *Crit Rev Toxicol.* 26:585–616. 1996. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 33 Kear BH, Skandan G and Sadangi RK: Factors controlling decarburization in HV OF sprayed nano-WC/Co hardcoatings. *Scripta Materialia.* 44:1703–1707. 2001. [View Article](#) : [Google Scholar](#)
 - 34 Busch W, Kuhnel D, Schirmer K and Scholz S: Tungsten carbide cobalt nanoparticles exert hypoxia-like effects on the gene expression level in human keratinocytes. *BMC Genomics.* 11:652010. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 35 IARC Working Group on the Evaluation of Carcinogenic Risks to Humans: Cobalt in hard metals and cobalt sulfate, gallium arsenide, indium phosphide and vanadium pentoxide. *IARC Monogr Eval Carcinog Risks Hum.* 86:1–294. 2006. [PubMed/NCBI](#)
 - 36 Helewski KJ, Kowalczyk-Ziomek GI and Konecki J: Apoptosis and necrosis – two different ways leading to the same target. *Wiad Lek.* 59:679–684. 2006. (In Polish).
 - 37 Tozawa T, Kitamura H, Koshi K, Ikemi Y and Ambe K: Experimental pneumoconiosis induced by cemented tungsten and sequential concentrations of cobalt and tungsten in the lungs of the rat. *Sangyo Igaku.* 23:216–226. 1981. (In Japanese).
 - 38 Goldberg M, Harel A and Gruenbaum Y: The nuclear lamina: molecular organization and interaction with chromatin. *Crit Rev Eukaryot Gene Expr.* 9:285–293. 1999. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 39 Broers JL, Bronnenberg NM, Kuijpers HJ, Schutte B, Hutchison CJ and Ramaekers FC: Partial cleavage of A-type lamins concurs with their total disintegration from the nuclear lamina during apoptosis. *Eur J Cell Biol.* 81:677–691. 2002. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 40 Janmey PA: The cytoskeleton and cell signaling: component localization and mechanical coupling. *Physiol Rev.* 78:763–781. 1998. [PubMed/NCBI](#)
 - 41 White SR, Williams P, Wojcik KR, et al: Initiation of apoptosis by actin cytoskeletal derangement in human airway epithelial cells. *Am J Respir Cell Mol Biol.* 24:282–294. 2001. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 42 Gourlay CW and Ayscough KR: The actin cytoskeleton: a key regulator of apoptosis and ageing? *Nat Rev Mol Cell Biol.* 6:583–589. 2005. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 43 Ly JD, Grubb DR and Lawen A: The mitochondrial membrane potential ($\Delta\psi$) in apoptosis; an update. *Apoptosis.* 8:115–128. 2003. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 44 Siskind LJ, Kolesnick RN and Colombini M: Ceramide forms channels in mitochondrial outer membranes at physiologically relevant concentrations. *Mitochondrion.* 6:118–125. 2006. [View Article](#) : [Google Scholar](#) : [PubMed/NCBI](#)
 - 45 Landshamer S, Hoehn M, Barth N, et al: Bid-induced release of AIF from mitochondria.

- 46 Garland JM and Rudin C: Cytochrome c induces caspase-dependent apoptosis in intact hematopoietic cells and overrides apoptosis suppression mediated by bcl-2, growth factor signaling, MAP-kinase-kinase, and malignant change. *Blood*. 92:1235–1246. 1998.

Related Articles

- Induction of apoptosis in MDA-MB-231 human breast carcinoma cells with an ethanol extract of *Cyperus rotundus* L. by activating caspases
- 3-Hydroxyterphenyllin, a natural fungal metabolite, induces apoptosis and S phase arrest in human ovarian carcinoma cells
- The newly synthesized 2-[3-hydroxy-5-methoxyphenyl]-6,7-methylenedioxyquinolin-4-one triggers cell apoptosis through induction of oxidative stress and upregulation of the p38 MAPK signaling pathway in HL-60 human leukemia cells
- Pregnenolone, a cholesterol metabolite, induces glioma cell apoptosis via activating extrinsic and intrinsic apoptotic pathways
- Mechanism of N-acetyl-cysteine inhibition on the cytotoxicity induced by titanium dioxide nanoparticles in JB6 cells transfected with activator protein-1



DEMETRIOS A. SPANDIDOS

THE CANCER STORY

THE DISCOVERY OF
CELLULAR ONCOGENES

There is no doubt that the discovery of cellular oncogenes changed our views on the most devastating of human diseases: cancer. That is the reason why it is considered the most significant discovery of the 20th century. From the time of Hippocrates, the father of medicine who first described cancer, until the discovery of cellular oncogenes, very little progress had been made concerning an explanation to the causes of cancer, its diagnosis and treatment. The fact that under the category of cancer approximately 200 forms of diseases are included affecting almost every part of the human body, made it difficult to pinpoint the cause as well as control its result. However, the development of a variety of technological achievements, namely tissue culture, recombinant DNA technology, gene transfer techniques and so on, made the ideological approach feasible. By using the above techniques, it was possible to isolate human oncogenes; in other words, the genes responsible for the creation of benign or malignant growth of human tumours (oncos).

Download pdf



26th International Symposium on Molecular Medicine

Dates to be announced

GRAND HOTEL DUCA D'ESTE
Tivoli, Rome, ITALY

MYSTRAS
GRAND PALACE
RESORT & SPA

ALL LAWS AND EVERY
LAW'S TASK SHOULD AIM
TO EDUCATION

LYKOURGOS 800-730 BC

New Dates to be announced



28th International Symposium on Molecular Medicine

date to be announced

Guangzhou
CHINA



25th International Symposium on Molecular Medicine

new dates to be announced

PARK PLAZA HOTEL
SANTIAGO, CHILE



WORLD ACADEMY
OF SCIENCES

Copyright © Spandidos Publications 2022. All rights reserved.

[About](#) | [Contact](#) | [Help](#) | [Cookie Policy](#) | [Privacy Policy](#)



This site uses cookies

You can change your cookie settings at any time by following the instructions in our [Cookie Policy](#). To find out more, you may read our [Privacy Policy](#). [I agree](#)