

# Association Between Work–Family Conflict and Musculoskeletal Pain Among Hospital Patient Care Workers

**Seung-Sup Kim, ScD,<sup>1\*</sup> Cassandra A. Okechukwu, ScD,<sup>2</sup> Orfeu M. Buxton, PhD,<sup>3,4</sup>  
Jack T. Dennerlein, PhD,<sup>5</sup> Leslie I. Boden, PhD,<sup>6</sup> Dean M. Hashimoto, MD,<sup>3,4,5,7</sup>  
and Glorian Sorensen, PhD<sup>2,8</sup>**

**Background** A growing body of evidence suggests that work–family conflict is an important risk factor for workers’ health and well-being. The goal of this study is to examine association between work–family conflict and musculoskeletal pain among hospital patient care workers.

**Methods** We analyzed a cross-sectional survey of 1,119 hospital patient care workers in 105 units in two urban, academic hospitals. Work–family conflict was measured by 5-item Work–Family Conflict Scale questionnaire. Multilevel logistic regression was applied to examine associations between work–family conflict and self-reported musculoskeletal pain in the past 3 months, adjusting for covariates including work-related psychosocial factors and physical work factors.

**Results** In fully adjusted models, high work–family conflict was strongly associated with neck or shoulder pain (OR: 2.34, 95% CI: 1.64–3.34), arm pain (OR: 2.79, 95% CI: 1.64–4.75), lower extremity pain (OR: 2.20, 95% CI: 1.54–3.15) and any musculoskeletal pain (OR: 2.45, 95% CI: 1.56–3.85), and a number of body areas in pain (OR: 2.47, 95% CI: 1.82–3.36) in the past 3 months. The association with low back pain was attenuated and became non-significant after adjusting for covariates.

**Conclusions** Given the consistent associations between work–family conflict and self-reported musculoskeletal pains, the results suggest that work–family conflict could be an important domain for health promotion and workplace policy development among hospital patient care workers. *Am. J. Ind. Med.* 56:488–495, 2013.

© 2012 Wiley Periodicals, Inc.

**KEY WORDS:** *work–family conflict; musculoskeletal symptoms; low back pain; hospital worker*

<sup>1</sup>Department of Environmental and Occupational Health, The George Washington University School of Public Health and Health Services, Washington, District of Columbia

<sup>2</sup>Department of Society, Human Development, and Health, Harvard School of Public Health, Boston, Massachusetts

<sup>3</sup>Department of Medicine, Brigham and Women’s Hospital, Boston, Massachusetts

<sup>4</sup>Harvard Medical School, Boston, Massachusetts

<sup>5</sup>Department of Environmental Health, Harvard School of Public Health, Boston, Massachusetts

<sup>6</sup>Department of Environmental Health, Boston University School of Public Health, Boston, Massachusetts

<sup>7</sup>Occupational Health Services, Partners HealthCare System, Boston, Massachusetts

<sup>8</sup>Dana-Farber Cancer Institute, Center for Community-Based Research, Boston, Massachusetts

Contract grant sponsor: National Institute for Occupational Safety and Health; Contract grant number: U19 OH008861.

Contract grant sponsor: National Heart, Lung and Blood Institute; Contract grant number: R01HL107240.

Disclosure Statement: Dr. Hashimoto is employed by Partners HealthCare System, and the two studied hospitals are Partners hospitals. There are no other possible conflicts of interest to report.

\*Correspondence to: Dr. Seung-Sup Kim, ScD, Department of Environmental and Occupational Health, The George Washington University School of Public Health and Health Services, 2300 Eye Street, NW, Washington, DC 20037. E-mail: sskim@gwu.edu

Accepted 20 August 2012

DOI 10.1002/ajim.22120. Published online 27 September 2012 in Wiley Online Library (wileyonlinelibrary.com).

## INTRODUCTION

A growing body of evidence demonstrates that work-family conflict, defined as a conflict between work and family roles, is an important risk factor for workers' health and well-being [Allen and Armstrong, 2006; Amstad et al., 2011]. Previous studies have focused on the impacts of work-family conflicts on workers' well-being (e.g., job satisfaction [Cortese et al., 2010], intention to turnover [Fuss et al., 2008], and health satisfaction [Knecht et al., 2011]) and mental health outcomes (e.g., depression [Frone et al., 1996; Hammer et al., 2005; Okechukwu et al., 2012], substance dependence disorder [Frone, 2000], and mood/anxiety disorders [Wang et al., 2007]).

Recently, several studies have suggested that work-family conflict could be detrimental to workers' physical health outcomes as well [Jansen et al., 2006; Clays et al., 2009; Berkman et al., 2010; Ohta et al., 2011; Sabbath et al., 2012]. For example, in a prospective study of over 6,000 employees, Jansen et al. [2006] showed that workers with high work-family conflict were more likely to have long duration of sickness absence than workers with low-medium work-family conflict after adjusting for confounders including work schedule and having dependent children. Also, Berkman et al. [2010] found that working with managers who are supportive about work-family need is associated with reduced cardiovascular disease risk.

Work-family conflict is of increasing concern for workers who provide direct patient care in hospitals [Grzywacz et al., 2006]. For example, one U.S. survey of nurses showed that more than 90% experienced work-family conflict more than once over the past 6 months and fully half reported experiencing work-family conflict more than 1 day per week [Grzywacz et al., 2006]. Previous studies have also found that work-family conflict is associated with various negative outcomes such as job dissatisfaction [Cortese et al., 2010], lower sense of coherence [Takeuchi and Yamazaki, 2010], and increase of perceived work demand [Camerino et al., 2010] among patient care workers in hospitals.

However, little is known about how work-family conflict is related to physical health outcomes such as musculoskeletal disorders among hospital patient care workers despite their high prevalence of musculoskeletal disorders [Engels et al., 1996]. In two studies, Hammig et al. investigated the association between work-family conflict and musculoskeletal pain [Hammig and Bauer, 2009; Hammig et al., 2011]. They found an association between high work-family conflict and self-reported neck/shoulder pain and back pain among workers across four different industries including the healthcare sector [Hammig et al., 2011]. However, this study did not adjust for working conditions such as night shift work and number of work

hours. These factors are particularly critical for hospital health care workers in relation to both their experiences of work-family conflict and their potential exposure to musculoskeletal pain [Eriksen et al., 2004; Dex and Bond, 2005; Trinkoff et al., 2006; Camerino et al., 2010]. In addition, the focus on neck, shoulder and back pain may miss other types of musculoskeletal pain common among hospital patient care workers, who are known to experience a high rate of musculoskeletal pain in other body areas such as arm and low extremities [Ando et al., 2000; Smith et al., 2003].

In this study, after adjusting for covariates including six physical work factors and four work-related psychosocial factors, we examined whether work-family conflict is associated with self-reported musculoskeletal pain (i.e., neck or shoulder pain, arm pain, low back pain, leg pain, any musculoskeletal pain, and a number of body areas in pain) in the past 3 months among hospital patient care workers.

## MATERIALS AND METHODS

### Study Population

We conducted a cross-sectional survey of 2,000 randomly selected patient care workers in two large academic hospitals in the metropolitan Boston area between October 2009 and February 2010. This survey was part of the "Be Well Work Well" study, conducted by the Harvard School of Public Health Center for Work, Health and Wellbeing. Eligible participants included registered nurses, licensed practical nurses, and patient care/nursing assistants with direct patient care responsibilities. A detailed description of the study design, protocol and process of data collection is provided elsewhere [Sorensen et al., 2011; Dennerlein et al., 2012]. All project materials and methods were approved by the Institutional Review Board of Partners HealthCare.

Among 2,000 workers, 1,572 (79%) health care workers completed the survey. After excluding workers with missing values either for work-family conflict or for any covariate, we analyzed a total of 1,199 patient care workers in 105 units with 12 different types of units (i.e., emergency department, operating room, adult medical/surgical, adult intensive care, step-down, pediatric medical/surgical, pediatric/neonatal intensive care, psychiatry, obstetrics/postpartum, float pool, ambulatory units, and orthopedics).

### Measures

#### Work-family conflict

Work-family conflict was measured by the Work-Family Conflict Scale questionnaire, which was developed

by Netemeyer to address how work can interfere with family life [Netemeyer et al., 1996]. We modified wording of the questionnaire so that workers without family member in their household could also be eligible to answer the questions. The survey questions include: How much do you agree or disagree with the following statements: (1) The demands of my work interfere with my family or personal time. (2) The amount of time my job takes up makes it difficult to fulfill family or personal responsibilities. (3) Things I want to do at home do not get done because of the demands my job puts on me. (4) My job produces strain that makes it difficult to fulfill my family or personal duties. (5) Due to work-related duties, I have to make changes to my plans for family or personal activities. For each question, five ordinal answers were available (1: strongly disagree, 2: disagree, 3: neither disagree nor agree, 4: agree, 5: strongly agree). We classified the summed responses of workers into three categories (5–12: low, 13–17: intermediate, 18–25: high). Cronbach's alpha among five questions used to measure work–family conflict was 0.89.

### ***Self-reported musculoskeletal pains in the past 3 months***

Musculoskeletal pains were assessed for the past 3 months based on the Standardized Nordic Questionnaire for musculoskeletal symptoms [20–21]. Specifically, we asked, "During the past 3 months, have you had pain or aching in any of the areas shown on the diagram?" The areas were neck, shoulder, lower back, wrist or forearm, knee, ankle or feet, and none of the above. We combined responses for the neck and shoulder into a single neck/shoulder area and the responses for the knees, ankle and feet were grouped into a single lower extremity body area. We then created an ordinal variable of the number of body areas in pain (0–4) by summing the responses. We also created the variable "any musculoskeletal pain" to assess whether the worker had a pain in at least one body part with those who did not have any pain as a reference.

### ***Covariates***

We first included basic socio-demographic variables as covariates: age (18–24, 25–34, 35–44, 45–54,  $\geq 55$  years); race (Hispanic, White, Black, and mixed race/others), capability to pay bills (0: great deal of difficulty, 1: some difficulty, 2: a little difficulty, 3: no difficulty, 4: don't know, 5: refused), the number of children under 5 years old in home (0–5 children). In addition, we included job title (staff nurse, patient care associate, and others) because workers in the same job title are expected to have similar expected workloads that could be associated with both work–family conflict and musculoskeletal pain.

To find other relevant work-related covariates, we selected variables that are reported to be associated with work–family conflict and musculoskeletal pain based on previous reports [Ariens, 2001; Eriksen et al., 2004; Dex and Bond, 2005; Trinkoff et al., 2006; Grönlund, 2007; Pisarski et al., 2008; Camerino et al., 2010]. As a result, we included seven work-related variables including four psychosocial factors. We included worked hours per week five categories (<30, 30–34, 35–39, 40–44, 45–50, and >50 hr) [Dex and Bond, 2005; Trinkoff et al., 2006]. Since worked hours per week does not include working time in a second job, we included having a second job or not because it reflects extra working hours and potential risk burden. Shift work was dichotomized into having a night shift versus working only on daytime [Eriksen et al., 2004; Camerino et al., 2010]. Four work-related psychosocial factors were measured using a modified the Job Content Questionnaire [Karasek, 1979; Karasek et al., 1998; Landsbergis et al., 2002]: psychological demands (five items) and decision latitude (nine items) [Ariens, 2001; Grönlund, 2007], supervisor support (two items) and co-worker support (three items) [Eriksen et al., 2004; Pisarski et al., 2008].

To assess organizational factors related to ergonomic practices in a hospital unit, we included an organizational ergonomic policies and practices questionnaire [Amick et al., 2000] modified as described previously [Dennerlein et al., 2012]. The modified questionnaire is composed of six questions to assess workers' perceptions about organizational practices around patient lifting and equipment lifting; pushing pulling, bending, reaching and stooping; workstation design; and purchasing tools, equipment or furniture. Workers could answer on a five-point scale for each of six questions with higher score indicating better ergonomic practice. Summed responses ranged from 6 to 30.

To take physical work factors into account, we also adjusted for physical activity on the job and use of a lifting device. Physical activity on the job was assessed using five questions asking patient care workers to "estimate how much of a typical shift you spend: (a) sitting, (b) standing, (c) walking, (d) lifting and carrying, (e) pushing and pulling" [Reis et al., 2005]. Workers could answer in a five-point ordinal scale for each of five questions (all, more than half, about half, less than half, and none). The response was collapsed into three levels: all and more than half, about half, less than half and none. Use of a lifting device was measured using a question "In general, when a patient needs to be moved, how often do you use a lifting device?" Workers could answer in a five-point scale from "never" to "always" or indicate that they do not move patients on their job. The response was categorized into four groups: low (never and rarely), medium (sometimes), high (often and always), and not applicable.

## Data Analyses

Because patient care workers were nested within hospital unit, prevalence of musculoskeletal pain among workers in the same unit could be highly correlated with each other. To take those correlations and a hierarchical data structure into account, we applied multilevel logistic regression with a random intercept in a hospital unit to examine associations between work-family conflict and musculoskeletal pain. The number of body areas in pain was assessed as an ordinal scale; therefore, multilevel ordinal logistic regressions were applied to examine the association. All analyses were performed using STATA/SE version 11.0 (StataCorp, College Station, TX).

## RESULTS

Table I shows the distribution of the study population and the prevalence of any self-reported musculoskeletal pain in the past 3 months and the mean score of work-family conflict by key covariates. Among 1,199 hospital patient care workers, 89.9% were female, 73.9% were staff nurse, 28.4% worked some night shift. Female workers (77.0%) were more likely to report any musculoskeletal pain in the past 3 months compared to male workers (66.7%). Workers who had a second job or who worked night shift reported significantly higher levels of work-family conflict compared to their counterparts. For the prevalence of musculoskeletal pain in the past 3 months,

**TABLE I.** Distribution of Study Population and Prevalence of Any Musculoskeletal Pain in the Past 3 Months and Mean Score of Work-Family Conflict by Key Covariates (N = 1,199)

Characteristics	Distribution		P-value <sup>a</sup>	Work-family conflict (5–25)	
	N (%)	N (%)		Mean (SD)	P-value <sup>b</sup>
Gender			0.012		0.906
Male	120 (9.9)	80 (66.7)		13.6 (4.1)	
Female	1,079 (89.9)	831 (77.0)		13.6 (4.4)	
Age (years)			0.059		0.008
18–24	90 (7.5)	68 (76.0)		14.4 (4.1)	
25–34	339 (28.3)	265 (78.2)		14.0 (4.0)	
35–44	287 (24.0)	200 (69.7)		13.3 (4.4)	
45–54	319 (26.6)	253 (79.3)		13.4 (4.4)	
55	164 (13.7)	125 (76.2)		12.9 (4.7)	
Race			0.016		<0.001
Hispanic	45 (3.8)	35 (77.8)		12.1 (4.1)	
White, non-Hispanic	985 (82.2)	761 (77.3)		13.8 (4.3)	
Black, non-Hispanic	103 (8.6)	65 (63.1)		11.7 (4.0)	
Other	66 (5.5)	50 (75.8)		13.2 (4.2)	
Job title			<0.001		0.001
Staff nurse	886 (73.9)	700 (79.1)		13.7 (4.2)	
Patient care associate	70 (5.8)	50 (71.4)		12.0 (4.5)	
Others	243 (20.3)	161 (66.3)		13.2 (4.7)	
Hours worked per week			0.748		<0.001
<30	276 (23.0)	208 (75.4)		13.0 (3.9)	
30–34	144 (12.0)	111 (77.1)		13.8 (4.5)	
35–39	361 (30.1)	276 (76.5)		13.6 (4.0)	
40–44	364 (30.4)	279 (76.7)		13.5 (4.6)	
≥45	54 (4.5)	37 (68.5)		16.5 (5.2)	
Having a second job			0.366		<0.001
Yes	172 (14.4)	126 (73.6)		14.6 (4.4)	
No	1,027 (85.7)	785 (76.4)		13.4 (4.3)	
Having only day shift, %			0.643		0.001
Yes	341 (28.4)	256 (75.1)		12.9 (4.5)	
No (also night shift)	858 (71.6)	655 (76.3)		13.8 (4.3)	

<sup>a</sup>P-value of the Chi-square test comparing the prevalence of any musculoskeletal pain across the different groups.

<sup>b</sup>P-value of the Analysis of Variance test comparing the mean scores of work-family conflict across the different groups.

**TABLE II.** Distribution of Self-Reported Musculoskeletal Pain in the Past 3 Months Among Hospital Patient Care Workers (N = 1,119)

Self-reported musculoskeletal pain	Distribution	
	N	%
Neck/shoulder pain	537	44.8
Arm pain	126	10.5
Low back pain	662	55.2
Lower extremity pain	454	37.8
Any musculoskeletal pain	911	76.0
Number of body areas in pain		
0	288	24.0
1	323	27.0
2	353	29.4
3	190	15.9
4	45	3.8

the highest prevalence was for back pain (55.2%) while the lowest prevalence was for arm pain (10.5%; Table II). Overall, 76.0% of hospital patient care workers reported pain in at least one body part and 3.8% reported pain in all of four body parts (i.e., neck/shoulder, arm, low back, and lower extremity pain).

As shown in Table III, after adjusting for covariates work–family conflict remained statistically significantly associated with all indicators of musculoskeletal pain in the past 3 months except low-back pain. We found the strong associations of high work–family conflict with neck or shoulder pain (OR: 2.34, 95% CI: 1.64–3.34), arm pain (OR: 2.79, 95% CI: 1.64–4.75), lower extremity pain (OR: 2.20, 95% CI: 1.54–3.15) and any musculoskeletal pain (OR: 2.45, 95% CI: 1.56–3.85), and a number of body areas in pain (OR: 2.47, 95% CI: 1.82–3.36) in the past 3 months in the fully adjusted model. However, the association between high work–family conflict and low back pain was attenuated and became non-significant after adjustment for covariates (OR: 1.22, 95% CI: 0.86–1.74); whereas, the association with intermediate work–family conflict remained significant (OR: 1.49, 95% CI: 1.12–1.98).

## DISCUSSION

We found consistent associations between work–family conflict and self-reported musculoskeletal pain (i.e., neck or shoulder pain, arm pain, leg pain and any musculoskeletal pain, and a number of body areas in pain) in the past 3 months among hospital patient care workers. This analysis adjusted for relevant covariates, including physical work factors and work-related psychosocial factors. These results extend the findings of previous research reporting that work–family conflict is associated with

neck/shoulder pain and low back pain [Hammig et al., 2011], and are consistent with prior findings that work–family conflict is associated with a variety of negative consequences including depression [Hammig and Bauer, 2009], substance dependence disorder [Frone, 2000], cardiovascular disease risk [Berkman et al., 2010] and sickness absence [Clays et al., 2009; Sabbath et al., 2012].

Although, we found that high work–family conflict was associated with most indicators of musculoskeletal pain, the association between high levels of work–family conflict and self-reported low back pain was attenuated and became non-significant in the adjusted model, whereas the association between intermediate levels of work–family conflict and low back pain remained significant. This finding is different from findings from previous studies that reported an association between work–family conflict and low back pain, and may reflect previous studies' lack of full adjustment for working conditions such as shift work, worked hours per week, job title, physical work factors [Hammig et al., 2011] or other relevant psychosocial factors such as job demand, decision latitude, and supervisor support [Hammig et al., 2009]. In addition, future research needs to examine why the relationship of work–family conflict with low-back pain could be potentially different from the relationships with other musculoskeletal pains.

There could be several explanations of pathways linking work–family conflict and musculoskeletal pain. When workers experience work–family conflict, workers could be preoccupied at the workplace with home-related duties. Accordingly, it could influence psychosocial working conditions, such as job demand, which could then impact musculoskeletal pain [Bongers et al., 2002; Lang et al., 2012]. However, the adjustment of covariates including work-related psychosocial stressors (i.e., job demand, job control, supervisor support, and co-worker support), work physical factors (i.e., sitting, standing, walking, lifting and carrying, pushing, and pulling), and work characteristics (i.e., job title, having a second job or not, only day shift or not, worked hours per week) attenuated less than 20% of the crude associations for most of musculoskeletal outcomes in this study, this suggests that the association would not be wholly explained by psychosocial and other working conditions.

It is also possible that psychological stress play a role in the pathway because experiences of work–family conflict could increase physiological stress reaction such as increased muscle tension which can lead to musculoskeletal pain [Lundberg et al., 2002]. Hammig et al. [2011] examined a hypothesis that the association between work–family conflict and musculoskeletal pain is mediated by general stress, showing that the association was substantially attenuated when they adjusted for general stress in addition to other covariates. However, in a post-hoc

**TABLE III.** Association Between Work-Family Conflict and Musculoskeletal Pain in the Past 3 Months Among Hospital Patient Care Workers (N = 1,199)

Work-family conflict (5–25)	Neck/Shoulder pain				Arm pain			
	Unadjusted		Fully adjusted <sup>a</sup>		Unadjusted		Fully adjusted <sup>a</sup>	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
Low (5–12)	1	Referent	1	Referent	1	Referent	1	Referent
Moderate (13–17)	1.63**	1.25, 2.14	1.50**	1.13, 2.01	1.53	0.97, 2.42	1.55	0.95, 2.53
High (18–25)	2.58**	1.88, 3.55	2.34**	1.64, 3.34	2.88**	1.82, 4.56	2.79**	1.64, 4.75
Low back pain								
Work-family conflict (5–25)	Unadjusted		Fully adjusted <sup>a</sup>		Unadjusted		Fully adjusted <sup>a</sup>	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
	Low (5–12)	1	Referent	1	Referent	1	Referent	1
Moderate (13–17)	1.84**	1.40, 2.40	1.49**	1.12, 1.98	1.80**	1.27, 2.30	1.71**	1.32, 2.30
High (18–25)	1.63*	1.19, 2.22	1.22	0.86, 1.74	2.32**	1.69, 3.17	2.20**	1.54, 3.15
Any musculoskeletal pain								
Work-family conflict (5–25)	Unadjusted		Fully adjusted <sup>a</sup>		Unadjusted		Fully adjusted <sup>a</sup>	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
	Low (5–12)	1	Referent	1	Referent	1	Referent	1
Moderate (13–17)	2.22**	1.61, 3.05	1.89**	1.34, 2.67	2.01**	1.61, 2.58	1.78**	1.39, 2.28
High (18–25)	2.87**	1.92, 4.30	2.45**	1.56, 3.85	2.86**	2.16, 3.79	2.47**	1.82, 3.36

\*P < 0.01.

\*\*P < 0.001.

<sup>a</sup>Fully adjusted: Adjusted for age, race, gender, job title, having a second job or not, only day shift or not, worked hours per week, job demands, job control, supervisor support, co-worker support, use of a lifting device, physical activities on the job (i.e., sitting, standing, walking, lifting and carrying, pushing, and pulling), organizational ergonomic practice, and the number of children under 5 years old.

analysis when we adjusted for non-specific psychological distress assessed by the six-item Kessler scale [Kessler et al., 2002] in addition to other covariates, we found only slight attenuation in the associations between work-family conflict and musculoskeletal pain. Because there could be remained exposure to stress which could not be captured using a self-reported questionnaire, future research is required to examine the role of stress in the association between work-family conflict and musculoskeletal pain as a mediator.

This study also found that workers who worked more than 45 hr per week or who has a second job or who worked at night shift were more likely to experience work-family conflict than their counterparts, which is consistent with the findings of previous literature [Dex and Bond, 2005; Camerino et al., 2010]. We also found that staff nurse reported higher level of work-family conflict than others. The difference of work-family conflict level by job title may be explained by our finding that the prevalence of “having only a day shift” is much lower among staff nurse (22.6%) compared to patient care associate

(34.2%) and others (49.2%). Future studies with larger samples of different types of workers are needed to understand joint differences in work-family conflict and health-related associations by job title and other work characteristics. Future studies are also needed to investigate the impact of hospital-level policies, such as access to paid leave, on the association between work-family conflict and injury outcomes [Asfaw et al., 2012].

Several limitations to this study should be noted. First, temporal precedence cannot be established due to the cross sectional design of the study. We cannot rule out the possibility of reverse causation in which workers with musculoskeletal pain reported more work-family conflict. Second, although we had a strong adjustment of potential covariates related to working conditions, there could be residual confounding pertaining to worker’s health condition because workers with poor health conditions may be more likely to report musculoskeletal pain and also more likely to report high work-family conflict. This is particularly possible because both of exposure and outcome variables were assessed using a self-reported questionnaire.

Another limitation would be that, because of the high prevalence of several outcome variables such as low back pain and any musculoskeletal pain, the estimated odds ratio could overestimate the ratio that would have been observed had adjusted prevalence ratios been computed. However, we found strong relationship of work–family conflict with relatively low prevalent outcomes like arm pain as well as other highly prevalent outcomes, the observed strong association could not be wholly explained by the potential overestimation. Finally, work–life conflict consists of two reciprocal components: work–family conflict that work interferes with family life and family–work conflict that family interferes with work, this study only assessed work–family conflict. A previous study, which measured both of these components and examined their associations with musculoskeletal pain, found that family–work conflict is not associated with musculoskeletal pain whereas work–family conflict is a strong predictor of musculoskeletal pain [Hammig et al., 2011], as we have observed.

There are also several strengths in this study. First, we could adjust for covariates including working conditions, other work-related psychosocial constructs, organizational practice, physical work factors, and the number of children under 5 years old at home. Second, we applied multilevel logistic regression with a random intercept at the work unit level to take into account the hierarchical data structure.

Our study found that work–family conflict could be a potential risk factor of musculoskeletal pain among hospital health care workers. These findings remained significant after adjustments for working conditions (i.e., job title, having a second job or not, working only day shift or not, hours worked per week), psycho-social factors (i.e., demand, decision latitude, co-worker support, and supervisor support), organizational ergonomic practice, and physical work factors. Work–family conflict could be proposed as an important domain for health promotion strategy among hospital patient care workers. However, it will be important for future studies to first replicate these findings with a prospective study design.

## ACKNOWLEDGMENTS

This work was supported by a grant from the National Institute for Occupational Safety and Health (U19 OH008861) for the Harvard School of Public Health Center for Work, Health and Well-being. This study would not have been accomplished without the participation of Partners HealthCare System and leadership from Dennis Colling, Sree Chaguturu, and Kurt Westerman. The authors would like to thank Partners Occupational Health Services including Marlene Freeley for her guidance, as well as Elizabeth Taylor, Elizabeth Tucker O'Day, and

Terry Orechia. We also thank individuals at each of the hospitals including Jeanette Ives Erickson and Jacqueline Somerville in Patient Care Services leadership, and Jeff Davis and Julie Celano in Human Resources. Additionally, we wish to thank Charlene Feilteau, Mimi O'Connor, Margaret Shaw, Eddie Tan and Shari Weingarten for assistance with supporting databases. We also thank Chris Kenwood of NERI for his statistical and programming support, Evan McEwing, Project Director, and Linnea Benson-Whelan for her assistance with the production of this manuscript. Our thanks also go to Anne M. Stoddard for her comments about data analyses.

## REFERENCES

Allen TD, Armstrong J. 2006. Further examination of the link between work–family conflict and physical health. *Am Behav Sci* 49:1204–1221.

Amick BC, Habeck RV, Hunt A, Fossel AH, Chapin A, Keller RB, Katz JN. 2000. Measuring the impact of organizational behaviors on work disability prevention and management. *J Occup Rehabil* 10: 21–38.

Amstad FT, Meier LL, Fasel U, Elfering A, Semmer NK. 2011. A meta-analysis of work–family conflict and various outcomes with a special emphasis on cross-domain versus matching-domain relations. *J Occup Health Psychol* 16:151–169.

Ando S, Ono Y, Shimaoka M, Hiruta S, Hattori Y, Hori F, Takeuchi Y. 2000. Associations of self estimated workloads with musculoskeletal symptoms among hospital nurses. *Occup Environ Med* 57:211–216.

Ariens G. 2001. Psychosocial risk factors for neck pain: A systematic review. *Am J Ind Med* 39:180–193.

Asfaw A, Pana-Cryan R, Rosa R. 2012. Paid sick leave and nonfatal occupational injuries. *Am J Public Health* 102:e59–e64.

Berkman LF, Buxton O, Ertel K, Okechukwu C. 2010. Managers' practices related to work–family balance predict employee cardiovascular risk and sleep duration in extended care settings. *J Occup Health Psychol* 15:316–329.

Bongers PM, Kremer AM, Laak Jt. 2002. Are psychosocial factors, risk factors for symptoms and signs of the shoulder, elbow, or hand/wrist?: A review of the epidemiological literature. *Am J Ind Med* 41:315–342.

Camerino D, Sandri M, Sartori S, Conway PM, Campanini P, Costa G. 2010. Shiftwork, work–family conflict among Italian nurses, and prevention efficacy. *Chronobiol Int* 27:1105–1123.

Clays E, Kittel F, Godin I, Bacquer DD, Backer GD. 2009. Measures of work–family conflict predict sickness absence from work. *J Occup Environ Med* 51:879–886.

Cortese CG, Colombo L, Ghislieri C. 2010. Determinants of nurses' job satisfaction: The role of work–family conflict, job demand, emotional charge and social support. *J Nurs Manag* 18:35–43.

Dennerlein JT, Hopcia K, Sembajwe G, Kenwood C, Stoddard AM, Tveito TH, Hashimoto DM, Sorensen G. 2012. Ergonomic practices within patient care units are associated with musculoskeletal pain and limitations. *Am J Ind Med* 55:107–116.

Dex S, Bond S. 2005. Measuring work–life balance and its covariates. *Work Employ Soc* 19:627–637.

Engels JA, van der Gulden JW, Senden TF, van't Hof B. 1996. Work related risk factors for musculoskeletal complaints in the nursing profession: Results of a questionnaire survey. *Occup Environ Med* 53:636–641.

Eriksen W, Bruusgaard D, Knardahl S. 2004. Work factors as predictors of intense or disabling low back pain: A prospective study of nurses' aides. *Occup Environ Med* 61:398–404.

Frone MR. 2000. Work-family conflict and employee psychiatric disorders: The National Comorbidity Survey. *J Appl Psychol* 85: 888–895.

Frone MR, Russell M, Barnes GM. 1996. Work-family conflict, gender, and health-related outcomes: A study of employed parents in two community samples. *J Occup Health Psychol* 1:57–69.

Fuss I, Nubling M, Hasselhorn HM, Schwappach D, Rieger MA. 2008. Working conditions and work-family conflict in German hospital physicians: Psychosocial and organisational predictors and consequences. *BMC Public Health* 8:353.

Grönlund A. 2007. More control, less conflict? Job demand-control, gender and work-family conflict. *Gender Work Organ* 14:476–497.

Grzywacz JG, Frone MR, Brewer CS, Kovner CT. 2006. Quantifying work-family conflict among registered nurses. *Res Nurs Health* 29:414–426.

Hammer LB, Cullen JC, Neal MB, Sinclair RR, Shafiro MV. 2005. The longitudinal effects of work-family conflict and positive spillover on depressive symptoms among dual-earner couples. *J Occup Health Psychol* 10:138–154.

Hammig O, Bauer G. 2009. Work-life imbalance and mental health among male and female employees in Switzerland. *Int J Public Health* 54:88–95.

Hammig O, Gutzwiller F, Bauer G. 2009. Work-life conflict and associations with work- and nonwork-related factors and with physical and mental health outcomes: A nationally representative cross-sectional study in Switzerland. *BMC Public Health* 9:435.

Hammig O, Knecht M, Laubli T, Bauer G. 2011. Work-life conflict and musculoskeletal disorders: A cross-sectional study of an unexplored association. *BMC Musculoskeletal Disord* 12:60.

Jansen NW, Kant IJ, van Amelsvoort LG, Kristensen TS, Swaen GM, Nijhuis FJ. 2006. Work-family conflict as a risk factor for sickness absence. *Occup Environ Med* 63:488–494.

Karasek RA, Jr. 1979. Job demands, job decision latitude, and mental strain: Implications for job redesign. *Adm Sci Q* 24:285–308.

Karasek R, Brisson C, Kawakami N, Houtman I, Bongers P, Amick B. 1998. The Job Content Questionnaire (JCQ): An instrument for internationally comparative assessments of psychosocial job characteristics. *J Occup Health Psychol* 3:322–355.

Kessler RC, Andrews G, Colpe LJ, Hiripi E, Mroczek DK, Normand SL, Walters EE, Zaslavsky AM. 2002. Short screening scales to monitor population prevalences and trends in non-specific psychological distress. *Psychol Med* 32:959–976.

Knecht MK, Bauer GF, Gutzwiller F, Hammig O. 2011. Persistent work-life conflict and health satisfaction—A representative longitudinal study in Switzerland. *BMC Public Health* 11:271.

Landsbergis PA, Schnall PL, Pickering TG, Schwartz JE. 2002. Validity and reliability of a work history questionnaire derived from the Job Content Questionnaire. *J Occup Environ Med* 44:1037–1047.

Lang J, Ochsmann E, Kraus T, Lang JWB. 2012. Psychosocial work stressors as antecedents of musculoskeletal problems: A systematic review and meta-analysis of stability-adjusted longitudinal studies. *Soc Sci Med* 75:1163–1174.

Lundberg U, Forsman M, Zachau G, Eklöf M, Palmerud G, Melin B, Kadefors R. 2002. Effects of experimentally induced mental and physical stress on motor unit recruitment in the trapezius muscle. *Work Stress* 16:166–178.

Netemeyer RG, Boles JS, McMurrain R. 1996. Development and validation of work-family conflict and family-work conflict scales. *J Appl Psychol* 81:400–410.

Ohta H, Wada K, Kawashima M, Arimatsu M, Higashi T, Yoshikawa T, Aizawa Y. 2011. Work-family conflict and prolonged fatigue among Japanese married male physicians. *Int Arch Occup Environ Health* 84:937–942.

Okechukwu CA, El Ayadi AM, Tamers SL, Sabbath EL, Berkman L. 2012. Household food insufficiency, financial strain, work-family spillover, and depressive symptoms in the working class: The work, family, and health network study. *Am J Public Health* 102:126–133.

Pisarski A, Lawrence SA, Bohle P, Brook C. 2008. Organizational influences on the work life conflict and health of shiftworkers. *Appl Ergon* 39:580–588.

Reis JP, Dubose KD, Ainsworth BE, Macera CA, Yore MM. 2005. Reliability and validity of the occupational physical activity questionnaire. *Med Sci Sports Exerc* 37:2075–2083.

Sabbath EL, Melchior M, Goldberg M, Zins M, Berkman LF. 2012. Work and family demands: Predictors of all-cause sickness absence in the GAZEL cohort. *Eur J Public Health* 22:101–106.

Smith DR, Ohmura K, Yamagata Z, Minai J. 2003. Musculoskeletal disorders among female nurses in a rural Japanese hospital. *Nurs Health Sci* 5:185–188.

Sorensen G, Stoddard AM, Stoffel S, Buxton O, Sembajwe G, Hashimoto D, Dennerlein JT, Hopcia K. 2011. The role of the work context in multiple wellness outcomes for hospital patient care workers. *J Occup Environ Med* 53:899–910.

Takeuchi T, Yamazaki Y. 2010. Relationship between work-family conflict and a sense of coherence among Japanese registered nurses. *Jpn J Nurs Sci* 7:158–168.

Trinkoff AM, Le R, Geiger-Brown J, Lipscomb J, Lang G. 2006. Longitudinal relationship of work hours, mandatory overtime, and on-call to musculoskeletal problems in nurses. *Am J Ind Med* 49:964–971.

Wang J, Afifi TO, Cox B, Sareen J. 2007. Work-family conflict and mental disorders in the United States: Cross-sectional findings from The National Comorbidity Survey. *Am J Ind Med* 50:143–149.