

High hsCRP Is Associated With Reduced Lung Function in Structural Firefighters

Denise M. Gaughan, SD,^{1,2} David C. Christiani, MD, MPH, MS,² Michael D. Hughes, PhD,³ Dorothee M. Baur, MD, MSc,² Lester Kobzik, MD,⁴ Gregory R. Wagner, MD,² and Stefanos N. Kales, MD, MPH^{2*}

Background *To assess the association between markers of systemic inflammation and pulmonary function in a population of structural firefighters.*

Methods *We studied male career members of a large Midwestern fire department with questionnaires, spirometry, and high-sensitivity C-reactive protein (hsCRP) as a biomarker of systemic inflammation. We examined percent predicted forced expiratory volume in 1 s (FEV₁%-predicted) and forced vital capacity (FVC%-predicted).*

Results *Complete data were available for 401 firefighters. Higher hsCRP levels were associated with lower lung function values, after adjusting for confounding variables. Specifically, for every twofold increase in log₁₀-hsCRP, FEV₁%-predicted decreased by a mean 1.5% (95% CI: 0.4, 2.6%) and FVC%-predicted decreased by a mean 1.4% (95% CI: 0.4, 2.3%).*

Conclusion *hsCRP as a biomarker of systemic inflammation may indicate reduced lung function in structural firefighters. Am. J. Ind. Med. 57:31–37, 2014. © 2013 Wiley Periodicals, Inc.*

KEY WORDS: *percent predicted forced expiratory volume in 1 s (FEV₁%-predicted); percent predicted forced vital capacity (FVC%-predicted); high-sensitivity C-reactive protein (hsCRP)*

INTRODUCTION

Chronic obstructive pulmonary disease (COPD) is the fourth leading cause of death in the U.S. and a major cause of long-term disability. While cigarette smoking is the probable

cause of COPD in the majority of cases, it has been estimated that about 15% of COPD in the overall population may be due to occupational exposures, and a much higher portion of COPD among non-smokers may be due to occupational exposures [Hnizdo et al., 2002]. Firefighters, both municipal (structural) and wildland are known to face respiratory hazards [Musk et al., 1979; Chia et al., 1990; Materna et al., 1992; Scannell and Balmes, 1995; Bergstrom et al., 1997; Burgess et al., 2001; Prezant et al., 2002; Greven et al., 2012]. Researchers have identified particulates and other exposures, heavy physical exertion and cardiovascular strain to be among the chief, non-traumatic health hazards associated with structural firefighting [Guidotti and Clough, 1992; Austin et al., 2001; Fahy, 2005].

Various studies have associated increased serum concentrations of C-reactive protein (CRP), a chemokine, primarily synthesized in the liver but also produced in the respiratory epithelium [Gould and Weiser, 2001], to reduced lung function in healthy populations [Mendall et al., 2000; Cirillo et al., 2002; Mannino et al., 2003; Aronson et al., 2004; Kony et al., 2004; Gan et al., 2005; Walter et al., 2008; Thorleifsson et al., 2009]. Kony and coworkers studied serum

¹Department of Preventive Medicine, Institute for Translational Epidemiology, Icahn School of Medicine at Mount Sinai, New York, New York

²Department of Environmental Health (Environmental and Occupational Medicine and Epidemiology Program), Harvard School of Public Health, Boston, Massachusetts

³Department of Biostatistics, Harvard School of Public Health, Boston, Massachusetts

⁴Department of Environmental Health (Molecular and Integrative Physiological Sciences), Harvard School of Public Health, Boston, Massachusetts

Contract grant sponsor: Federal Emergency Management Agency (FEMA); Contract grant sponsor: Assistance to Firefighters Grant (AFG); Contract grant numbers: EMW-2006-FP-01493; EMW-2009-FP-00835.

Disclosure Statement: The authors report no conflicts of interests.

*Correspondence to: Stefanos N. Kales, Department of Environmental Health (Environmental and Occupational Medicine and Epidemiology Program), Building 1, Room 1407, Harvard School of Public Health, Boston, MA. E-mail: skales@hsph.harvard.edu

Accepted 23 August 2013

DOI:10.1002/ajim.22260. Published online 20 September 2013 in Wiley Online Library (wileyonlinelibrary.com).

CRP levels, forced expiratory volume in 1 s (FEV₁), and bronchial hyperresponsiveness (BHR) as measured by methacholine challenge, in 259 adults free of cardiovascular disease or respiratory infection. The authors observed that the mean FEV₁ (adjusted for potential confounders such as age, gender, height and smoking status) was lower in subjects with a high CRP level and BHR was more frequent (41.9% vs. 24.9%) in these individuals than in subjects with lower CRP levels [Kony et al., 2004]. Additionally, Aronson et al. [2004] also observed a strong inverse association between CRP levels and quartiles of FEV₁ as well as forced vital capacity (FVC).

Data collected in the Third National Health and Nutrition Examination Survey (NHANES III) also found associations between increased CRP level and reduced pulmonary function [Cirillo et al., 2002; Mannino et al., 2003; Gan et al., 2005]. Moreover, when Gan et al. [2005] focused on smoking status, the authors observed an additive effect of current smoking on the inverse relationship of CRP, FEV₁ observed previously in this cohort. Similar results were recently observed in a cohort of FDNY World Trade Center disaster first responders [Weiden et al., 2013]. The authors report that individuals with elevated CRP levels obtained within 6 months of 9/11/01 had a significantly increased risk of developing decreased lung function (FEV₁) at their subsequent pulmonary testing, independent of a history of CVD. Two case-control studies of both childhood and adult asthmatics also found increased CRP levels correlated with decreased pulmonary function in steroid-naïve participants [Takemura et al., 2006; Deraz et al., 2012]. Takemura and coworkers additionally observed a strong correlation between sputum eosinophils and CRP levels in adult asthmatics, however this was not observed in children [Deraz et al., 2012], possible reflecting the differences in disease etiology.

Genetic studies to date of hsCRP polymorphisms or haplotypes have yielded inconclusive results [Yanbaeva et al., 2009; van Durme et al., 2009; Bolton et al., 2011]. Specifically, while both Bolton and van Durme observed strong inverse associations between CRP and lung function, neither study was able to identify a CRP polymorphism or haplotype associated with pulmonary dysfunction. Thus, while the association has been described in the literature, the mechanism remains unclear.

We hypothesized that hsCRP as a biomarker of systemic inflammation may indicate reduced lung function in structural firefighters.

MATERIALS AND METHODS

Subjects

We studied male career members of a large Midwestern municipal fire department with spirometry, blood, exercise tolerance, and questionnaire data from 2007 to 2009. The

study was approved by the Harvard School of Public Health Human Subjects Committee and the local institutional review board. Informed consent was obtained in writing from each participant prior to data collection. Information on firefighters who chose not to consent or participate was not available.

Venous Blood Samples

Whole non-fasting blood samples were collected by venous phlebotomy in EDTA tubes, and buffy coat were extracted and stored in cell lyses solution at -20°C for analyses. High sensitivity CRP concentration was determined by immunochemiluminometric assay on a Roche COBAS Integra[®] (detection limit of 0.1 mg/L). These analyses were performed by LabCorp (Burlington, NC).

Questionnaire

Participants were administered a health, lifestyle, and occupational history questionnaire while undergoing their fire department's standard medical examination [Durand et al., 2011]. For the purposes of this analysis, we examined responses to history of pulmonary disease including asthma, COPD, diabetes, heart disease, or hypertension.

Spirometry

FEV₁ and FVC were obtained from participants in the standing position by trained technicians following ATS guidelines [Miller et al., 2005] using a Welch Allyn[®] PC-Based Spirometer Model SP (Welch Allyn, Inc., Skaneateles Falls, NY). Predicted values were derived from Knudson's 1983 tables [Knudson et al., 1983]. All measurements were taken pre-bronchodilator.

Cardiorespiratory Fitness

Cardiorespiratory fitness was determined from a symptom limited maximal treadmill exercise testing with electrocardiogram monitoring and estimation of oxygen consumption (METS) following the Bruce protocol. The participants were encouraged to continue exercise until volitional exhaustion, even after exceeding 85% of their maximum predicted heart rate defined as (220-age). The cohort achieved an average of 98.9% (SD 5.9) of maximal age-predicted heart rate on these tests. During the exercise test, total treadmill time in seconds (ETTT) and maximum METS (maxMETS) achieved were recorded. The details of this analysis in this cohort have been previously reported [Baur et al., 2011].

Statistical Methods

Descriptive statistics were calculated for demographic and clinical variables. Ordinary least squares regression techniques were used to examine associations between percent predicted values of FEV₁ (FEV₁%-predicted), predicted values of FVC (FVC%-predicted), as well as the ratio of FEV₁ to FVC using the following predictor variables: current smoker, history of pulmonary disease, body mass index (BMI) which was calculated by dividing the weight (kg) by height (cm squared), maxMETs, resting systolic and diastolic blood pressures, current hypertension (resting systolic blood pressure \geq 140 mm Hg or resting diastolic blood pressure \geq 90 mm Hg). We also examined hsCRP levels and weight (kg), which were not normally distributed and were log₁₀-transformed to achieve an approximate normal distribution. HsCRP was additionally categorized and examined based on distributional quartiles. Specifically, \leq 0.57 mg/L (n = 103) (referent category), $>$ 0.57 to \leq 1.08 (n = 101) (Q2), $>$ 1.08 to \leq 2.48 mg/L (n = 102) (Q3), and $>$ 2.48 mg/L (n = 97) (Q4). Multivariable models were chosen based on an initial evaluation using stepwise selection techniques. Variables that were retained were significant with $P < 0.05$. Overall goodness of fit of the models was assessed using the likelihood ratio test, a statistical tool that compares the fit of the null and alternative models.

Airways obstruction on spirometry was defined according to the Global Initiative for Chronic Obstructive Lung Disease criteria for COPD using the FEV₁, FVC, and FEV₁/FVC. Severe COPD (FEV₁/FVC $<$ 0.70 and FEV₁ $<$ 50% predicted), moderate COPD (FEV₁/FVC $<$ 0.70 and FEV₁ \geq 50% to $<$ 80% predicted), and mild COPD (FEV₁/FVC $<$ 0.70 and FEV₁ = 80%). Restriction was defined as an FEV₁ \geq 70% and FVC $<$ 80% predicted [Pauwels et al., 2001]. All analyses were performed using SAS statistical software (version 9.3).

RESULTS

Characteristics of the population are summarized in Table I. Complete data were available for 401 individuals. Participants had a median age of 36 years. Sixteen (4%) reported a history of pulmonary disease (e.g., asthma and or chronic bronchitis) and 16 (4%) reported active pulmonary disease. One participant reported a history of heart disease. We did not record information on race/ethnicity; however, we know that fire departments where our subjects work are 85–90% White, as previously documented [Poston et al., 2011].

Pulmonary Function

Pulmonary function tests results are summarized in Table II. The median FEV₁%-predicted was 95% (10th, 90th

TABLE I. Subject Characteristics at Examination

Variable	N = 401
Age (years)	36 (27, 47) ^a
Male gender, n (%)	401 (100)
Body mass index (kg/m ²)	29 (24, 36) ^a
Weight (kg)	91.6 (74.4, 113) ^a
Resting diastolic blood pressure (mm Hg)	80 (70, 90) ^a
Resting systolic blood pressure (mm Hg)	120 (108, 138) ^a
MaxMETs achieved during ETT ^b	10.1 (7.9, 13.2) ^a
Current smoker, n (%)	34 (8.5)
Former smoker, n (%)	55 (14)
Pulmonary disease (ever), n (%) ^c	16 (4)
Current hypertension, n (%) ^d	73 (18)
History of heart disease, n (%) ^e	1 (0.2)
High sensitivity C-reactive protein (μ g/ml)	1.06 (0.3, 4.9) ^a

^aMedian and 10th, 90th percentiles.

^bMaximum metabolic equivalent (MET) achieved during exercise tolerance test (ETT).

^cHistory of pulmonary disease including asthma and COPD.

^dResting systolic blood pressure \geq 140 mm Hg or resting diastolic blood pressure \geq 90 mm Hg.

^eHistory of heart disease includes diabetes, heart disease, or hypertension.

percentiles: 80%, 115%), the median FVC%-predicted was 99% (10th, 90th percentiles: 84%, 117%) and the median ratio was 81% (10th, 90th percentiles: 73%, 86%). Twenty-four participants (6%) had airways obstruction on spirometry (10 moderate, 14 mild) and four other participants' spirometry indicated a restrictive pattern. There were no subjects with mixed disease.

Systemic Inflammation

The median hsCRP concentration level was 1.06 mg/L (10th, 90th percentiles: 0.31 mg/L, 4.85 mg/L). High hsCRP values, defined as greater than 3 mg/L [Pearson et al., 2003], were observed in 83 participants (21%). There were no hsCRP values below the limit of detection.

Pulmonary Function and Systemic Inflammation

Figure 1 shows mean FEV₁%-predicted and FVC %-predicted values by hsCRP quartiles. Higher mean hsCRP levels were associated with lower mean lung function values. The mean FEV₁ for participants in the highest hsCRP quartile (93.3%), was significantly lower than for individuals in the lowest quartile (99.4%; $P < 0.01$). This difference was even more pronounced for FVC%-predicted. Specifically, the mean FVC%-predicted for participants in the highest hsCRP quartile (94.8%) was significantly lower than the

TABLE II. Pulmonary Function Test Results

Variable	N = 401
Percent predicted forced expiratory volume in 1 s (FEV ₁ %-predicted)	95 (80, 115) ^a
Percent predicted forced vital capacity (FVC%-predicted)	99 (84, 117) ^a
FEV ₁ /FVC (%)	81 (73, 86) ^a
Moderate COPD, n (%)	10 (3)
Mild COPD, n (%)	14 (4)
Restrictive lung disease, n (%)	4 (1)

^aMedian and 10th, 90th percentiles.

lowest quartile (104%) ($P < 0.001$), as well as values in the second (99.7%; $P < 0.05$) and third quartiles (100%) ($P < 0.05$).

In multivariable analysis, higher hsCRP levels were associated with lower lung function values and are detailed in Tables III and IV. Specifically, for every 1 log₁₀ mg/L increase log₁₀ hsCRP, mean FEV₁%-predicted decreased by 5.3% (95% CI: 1.6, 8.9%) after adjusting for BMI, resting systolic blood pressure, and achieved MaxMETs. This corresponds to an adjusted mean reduction of 1.6% (95% CI: 0.5, 2.7%) for each twofold increase in hsCRP. Current smoker, history of pulmonary disease, current hypertension, log₁₀ weight and resting diastolic blood pressure were not associated with FEV₁%-predicted in this cohort. Moreover, for every twofold increase in hsCRP, FVC%-predicted decreased by a mean 1.4% (95% CI: 0.4, 2.4%) (regression estimate for log₁₀ hsCRP: -0.47 (95% CI: -0.79, -0.15)) after adjusting for log₁₀ weight. Current smoker, history of pulmonary disease, current hypertension, achieved MaxMETs, resting systolic blood pressure, resting diastolic blood pressure and BMI were not associated FVC%-predicted in this cohort.




FIGURE 1. Mean FEV₁ and FVC%-predicted values by hsCRP category. ***Significant from Q1 (referent) at <0.001 . **Significant from Q1 (referent) at <0.01 . *Significant from Q1 (referent) at <0.05 .

DISCUSSION

Lung function declines, specifically FEV₁, are established risk factors for CVD, rivaling that of cholesterol [Sin et al., 2005]. Similar to other studies, we observed an inverse relationship between higher hsCRP and lung function as measured by FEV₁%-predicted and FVC%-predicted, after adjusting for established potential confounders. Unlike Gan et al. we did not observe an additive effect of current smoking on FEV₁%-predicted in our cohort. However, given the limited number of subjects currently smoking, we may not have had sufficient statistical power to observe such a difference.

While numerous studies have found a negative association between lung function reduction and increasing CRP, the mechanism remains unclear. Genetic studies to date of hsCRP polymorphisms or haplotypes have not helped elucidate the mechanism [Yanbaeva et al., 2009; van Durme et al., 2009; Bolton et al., 2011]. In the absence of definitive genetic studies some researchers have suggested reverse causality [Bolton et al., 2011] or an undefined mediator between lung function and CRP may be driving the association between the two outcomes. CRP has also been linked to the development of ischemic heart disease and stroke, the two primary causes of death in individuals with COPD. Higher levels of systemic inflammation may simply represent spill over from inflammation in the airways. Concurrent examination of systemic inflammation, eosinophilic and neutrophilic, from induced sputum, in this population may assist in answering this question.

As previously noted, CVD events are the leading cause of morbidity and mortality in structural firefighters affecting 1 in 1,000 structural firefighters annually [Fahy, 2005; Soteriades et al., 2011]. Higher BMI and lower cardiorespiratory fitness, as measured by achieved MaxMETs are among the factors associated with an elevated CVD risk profile in firefighters [Baur et al., 2011]. We observed a borderline significant association between higher cardiorespiratory fitness, as measured by achieved MaxMETs and higher FEV₁%-predicted values but not vital capacity. We observed an association between higher FEV₁%-predicted and higher BMI and an inverse relationship between higher FEV₁%-predicted and resting systolic blood pressure after adjusting for hsCRP and MaxMETs.

There is evidence that particulate matter (PM) produced by fires may be more toxic than PM from ambient air due to chemical components found in the smoke [Wegesser et al., 2009]. Leonard et al. [2007] examined six sets of aerodynamically size-selected aerosol samples at a large fire. The authors found smoke particles in all size fractions, with highest overall mass fraction on filters in the fine range. Ultrafine particles ($<0.1 \mu\text{m}$ diameter) may produce greater inflammation than larger particles, simply due to a much larger total surface area to volume ratio of the former. Ultrafine particles may escape phagocytosis by alveolar

TABLE III. Predictors of FEV₁%-Predicted, OLS Estimates and 95% CIs

Variable ^a	Unadjusted	Adjusted ^b
HsCRP (per 1 log ₁₀ mg/L higher)	-4.2 (-7.3, -1.0)	-5.3 (-8.9, -1.6)
Body mass index (per 1 kg/m ² higher)	0.1 (-0.3, 0.4)	0.6 (0.2, 1.0)
Resting systolic blood pressure (per 10 mm Hg higher)	-1.4 (-2.6, -0.1)	-1.7 (-2.9, -0.4)
Achieved maxMETs (per 1 MET higher)	0.6 (-0.1, 1.3)	0.8 (-0.0, 1.7)

OLS, Ordinary least squares.

^aCurrent smoker, history of pulmonary disease, current hypertension, log₁₀ weight and resting diastolic blood pressure were not associated with FEV₁%-predicted in this cohort and so were not included in the regression model.

^bEstimates from the regression model that included all four variables shown.

macrophages, easily gaining access to the pulmonary interstitium. The severity of airways inflammation due to particulate exposures is correlated with particle surface area [Brown et al., 2001; Oberdorster, 2001].

Additionally, smoke particulate may be associated with adverse respiratory health outcomes as a result of the generation of toxic free radicals [Bankson et al., 1993; Jankovic et al., 1993; Leonard et al., 2000]. Leonard et al. [2007] used electron spin resonance to measure carbon-centered as well as hydroxyl radicals produced by a Fenton-like reaction with smoke. The authors observed that OH radical precursors were more prevalent (per unit mass) in the ultrafine-sized particles and caused a significant increase in cellular H₂O₂ production, DNA strand breaks, and reactive oxygen species (ROS)-induced lipid peroxidation as measured by malondialdehyde (MDA). Thus, exposure to chemicals and fine particulates routinely encountered during structural firefighters overhaul operations, where respiratory protection is generally not donned [Burgess et al., 2001] could generate ROS, potentially leading to oxidative damage and inflammation and may explain the increased risk of lower lung dysfunction in this healthy worker population comprising primarily non-current smokers. Moreover, Weiden et al., observed a significant inverse association between CRP and

TABLE IV. Predictors of FVC%-Predicted, OLS Estimates and 95% CIs

Variable ^a	Unadjusted	Adjusted ^b
HsCRP (per 1 log ₁₀ mg/L)	-5.9 (-8.8, -3.0)	-4.7 (-7.9, -1.5)
Weight (per 1 log ₁₀ lb higher)	-30.6 (-48.8, -12.5)	-19.6 (-39.1, -0.05)

OLS, ordinary least squares.

^aCurrent smoker, history of pulmonary disease, current hypertension, achieved maxMETs, resting systolic blood pressure, resting diastolic blood pressure and BMI were not associated FVC%-predicted in this cohort and so were not included in the model.

^bEstimate from the regression model that included both variables shown.

FEV₁ in firefighters exposed to PM generated by the World Trade Center Disaster [Weiden et al., 2013]. The authors hypothesized that the increased systemic CRP levels could reflect pulmonary vessel damage. It is of note, that some of the mean predicted values for lung function we observed were lower than the referent population. However, our study lacked a means to quantify this type of exposure. Further research is needed to examine how our findings might relate to exposure.

We examined hsCRP which minimizes the biological variation in the samples. Only obtaining one measure of hsCRP, which varies by season [Sung, 2006], could have affected our results. However, the testing took place over the course of a few years year so any potential misclassification would be non-differential.

We did not observe a difference among individuals with a history of pulmonary or cardiovascular disease. However, given the small number of individuals with these outcomes, 16 and one respectively, we may have lacked sufficient statistical power to observe one if it did exist.

Our study has a number of limitations. We examined only one biomarker of inflammation. Future studies should examine sputum biomarkers of neutrophilic inflammation, such as myeloperoxidase, which has been linked to occupationally-related airways obstruction [Jung and Park, 1999]. We do not have exposure data on participants which limited our ability to examine dose-response relationships. Moreover, we lack lower respiratory symptom data such as wheeze and shortness of breath, which limits the interpretation of our objective findings as they relate to overall well-being and function. Finally, our study population (n = 401) with complete data that was analyzed constituted 98% of the male members of the fire department who consented to participate. While the possibility of selection bias does exist, we do know that our consented population has a very similar age, BMI, and expected ethnic composition to that found in a randomly sampled, population-based study of career firefighters [Poston et al., 2011].

hsCRP, a biomarker of systemic inflammation potentially generated from lifestyle and occupational exposures including smoke particulate, is associated with diminished exercise tolerance and oxidative stress. Damage reflected in elevated HsCRP may result in reduced lung function in structural firefighters. Our results corroborate the findings of an inverse relationship between CRP and lung function observed in both occupational settings and the general population. The long-term significance of these findings to the cardiopulmonary health of structural firefighters, including the relationship to exposure, should be investigated further.

ACKNOWLEDGEMENTS

This work was supported by Federal Emergency Management Agency (FEMA) (Washington D.C., USA),

Assistance to Firefighters Grant (AFG) program's awards EMW-2006-FP-01493, and EMW-2009-FP-00835. And NIH center grant support ES00002. Dr. Gaughan additionally received training-grant support (T42 OH008416). The authors additionally thank Elena Rahona for assistance with graphics.

REFERENCES

- Aronson D, Roterman I, Yigla M, Kerner A, Avizohar O, Sella R, Bartha P, Levy Y, Markiewicz W. 2004. Inverse association between pulmonary function and C-reactive protein in apparently healthy subjects. *Inhal Toxicol* 16:607–613.
- Austin C, Wang D, Ecobichon DJ, Dussault G. 2001. Characterization of volatile organic compounds in smoke at municipal structural fires. *J Toxicol Environ Health* 63:437–458.
- Bankson DD, Kestin M, Rifai N. 1993. Role of free radicals in cancer and atherosclerosis. *Clin Lab Med* 13:463–480.
- Baur DM, Christophi CA, Tsismenakis AJ, Cook EF, Kales SN. 2011. Cardiorespiratory fitness predicts cardiovascular risk profiles in career firefighters. *J Occup Environ Med* 53:1155–1160.
- Bergstrom CE, Eklund A, Skold M, Tornling G. 1997. Bronchoalveolar lavage findings in firefighters. *Am J Ind Med* 32:332–336.
- Bolton CE, Schumacher W, Cockcroft JR, Timpson NJ, Smith GD, Gallacher J, Rumley A, Lowe GD, Ebrahim S, Shale DJ, Ben-Shlomo Y. 2011. The CRP genotype, serum levels and lung function in men: The Caerphilly Prospective Study. *Clin Sci (Lond)* 120:347–355.
- Brown DM, Wilson MR, MacNee W, Stone V, Donaldson K. 2001. Size-dependent proinflammatory effects of ultrafine polystyrene particles: A role for surface area and oxidative stress in the enhanced activity of ultrafines. *Toxicol Appl Pharmacol* 175:191–199.
- Burgess JL, Nanson CJ, Bolstad-Johnson DM, Gerkin R, Hysong TA, Lantz RC, Sherrill DL, Crutchfield CD, Quan SF, Bernard AM, Witten ML. 2001. Adverse respiratory effects following overhaul in firefighters. *J Occup Environ Med* 43:467–473.
- Chia KS, Jeyaratnam J, Chan TB, Lim TK. 1990. Airway responsiveness of firefighters after smoke exposure. *Br J Ind Med* 47:524–527.
- Cirillo DJ, Agrawal Y, Cassano PA. 2002. Lipids and pulmonary function in the Third National Health and Nutrition Examination Survey. *Am J Epidemiol* 155:842–848.
- Deraz TE, Kamel TB, El-Kerdany TA, El-Ghazoly HM. 2012. High-sensitivity C reactive protein as a biomarker for grading of childhood asthma in relation to clinical classification, induced sputum cellularity, and spirometry. *Pediatr Pulmonol* 47:220–225.
- Durand G, Tsismenakis AJ, Jahnke SA, Baur DM, Christophi CA, Kales SN. Firefighters' physical activity: Relation to fitness and cardiovascular disease risk. *Med Sci Sports Exerc* 2011. 43:1752–1759.
- Fahy R. 2005. U.S. firefighters fatalities due to sudden cardiac death, 1995–2004. Quincy, MA: National Fire Protection Association. Accessed on-line 1/23/13, <http://www.nfpa.org/assets/files/PDF/OSCardiacDeath.pdf>
- Gan WQ, Man SF, Sin DD. 2005. The interactions between cigarette smoking and reduced lung function on systemic inflammation. *Chest* 127:558–564.
- Gould JM, Weiser JN. 2001. Expression of C-reactive protein in the human respiratory tract. *Infect Immun* 69:1747–1754.
- Greven FE, Krop EJ, Spithoven JJ, Burger N, Rooyackers JM, Kerstjens HA, van der Heide S, Heederik DJ. 2012. Acute respiratory effects in firefighters. *Am J Ind Med* 55:54–62.
- Guidotti TL, Clough VM. 1992. Occupational health concerns of firefighting. *Annu Rev Public Health* 13:151–171.
- Hnizdo E, Sullivan PA, Bang KM, Wagner G. 2002. Association between chronic obstructive pulmonary disease and employment by industry and occupation in the US population: A study of data from the Third National Health and Nutrition Survey. *Am J Epidemiol* 156:738–746.
- Jankovic J, Jones W, Castranova V, Dalal N. 1993. Measurement of short-lived reactive species and long-lived free radicals in air samples from structural fires. *Appl Occup Environ Hyg* 8:650–654.
- Jung KS, Park HS. 1999. Evidence for neutrophil activation in occupational asthma. *Respirology* 4:303–306.
- Knudson RJ, Lebowitz MD, Holberg CJ, Burrows B. 1983. Changes in the normal maximal expiratory flow-volume curve with growth and aging. *Am Rev Respir Dis* 127:725–734.
- Kony S, Zureik M, Driss F, Neukirch C, Leynaert B, Neukirch F. 2004. Association of bronchial hyperresponsiveness and lung function with C-reactive protein (CRP): A population based study. *Thorax* 59:892–896.
- Leonard SS, Wang S, Shi X, Castranova V, Dubik MA. 2000. Wood smoke particles generate free radicals and cause lipid peroxidation, DNA damage, NF-kappaB activation and TNF-alpha release from macrophages. *Toxicology* 150:147–157.
- Leonard SS, Castranova V, Chen BT, Schwegler-Berry D, Hoover M, Piacitelli C, Gaughan DM. 2007. Particle size-dependent radical generation from wildland fire smoke. *Toxicology* 236:103–113.
- Mannino DM, Ford ES, Redd SC. 2003. Obstructive and restrictive lung disease and markers of inflammation: Data from the Third National Health and Nutrition Examination. *Am J Med* 114:758–762.
- Materna BL, Jones JR, Sutton PM, Rothman N, Harrison RJ. 1992. Occupational exposures in California wildland firefighting. *Am Ind Hyg Assoc J* 53:69–76.
- Mendall MA, Strachan DP, Butland BK, Ballam L, Morris J, Sweetnam PM, Elwood PC. 2000. C-Reactive protein: Relation to total mortality, cardiovascular mortality and cardiovascular risk factors in men. *Eur Heart J* 21:1584–1590.
- Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, Crapo R, Enright P, van der Grinten CP, Gustafsson P, Jensen R, Johnson DC, MacIntyre N, McKay R, Navajas D, Pedersen OF, Pellegrino R, Viegi G, Wang J, ATS/ERS Task Force. 2005. Standardisation of spirometry. *Eur Respir J* 26:319–338.
- Musk AW, Smith TJ, Peters JM, McLaughlin E. 1979. Pulmonary function in firefighters: Acute changes in ventilatory capacity and their correlates. *Br J Ind Med* 36:29–34.
- Oberdorster G. 2001. Pulmonary effects of inhaled ultrafine particles. *Int Arch Occup Environ Health* 74:1–8.
- Pauwels RA, Buist AS, Calverley PM, Jenkins CR, Hurd SS, Scientific GOLD Committee. 2001. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. NHLBI/WHO Global Initiative for Chronic Obstructive Lung Disease (GOLD) Workshop summary. *Am J Respir Crit Care Med* 163:1256–1276.
- Pearson TA, Mensah GA, Alexander RW, Anderson JL, Cannon RO III, Criqui M, Fadl YY, Fortmann SP, Hong Y, Myers GL, Rifai N, Smith SC, Jr., Taubert K, Tracy RP, Vinicor F, Centers for Disease Control and Prevention; American Heart Association. 2003. Markers of inflammation and cardiovascular disease: Application to clinical and public health practice: A statement for healthcare professionals from the Centers for

- Disease Control and Prevention and the American Heart Association. *Circulation* 107:499–511.
- Poston WS, Haddock CK, Jahnke SA, Jitnarin N, Tuley BC, Kales SN. 2011. The prevalence of overweight, obesity, and substandard fitness in a population-based firefighter cohort. *J Occup Environ Med* 53:266–273.
- Prezant DJ, Weiden M, Banauch GI, McGuinness G, Rom WN, Aldrich TK, Kelly KJ. 2002. Cough and bronchial responsiveness in firefighters at the World Trade Center site. *N Engl J Med* 347:806–815.
- Scannell CH, Balmes JR. 1995. Pulmonary effects of firefighting. *Occup Med* 10:789–801.
- Sin DD, Wu L, Man SFP. 2005. The relationship between reduced lung function and cardiovascular mortality: A population-based study and a systemic review of the literature. *Chest* 127:1952–1959.
- Soteriades ES, Smith DL, Tsismenakis AJ, Baur DM, Kales SN. 2011. Cardiovascular disease in US firefighters: A systematic review. *Cardiol Rev* 194:202–215.
- Sung KC. 2006. Seasonal variation of C-reactive protein in apparently healthy Koreans. *Int J Cardiol* 107:338–342.
- Takemura M, Matsumoto H, Niimi A, Ueda T, Matsuoka H, Yamaguchi M, Jinnai M, Muro S, Hirai T, Ito Y, Nakamura T, Mio T, Chin K, Mishima M. 2006. High sensitivity C-reactive protein in asthma. *Eur Respir J* 27:908–912.
- Thorleifsson SJ, Margretardottir OB, Gudmundsson G, Olafsson I, Benediktsdottir B, Janson C, Buist AS, Gislason T. 2009. Chronic airflow obstruction and markers of systemic inflammation: Results from the BOLD study in Iceland. *Respir Med* 103:1548–1553.
- Van Durme YM, Verhamme KM, Aarnoudse AJ, Van Pottelberge GR, Hofman A, Witterman JC, Joos GF, Brusselle GG, Stricker BH. 2009. C-reactive protein levels, haplotypes, and the risk of incident chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 179:375–382.
- Walter RE, Wilk JB, Larson MG, Vasani RS, Keaney JF Jr, Lipinska I, O'Connor GT, Benjamin EJ. 2008. Systemic inflammation and COPD: The Framingham Heart Study. *Chest* 133:19–25.
- Wegesser TC, Pinkerton KE, Last JA. 2009. California Wildfires of 2008: Coarse and fine particulate matter toxicity. *Environ Health Perspect* 117:893–897.
- Weiden MD, Naveed B, Kwon S, Cho SJ, Comfort AL, Prezant DJ, Rom WN, Nolan A. 2013. Cardiovascular biomarkers predict susceptibility to lung injury in World Trade Center dust-exposed firefighters. *Eur Respir J* 4:1023–1030.
- Yanbaeva DG, Dentener MA, Spruit MA, Houwing-Duistermaat JJ, Kotz D, Passos VL, Wouters EF. 2009. IL6 and CRP haplotypes are associated with COPD risk and systemic inflammation: A case-control study. *BMC Med Genet* 9:10–23.