

Presence of organophosphorus pesticide oxygen analogs in air samples

Jenna L. Armstrong ^{a,*}, Richard A. Fenske ^a, Michael G. Yost ^a, Kit Galvin ^a, Maria Tchong-French ^a, Jianbo Yu ^{a,b}

^a Department of Environmental and Occupational Health Sciences, University of Washington School of Public Health, Health Sciences Building, 1959 NE Pacific Street, Box 357234, Seattle, WA 98195, USA

^b Environmental Health Laboratory and Trace Organics Analysis Center, University of Washington, Health Sciences Building F455, 1959 NE Pacific Street, Seattle, WA 98195, USA

HIGHLIGHTS

- Chlorpyrifos-oxon is formed during air sampling with matrices containing XAD-2.
- Chemical analysis without measurement of oxon underestimates air concentrations.
- Comparisons of lab and field data suggest that oxon is present in community air.
- Accounting for chemical mixtures of parent OPs and oxons in air is important.
- Small amounts of oxon in air have a large effect on human health risk estimates.

ARTICLE INFO

Article history:

Received 5 February 2012

Received in revised form

26 June 2012

Accepted 28 June 2012

Keywords:

Air monitoring

Exposure

Organophosphorus pesticides

Oxon

Toxicity equivalent concentration

XAD resin

ABSTRACT

A number of recent toxicity studies have highlighted the increased potency of oxygen analogs (oxons) of several organophosphorus (OP) pesticides. These findings were a major concern after environmental oxons were identified in environmental samples from air and surfaces following agricultural spray applications in California and Washington State. This paper reports on the validity of oxygen analog measurements in air samples for the OP pesticide, chlorpyrifos. Controlled environmental and laboratory experiments were used to examine artificial formation of chlorpyrifos-oxon using OSHA Versatile Sampling (OVS) tubes as recommended by NIOSH method 5600. Additionally, we compared expected chlorpyrifos-oxon attributable to artificial transformation to observed chlorpyrifos-oxon in field samples from a 2008 Washington State Department of Health air monitoring study using non-parametric statistical methods. The amount of artificially transformed oxon was then modeled to determine the amount of oxon present in the environment. Toxicity equivalency factors (TEFs) for chlorpyrifos-oxon were used to calculate chlorpyrifos-equivalent air concentrations. The results demonstrate that the NIOSH-recommended sampling matrix (OVS tubes with XAD-2 resin) was found to artificially transform up to 30% of chlorpyrifos to chlorpyrifos-oxon, with higher percentages at lower concentrations ($<30 \text{ ng m}^{-3}$) typical of ambient or residential levels. Overall, the 2008 study data had significantly greater oxon than expected by artificial transformation, but the exact amount of environmental oxon in air remains difficult to quantify with the current sampling method. Failure to conduct laboratory analysis for chlorpyrifos-oxon may result in underestimation of total pesticide concentration when using XAD-2 resin matrices for occupational or residential sampling. Alternative methods that can accurately measure both OP pesticides and their oxygen analogs should be used for air sampling, and a toxicity equivalent factor approach should be used to determine potential health risks from exposures.

© 2012 Elsevier Ltd. All rights reserved.

1. Introduction

Although banned for residential use in 2000, the common organophosphorus pesticide chlorpyrifos (CPF) remains an

effective tool for agricultural protection from more than 200 pests. CPF continues to be widely used because few chemical substitutes are as effective. In Washington State, CPF is often applied in aerosolized form to tree fruits and vegetables using a large sprayer attached behind a tractor. A number of past studies have found that some semi-volatile organophosphorus pesticides (OP) can drift in air far from source of application, as indicated in past community studies (see Table 1). Recent monitoring in the agricultural regions

* Corresponding author.

E-mail address: jennaa@uw.edu (J.L. Armstrong).

Table 1

Occupational and community chlorpyrifos and chlorpyrifos-oxon air monitoring studies since 1996.

Air concentration range > MDL ^a (ng m ⁻³)		Sampling location	Sampling medium (resin)	Analytical method	MDL ^a (ng m ⁻³)	Reference
Chlorpyrifos	Chlorpyrifos-oxon					
Occupational studies						
48–2000	Not measured	Iowa, North Carolina	XAD-2	LC/MS/MS	NA	Thomas et al., 2010
13,000–54,000	Not measured	Egypt	XAD-2	LC/MS/MS	1.5	Farahat et al., 2010
22,000–56,000	Not measured	Thailand	XAD-2	GC/FPD	1.6	Jaipieam et al., 2010
Community studies						
10–230	Not measured	Iowa	XAD-2	GC–MS	3.5	Curwin et al., 2005
16–1340	16–230	Tulare County, CA	XAD-4	GC–MS	5.25	CARB, 1998b, 1999
7–150	10–28	Parlier, CA	XAD-4	LC/MS	5	CDPR, 2006, 2009
83 (Max)	8.5 (max)	Lompoc, CA	XAD-4	LC/MS	5	CDPR, 2003, 2009
1–2.9	Not measured	Iowa	XAD-2	GC–MS	0.13	Peck and Hornbuckle, 2005
5000 (Max)	Not measured	Thailand	XAD-2	GC/FPD	1.6	Jaipieam et al., 2010
9–494	2–108	Washington	XAD-2	LC/MS/MS	0.35	Fenske et al., 2009
Other studies		Sequoia National Park, CA	XAD-4	GC–MS	5×10^{-4} – 8×10^{-4} ^b	LeNoir et al., 1999

^a Method detection limit.^b High volume sampling.

of California and Washington has reported both the parent compound (CPF) and the presence of chlorpyrifos-oxon (CPF-O), the oxygen analog of CPF, in community air samples.

The airborne presence of the oxons of OP pesticides is a human health concern because *in vivo* toxicity studies have found that the toxicity of the oxon can be 5–100 times more toxic than the parent OP pesticide (Chambers and Carr, 1993; Cole et al., 2005, 2011; Huff et al., 1994; Timchalk et al., 2007). CPF-O is believed to pose a special risk for genetically susceptible individuals who have lower levels of the paraoxonase [PON-1(–/–)] enzyme playing a key role in metabolism of OPs in the body. In addition, young mice have been found to demonstrate changes in specific brain cells and irregular distribution of neurons in the cortical plate after exposures to chronic low doses of CPF-O (Furlong et al., 2005). The presence of oxon in community air samples is a concern because children may be particularly susceptible to its toxicity (Harnly et al., 2005; Costa et al., 2005) due to differences in metabolic functioning during development.

CPF-O may be present in the atmospheric environment, as the transformation of OP pesticides to their oxons has been documented for foliar residues (Popendorf and Leffingwell, 1978; Spear et al., 1978) and in air samples (Seiber et al., 1989). The environmental formation of oxons has been attributed to conditions such as higher temperatures, ozone, dry weather, or photodegradation via ultraviolet light (Spear et al., 1978; Aston and Seiber, 1997; Bavcon Kralj et al., 2007). Very little is known about transformation or changes in levels of CPF-O as the total pesticide mixture drifts from site of application to more residential areas.

CPF-O was identified in virtually all of the 145 air samples collected in agricultural communities of the recent 2008 Washington State Department of Health air monitoring study (Fenske et al., 2009), with higher proportions oxon being reported at residential locations >250 m from agricultural fields. This was a concern for public health because the study area is home to many farmworker families, and greater than a third of the population consists of children and adolescents <18 years of age (US CENSUS, 2010). The oxon represented as much as 94% percent of the total CPF in some cases; this raised the issue of potential transformation during air sampling. It is important to rule out sampling transformation in order for researchers to correctly assess health risk from exposure to airborne CPF-O.

The 2008 Washington study used OSHA Versatile Sampling (OVS) tubes, a popular sampling matrix that contains XAD-2 resin. The OVS tube is part of the standard method recommended by

National Institute for Occupational Safety and Health (NIOSH, 1994), and the XAD-2 resin is identified in current guidance documents as an appropriate sampling matrix by the U.S. Environmental Protection Agency and the American Society for Testing and Materials (USEPA, 1999; ASTM, 2011). The resin has been used in a large number of community and occupational air sampling studies (Table 1).

The primary objective of this study was to determine the degree to which the OP pesticide chlorpyrifos (CPF) can be transformed artificially to its oxygen analog form by air sampling with OVS tubes at levels representative of concentrations measured during community sampling at low and high flow rates. The second objective was to ascertain the extent to which CPF-O may have been environmentally present in the community air that was sampled in the 2008 Washington study. The final objective was to use this data to include the toxicity of CPF-O and estimate potential changes in community health risk resulting from respiratory exposure to the mixture of both parent CPF and CPF-O.

2. Sampling methods

2.1. Laboratory studies

Sampling was conducted according to NIOSH method 5600 for OP pesticides (NIOSH, 1994). OVS tubes containing XAD-2 sorbent were spiked in triplicates at levels of 0, 42, 210, and 2100 ng of 99.5% pure chlorpyrifos (ChemService, Inc. PS-674) in solution. A 25 μ l Hamilton™ positive displacement syringe was used to apply chlorpyrifos in acetone solution directly to the resin by inserting the needle beyond the quartz fiber pre-filter into the first bed of XAD-2. The back-up XAD-2 resin was not spiked because past studies have demonstrated that even higher concentrations of chlorpyrifos (>96%) are primarily trapped on the first resin bed (Shibamoto et al., 1996). Each OVS tube was paired with an SKC air sampling pump (224-PCXR8) operated at a flow rate of either two or 6 L per minute (LPM) for approximately 24 h. Two flow rates were tested to examine potential differences in artificial transformation at higher flow rates and to simulate popular air sampling procedures conducted in past community monitoring studies (CARB, 1998a; CDPR, 2003; Fenske et al., 2009). Larger spike masses were applied at higher flow rates (~6 LPM) to account for larger air sampling volumes. Pumps were pre- and post-calibrated between studies using a DryCal DC-Lite and flow rates were calculated separately for each sample in order to calculate the air volumes for

each spiked sample. Laboratory blanks, spikes, and storage spikes were included in the experiments for quality assurance purposes.

2.2. Field studies

The following spring, data were collected outdoors at a community air sampling site managed by the Washington Department of Ecology in Yakima Valley, WA. The site was previously utilized in the 2008 Washington State study to collect ambient concentrations and is located >1 km (1405 m) from the nearest tree fruit field. OVS tubes containing XAD-2 sorbent were spiked with chlorpyrifos in triplicates at levels of 0, 15, 30, 60, 200, 592, and 2628 ng following the procedures described in the laboratory studies. Previous data from the 2008 study emphasized the need to control for small background concentrations of CPF and CPF-O by including non-spiked samples. Triplicate spikes of 0 and 30 ng were deployed with no pump air flow. All SKC air sampling pumps were stored in a weather-proof container and operated at either 2 or 6 LPM for 24 h. Outdoor temperatures ranged from 4 to 12 °C during the sampling period. Samples were stored at a University of Washington field office in Yakima in a freezer at –10 °C until transport to the University of Washington Environmental Health Laboratory. Flow rates were calibrated and calculated separately in order to calculate the air volumes (m³) for each spiked sample.

The Environmental Health Laboratory performed chemical analysis for both CPF and CPF-O. A new LC-MS-MS method was developed through modifications of methods described by [Sancho et al. \(2000\)](#), [Yusa et al. \(2009\)](#), and NIOSH Method 5600. The quartz fiber filter and primary resin section was placed in a 10 mL vial, and separated from the secondary resin section to analyze for break-through. All samples were sonicated with an acetone/acetonitrile solution containing stable-isotope labeled internal standards of chlorpyrifos diethyl-D10, 99% (Cambridge Isotope Labs DLM-4360) and 13C2, 15N-Chlorpyrifos oxon (donated by Dow Agro Sciences LLC). The limit of detection (LOD) for both chlorpyrifos and CPF-O was 1 ng sample^{–1}. Fortification/recovery studies involved spiking matrix blanks with CPF and CPF-O at levels ranging from 5 to 1000 ng sample^{–1}; recoveries were 86.1–94.6% for CPF and 85.8–97.6% for CPF-O.

Quality assurance field spikes and blanks were carried into the field and handled in a manner similar to other samples except they were capped and did not have air drawn through them. Field spikes were prepared by introducing low levels CPF (20–50 ng sample^{–1}) into the front section of the XAD-2 resin with a micropipette, immediately recapping, followed by storage on ice. Field spike recoveries were 74.4–101.5% for CPF, and yielded no detectable CPF-O. Field blanks yielded no CPF or CPF-O. Static storage spikes were kept in the laboratory freezer at –10 °C for 2 months and recoveries ranged from 84 to 105%.

2.3. Determination of artificial and environmental CPF-O

Total chlorpyrifos was calculated for each sample by converting the measured chlorpyrifos-oxon to its chlorpyrifos equivalent using the ratio of molecular weights, and adding this value to the measured chlorpyrifos, as indicated in Equation (1):

$$\text{CPF}_{\text{Total}} = \text{CPF-O}_m * (\text{CPF}_{\text{MW}} / \text{CPF-O}_{\text{MW}}) + \text{CPF}_m \quad (1)$$

where $\text{CPF}_{\text{Total}}$ = total chlorpyrifos in sample (ng), CPF-O_m = mass of chlorpyrifos-oxon measured in sample (ng), CPF_{MW} = chlorpyrifos molecular weight (350.6 ng nmol^{–1}), CPF-O_{MW} = chlorpyrifos-oxon molecular weight (334.5 ng nmol^{–1}), and CPF_m = mass of chlorpyrifos measured in the sample (ng). To

control for small concentrations levels of CPF and CPF-O, outdoor recovery masses were corrected by subtracting background levels CPF and CPF-O from non-spiked samples in the outdoor environment to calculate total % recovery and % CPF-O.

The percent chlorpyrifos-oxon (% CPF-O) in each sample was calculated by dividing the mass of chlorpyrifos-oxon, expressed as chlorpyrifos equivalent, by total chlorpyrifos and multiplying by 100, as indicated in Equation (2):

$$\% \text{CPF-O} = \text{CPF-O}_m * (\text{CPF}_{\text{MW}} / \text{CPF-O}_{\text{MW}}) / \text{CPF}_{\text{Total}} * 100 \quad (2)$$

Next, % CPF-O (y-axis) was plotted against total mass per unit volume (x-axis), which was estimated by dividing $\text{CPF}_{\text{Total}}$ by specific air sampling volumes. A quadratic prediction plot with 95% confidence intervals was used to model expected oxon as an artifact of sampling ($R^2 = 0.288$) as a function of total CPF air concentration. Community measurements from ambient sites (>500 m from orchard) and near-field sites (<100 m from orchard) from the 2008 Washington State study were directly compared with the laboratory and field experimental data. A non-parametric statistical test was used to compare expected values determined from laboratory and field study samples to observed community measurements from Washington State study data. The Mann Whitney U-test was used since sample size and variability of the two groups differed and the observations were independent of each other ([Fay and Proschan, 2010](#)). The basic assumption of normality was questionable via Shapiro Wilk test and small experiment sample size ([Rosner and Grove, 1999](#)). Due to the variable nature of the relationship between oxon transformation and total concentration chlorpyrifos, statistical tests were performed for the entire data set and for each data tertile of pesticide concentrations. All calculations were performed using STATA™ 10.1 data analysis and statistical software (StataCorp LP College Station, Texas).

3. Results

Data from the laboratory and field experiments are presented in [Table 2](#). In the laboratory experiments, samples with low spike masses CPF (42 ng) had an average of 31.6% oxon transformation whereas higher spike masses CPF (~2100 ng) had an average of only 15.4% oxon transformation. Similarly in the field experiments, low spike masses (<30 ng) yielded an average of 24.4% oxon, whereas medium spike masses (60–200 ng) yielded only 9.9% oxon. A similar effect was found at higher flow rates of 6 LPM. Samples hung outdoors with no flow rate (0 LPM) resulted in no conversion to CPF-O. These results further emphasized that artificial transformation was occurring during the action of pulling air through the sampling tube.

An inverse relationship was observed in both the laboratory and field study experiments between total chlorpyrifos expressed as a mass per unit volume (ng m^{–3}) and the proportion of chlorpyrifos transformed to oxon on the XAD-2 resin sampling matrix ([Fig. 1](#)). A quadratic prediction plot model accounted for about 29% of the variability in these data ($R^2 = 0.288$). The 95% confidence intervals were generated from the model to assess differences between the transformation experiments and the 2008 Washington State study. In many instances the proportion of oxon found in the 2008 air monitoring samples exceeded the upper 95% C.I. of the experimental data. This was especially true at low concentrations ($\leq 10 \text{ ng m}^{-3}$) and in ambient samples that were located > 500 m from the nearest orchard. In fact, more than half (69%) of the ambient samples ($n = 32$) were higher than the predicted artificial transformation values. These samples warrant further investigation because this means significant airborne CPF-O in residential air. On the other hand, some of the percent oxon was less than expected at

Table 2

Spike masses (ng), air volume (m^3), recovery masses total chlorpyrifos (CPF), chlorpyrifos-oxon (CPF-O), and total mixture (CPF + CPF-O) (ng), % recovery total (CPF + CPF-O), and % oxon (CPF-O) in laboratory and field studies measured after 24 h air sampling with chlorpyrifos-spiked OVS tubes. All reported values are the mean of spiked samples run in triplicates. *Italicized* recovery masses are corrected by subtracting background levels CPF and CPF-O in the outdoor environment from non-spiked samples. These corrected values were used to calculate total % recovery and % CPF-O.

Spike mass (ng)	Sample air volume (m^3)	Recovery mass (ng)			% Recovery	% CPF-O ^b
		CPF _m	CPF-O _m	CPF _{Total} ^a (CPF + CPF-O)	Total	
Laboratory						
Low flow rate (2 LPM)						
0	2.81	<LOD	<LOD	<LOD	<LOD	NA
42	2.91	22	9.7	32.1	76.5	31.6
210	2.92	128	37.7	167.5	79.8	25.5
High flow rate (6 LPM)						
2100	8.25	1437	256.0	1706	81.2	15.4
Field^c						
No active flow rate (0 LPM) ^c						
0	0	<LOD	<LOD	<LOD	<LOD	NA
30	0	25	0	25	83.3	NA
Low flow rate (2 LPM)						
0 ^d	2.78	16.5	6.5	23.3	>100	30.5
15	3.03	7.5	2.0	11.4	76.0	18.4
30 ^d	2.88	19.5	0.5	20.6	68.7	2.5
60	2.86	52.0	3.5	55.5	92.5	6.3
200	3.2	161.5	12.5	169.7	84.9	7.3
High flow rate (6 LPM)						
0	10.1	48.5	19.5	70.96	>100	29.5
200	10.21	150.5	27.0	177.4	88.8	16.0
592	9.36	482	44.5	512.4	86.6	9.1
2628	9.58	2118	151	2228	84.8	7.1

^a Total chlorpyrifos mass (CPF_{Total}, ng) is the sum of CPF and CPF-O measured in samples, adjusted by molecular weight ratio (see Equation (1) in text).

^b % CPF-O is the amount of CPF-O divided the amount of total CPF times 100 (see Equation (2) in text).

^c These samples had no recorded flow rate because they were deployed on a sampling tube with no sampling pump attached.

^d Duplicate sample resulting from pump failure.

medium concentrations and in near-field samples <100 m of an orchard. Only 36% of near-field samples ($n = 113$) had higher percentages chlorpyrifos-oxon than predicted. Higher proportions CPF-O in the air samples farther in proximity from fields may be attributable to physiochemical transformation as pesticides drift further from their source of application. In the past, atmospheric scientists have used the term “aging” to describe similar processes on the local/regional scale for complex mixes of pollutants and transformation products (Demerjian, 2011).




Fig. 1. % Artificial transformation to oxon by Log mass per unit volume of total chlorpyrifos.

All the community samples from the Washington State study, both ambient and near-field, had a median of 14% greater proportion oxon (95 C.I. 12.8–20.3%) than could be explained by the artificial transformation experiments in the laboratory and field. This difference was statistically significant in a Mann Whitney test ($\alpha < 0.005$). When broken into tertiles (Fig. 2), the Washington study had a 15% greater proportion of oxon than expected (95 C.I. 10.04–18.9%; $\alpha < 0.001$) at low concentrations ($<10 \text{ ng m}^{-3}$) and 10% greater proportion of oxon than expected (95 C.I. 4.8–16.97%; $\alpha = 0.005$) at medium concentrations ($10–30 \text{ ng m}^{-3}$). The opposite was true for higher concentrations ($>30 \text{ ng m}^{-3}$); the Washington study actually had 3% less proportion of oxon than expected, but this was not statistically significant (95 C.I. -7.85–12.7%; $\alpha = 0.401$).

4. Discussion

4.1. Artificial transformation

In this study we found that CPF is artificially transformed on XAD-2 resin to CPF-O, but that it does not account for all CPF-O measured in the air of an agricultural community. This transformation is a major concern because Agency for Toxic Substances and Disease Registry (ATSDR) currently recommends the use of XAD-2 resin air sampling matrices. ATSDR notes that CPF may be converted to its oxygen analog under certain environmental conditions; however, it does not discuss the potential for artificial production of the oxon on the sampling matrix (ASTM, 2011). Neither the EPA/ASTM D 4861 international method nor the NIOSH 5600 method makes any mention of measuring for CPF-O. This study has demonstrated that if the oxon in air samples using XAD-2 resins is not measured, then true CPF air concentrations maybe underestimated by 5–30%.

Woodrow et al. (1978) noted the conversion of parathion to paraoxon on Amberlite XAD-4 resins during 1-h laboratory testing at high air sampling volumes and Seiber et al. (1989) also reported substantial conversion (up to 54%) of methyl parathion to its oxygen analog on XAD. In 1996, the California Air Resources Board (CARB) found substantial amounts of CPF-O in many of the samples collected in Tulare County (1998b, 1998c). CARB reported that when using XAD-4 resin, “conversion of chlorpyrifos to the oxon analog may take place on the trapping media during sampling.” However, CARB also reported that concurrent field spike studies showed “only insignificant conversion taking place under actual field conditions.” These results are puzzling in light of our finding of significant conversion of CPF to its oxon on XAD-2 resin in both the laboratory and in the field.

The act of drawing air through the sampling resin may be a causal factor, as demonstrated by no conversion in XAD resin at zero flow rate (Table 2). In the future, it will be important to test new sampling matrices or examine passive methods using diffusion or deposition to measure CPF and CPF-O in air. Arcury et al. (2006) have noted that many researchers and safety personnel continue to collect airborne OP pesticide samples without any harmonized environmental sampling and chemical analysis methods. Not only do problems exist with the current method, there is also a need for new guidelines on how to collect and measure both organophosphate pesticides and their oxygen analogs with minimal sampling artifacts.

4.2. Environmental chlorpyrifos-oxon (CPF-O)

A direct comparison to the 2008 Washington State study has also demonstrated that transformation of OP pesticides to their oxygen analogs may still be occurring in the atmospheric




Fig. 2. Mann Whitney U-test comparing observed vs. expected % chlorpyrifos-oxon at concentration tertiles; *p value ≤ 0.05 .

environment. Under these circumstances, the health risk associated with exposure to airborne OP pesticides will be higher than previously estimated due to increased potency of CPF-O in the airborne mixture.

We have demonstrated this phenomenon by using toxicity equivalence factor (TEF) values, which have been used in the past for compounds such as polychlorinated biphenyls (PCBs) and dioxins (Van den Berg et al., 1998) in chemical risk assessment. These numbers are drawn by two main *in vivo* toxicology studies that conducted side-by-side comparisons of brain acetylcholinesterase inhibition for both CPF and CPF-O (Chambers and Carr, 1993; Cole et al., 2005).

A toxicity equivalent concentration (TEC) can be calculated for the mixture of CPF and CPF-O by adjusting for the toxicity of the oxon formed in the environment and available for inhalation:

$$TEC_{CPF} = (CPF-O_e \cdot CPF_{MW}/CPF-O_{MW} \cdot TEF) + CPF \quad (3)$$

Where TEC = toxicity equivalent concentration of chlorpyrifos parent and oxon; CPF-O_e = mass of environmental chlorpyrifos-oxon; CPF_{MW} = molecular weight of chlorpyrifos ($350.6 \text{ ng nmol}^{-1}$); CPF-O_{MW} = molecular weight of chlorpyrifos-oxon ($334.5 \text{ ng nmol}^{-1}$); TEF = toxicity equivalence factor drawn from toxicology studies; and CPF = chlorpyrifos measured in the sample (ng).

In Table 3, we illustrate the possible magnitude of changes in toxicity from respiratory exposure to mixtures of the parent CPF and CPF-O. We conclude that even at low concentrations CPF-O (e.g., 2 ng m^{-3}), total toxicity can be increased by factors of one

to ten. If the higher toxicity equivalence factor values are correct, then toxicity could increase by more than 20-fold, putting some deceptively "low" concentrations above the acute screening level of 1200 ng m^{-3} as defined by the California EPA (CDPR, 2006).

Since many families in central Washington State live in communities with high agricultural density, we have found that pesticide levels with high amounts CPF-O ($>30 \text{ ng}$) would be a concern to human health. Neither artificial transformation nor toxicity equivalence of CPF-O was taken into account for the 2008 air monitoring study. Given its high potency, the failure to include analysis for CPF-O in air samples could lead to considerable underestimates of health risk, especially when considering populations of genetically susceptible individuals and young children.

Future research should continue to investigate the sources and causes of oxon formation in dusts and airborne particles, such as "aging", mixtures, photodegradation, temperature, and the presence of oxidizing compounds such as ozone (O_3) and sulfur dioxide (SO_2). Ozone levels in particular have been predicted to increase substantially in areas like central Washington State as a result of changes in the atmosphere and climate. Although ozone is known to damage vegetation and reduce crop yields, less is known about its potential airborne chemical interactions with pesticides and may become an increasing concern with climate change patterns.

These results will contribute to the much needed explanation of environmental fate and transport of CPF-O to complement recent toxicological and epidemiological studies that have examined numerous health effects. These experiments stress the importance of accurately defining oxon presence in the environment because low concentrations may change health risk assessments—especially at concentrations typical for residential atmospheres farther from agricultural fields, where many families and children live, work and play. These findings call for consideration of the presence of organophosphorus oxons in air when calculating residential risk to pesticide drift and aerial transport.

Table 3

Toxicity equivalent concentrations of chlorpyrifos for selected air concentrations after incorporation of chlorpyrifos-oxon using three toxicity equivalent factors. *Italics* are $>$ California EPA Screening Level of 1200 ng m^{-3} (CDPR, 2006).

Measured air concentration CPF (ng m ⁻³) ^a	Environmental oxon CPF-O (ng m ⁻³) ^a	Toxicity equivalent concentration ^b (ng m ⁻³)		
		5 × TEF ^c	10 × TEF ^c	100 × TEF ^c
15	2	25	36	225
25	2	35	46	235
50	11	108	165	1203
150	32	318	485	3500
250	25	381	512	2870

^a Corrected for artificial transformation.

^b $TEC_{CPF} = (CPF-O_e \cdot CPF_{MW}/CPF-O_{MW} \cdot TEF) + CPF$ (see Equation (3) in text).

^c TEF = toxicity equivalence factor; 5-fold value from Chambers and Carr (1993); 10 and 100-fold values from Cole et al. (2005), Sultatos et al. (1982).

Acknowledgments

This work was supported by the NIOSH Educational and Research Center Grant NIOSH-T42OH00843 and the Child Health Center Pesticide Exposures Pathways Project at the Center for Child Environmental Risks Research NEIHS-P01 ES009601, EPA-RD-83451401. Thanks to Lucio Costa, PhD for consultation on toxicity estimates.

References

Arcury, T., Quandt, S., Barr, D., Hoppin, J., McCauley, L., Grzywacz, J., Robson, M., 2006. Farmworker exposure to pesticides: methodologic issues for the collection of comparable data. *Environ. Health Perspect.* 114, 923–928.

ASTM, 2011. Standard Practice for Sampling and Selection of Analytical Techniques for Pesticides and Polychlorinated Biphenyls in Air. ASTM D4861-11. 2011 Annual Book of ASTM Standards. In: Atmospheric Analysis, vol. 11.07 (ASTM formerly known as American Society for Testing and Materials).

Aston, L., Seiber, J., 1997. Fate of summertime airborne organophosphate pesticide residues in the Sierra Nevada Mountains. *J. Environ. Qual.* 26, 1483–1492.

Bavcon Kralj, M., Franko, M., Trebs, P., 2007. Photodegradation of organophosphorus insecticides – investigations of products and their toxicity using gas chromatography–mass spectrometry and AChE-thermal lens spectrometric bioassay. *Chemosphere* 67, 99–107.

CARB, April 7 1998a. Report for the Application and Ambient Air Monitoring of Chlorpyrifos (and Oxon Analogue) in Tulare County during Spring/Summer 1996. California Air Resources Board, Sacramento, CA.

CARB, April 7 1998b. Appendices for the Report for the Application and Ambient Air Monitoring of Chlorpyrifos (and the Oxon Analogue) in Tulare County during Spring/Summer, 1996. California Air Resources Board, Sacramento, CA.

CARB, 1998c. Report for the Application (Kings County) and Ambient (Fresno County) Air Monitoring of Diazinon during Winter, 1998. C97-070/C97-069. California Air Resources Board, Sacramento, CA.

CDPR, March 2003. Report of Ambient Air Monitoring for Pesticides in Lompoc, California. Department of Pesticide Regulation, Sacramento, CA.

CDPR, December 2006. Environmental Justice Pilot Project: Pesticide Air Monitoring in Parlier, Second Progress Report. Attachment IV discusses screening levels. California Environmental Protection Agency, Department of Pesticide Regulation. http://www.cdpr.ca.gov/docs/envjust/pilot_proj/index.htm.

CDPR, December 2009. Pesticide Air Monitoring in Parlier, CA (Final Report). California Department of Pesticide Regulation, Sacramento, CA.

Chambers, J., Carr, R.L., 1993. Inhibition patterns of brain acetylcholinesterase and hepatic and plasma esterases following exposures to three phosphorothionate insecticides and their oxons in rats. *Toxicol. Sci.* 21, 111–119.

Cole, T., Walter, B., Shih, D., Tward, A., Lusis, A., Timchalk, C., Richter, R., Costa, L., Furlong, C., 2005. Toxicity of chlorpyrifos and chlorpyrifos oxon in a transgenic mouse model of the human paraoxonase (PON1) Q192 polymorphism. *Pharmacogenet. Genom.* 15, 589–598.

Cole, T.B., Beyer, R.P., Bammler, T.K., Park, S.S., Farin, F.M., Costa, L.G., Furlong, C.E., 2011. Repeated developmental exposure of mice to chlorpyrifos oxon is associated with paraoxonase 1 (PON1)-modulated effects on cerebellar gene expression. *Toxicol. Sci.* 123, 155–169.

Costa, L., Cole, T., Vitalone, A., Furlong, C., 2005. Measurement of paraoxonase 1 (PON1) status as a potential biomarker for organophosphate toxicity. *Clin. Chim. Acta* 352 (1–2), 37–47.

Curwin, B.D., Hein, M.J., Sanderson, W.T., Nishioka, M.G., Reynolds, S.J., Ward, E.M., Alavanja, M.C., 2005. Pesticide contamination inside farm and nonfarm homes. *J. Occup. Environ. Hyg.* 2, 357–367.

Demerjian, K., 2011. "Atmospheric Science of Air Pollution Phenomena—Current Directions toward Exposure Characterization", Technical Challenges of Multi-pollutant Air Quality Management. Springer Science + Business Media, pp. 231–259.

Farahat, F.M., Fenske, R.A., Olson, J.R., Galvin, K., Bonner, M.R., Rohlman, D.S., Farahat, T.M., Lein, P.J., Anger, W.K., 2010. Chlorpyrifos exposures in Egyptian cotton field workers. *Neurotoxicology* 31, 297–304.

Fay, M., Proschak, M., 2010. Wilcoxon–Mann–Whitney or t-test? On assumptions for hypothesis tests and multiple interpretations of decision rules. *Stat. Surv.* 4, 1–39.

Fenske, R.A., Yost, M., Galvin, K., Tchong, M., Negrete, M., Palmendez, P., Fitzpatrick, C., 2009. Organophosphorus Pesticide Air Monitoring Project, Final Report, University of Washington; Available from the Washington State Department of Health Pesticide Program. At: <http://www.doh.wa.gov/ehp/pest/uwdrift-report.pdf>.

Furlong, C., Cole, T., Jarvik, G., Pettan-Brewer, C., Geiss, G., Richter, R., Shih, D., Tward, A., Lusis, A., Costa, L., 2005. Role of paraoxonase (PON1) status in pesticide sensitivity: genetic and temporal determinants. *Neurotoxicology* 26, 651–659.

Harnly, M., McLaughlin, R., Bradman, A., Anderson, M., Gunier, R., 2005. Correlating agricultural use of organophosphates with outdoor air concentrations: a particular concern for children. *Environ. Health Perspect.* 113, 1184–1189.

Huff, R., Corcoran, J., Anderson, M., Abou-Donia, M., 1994. Chlorpyrifos oxon binds directly to muscarinic receptors and inhibits cAMP accumulation in rat striatum. *J. Pharmacol. Exp. Ther.* 269, 329–335.

Jaipieam, S., Visuthismajarn, P., Siriwong, W., Borjan, M., Robson, M.G., 2010. Inhalation exposure of organophosphate pesticides by vegetable growers in the Bang-Rieng subdistrict in Thailand. *J. Environ. Public Health* (published online February 7).

LeNoir, J., McConnell, L., Fellers, G., Cahill, T., Seiber, J., 1999. Summertime transport of current-use pesticides from California's Central Valley to the Sierra Nevada mountain range, USA. *Environ. Toxicol. Chem.* 18 (12), 2715–2722.

NIOSH, 1994. NIOSH Manual of Analytical Methods. Pub. No. 94-113, fourth ed. National Institute for Occupational Safety and Health Cincinnati, OH.

Peck, A., Hornbuckle, K., 2005. Gas-phase concentrations of current-use pesticides in Iowa. *Environ. Sci. Technol.* 39 (9), 2952–2959.

Popendorf, W., Leffingwell, T., 1978. Natural variations in the decay and oxidation of parathion foliar residues. *J. Agric. Food Chem.* 26, 437.

Rosner, B., Grove, D., 1999. Use of the Mann–Whitney U-test for clustered data. *Stat. Med.* 18, 1387–1400.

Sancho, J., Pozo, O., Hernández, F., 2000. Direct determination of chlorpyrifos and its main metabolite in human serum and urine by coupled column liquid chromatography/electrospray-tandem mass spectrometry. *Rapid Commun. Mass Spectrom.* 14, 1485–1490.

Seiber, J., McChesney, M., Woodrow, J., 1989. Airborne residues resulting from use of methyl parathion, molinate, and thiobencarb on rice in the Sacramento Valley, California. *Environ. Toxicol. Chem.* 8, 577–588.

Shibamoto, T., Mourer, C., Hall, G., Hengel, M., 1996. Method Development, Ambient Site and Application Site Monitoring for Chlorpyrifos and Chlorpyrifos Oxon in Air Samples Using XAD-4 Resin as Trapping Medium. California Air Resources Board.

Spear, R., Lee, Y., Leffingwell, J., Jenkins, D., 1978. Conversion of parathion to paraoxon in foliar residues: effects of dust level and ozone concentration. *J. Agric. Food Chem.* 26, 434.

Sultatos, L., Costa, L., Murphy, S., 1982. Factors involved in the differential acute toxicity of the insecticides chlorpyrifos and methyl chlorpyrifos in mice. *Toxicol. Appl. Pharmacol.* 65, 144–152.

Thomas, K.W., Dosemeci, M., Hoppin, J.A., Sheldon, L.S., Croghan, C.W., Gordon, S.M., Jones, M.L., Reynolds, S.J., Raymer, J.H., Akland, G.G., Lynch, C.F., Knott, C.E., Sandler, D.P., Blair, A.E., Alavanja, M.C., 2010. Urinary biomarker, dermal, and air measurement results for 2,4-D and chlorpyrifos farm applicators in the Agricultural Health Study. *J. Expo. Sci. Environ. Epidemiol.* 20, 119–134.

Timchalk, C., Busby, A., Campbell, J., Needham, L., Barr, D., 2007. Comparative pharmacokinetics of the organophosphorus insecticide chlorpyrifos and its major metabolites diethylphosphate, diethylthiophosphate and 3, 5, 6-trichloro-2-pyridinol in the rat. *Toxicology* 237, 145–157.

US CENSUS, 2010. United States Census Bureau, Washington, DC. <http://2010.census.gov/2010census/>.

USEPA, 1999. Method TO-10A: Determination of Pesticides and Polychlorinated Biphenyls in Ambient Air Using Low Volume Polyurethane Foam (PUF) Sampling Followed by Gas Chromatographic/Multi-Detector Detection (GC/MD). In: Compendium of Methods for the Determination of Toxic Organic Compounds in Ambient Air. United States Environmental Protection Agency. <http://www.epa.gov/ttnamti1/files/ambient/airtox/to-10ar.pdf> (accessed 07.09.11.).

Van den Berg, M., Birnbaum, L., Bosveld, A., Brunstrom, B., Cook, P., Feely, M., Giesy, J.P., Hanberg, A., Hasegawa, R., Kennedy, S.W., Tubiak, T., Larsen, J.C., van Leeuwen, F.X., Liem, A.K., Nolt, C., Peterson, R.E., Poellinger, L., Safe, S., Schrenk, D., Tillitt, D., Tyskild, M., Younes, M., Waern, F., Zacharewski, T., 1998. Toxic equivalency factors (TEFs) for PCBs, PCDDs, PCDFs for humans and wildlife. *Environ. Health Perspect.* 106 (12), 775–792.

Woodrow, J., Crosby, D., Mast, T., Moilanen, W., Seiber, J., 1978. Rates of transformation of trifluralin and parathion vapors in air. *J. Agric. Food Chem.* 26, 6.

Yusa, V., Coscolla, C., Mellouk, W., Pastor, A., de la Guardia, M., 2009. Sampling and analysis of pesticides in ambient air. *J. Chromatogr.* 1216, 2972–2983.