

## USING PCR TO DETECT INDOOR FUNGI

BT Chen<sup>1\*</sup>, J Keswani<sup>1</sup>, G Zhou<sup>2</sup>, and T Ong<sup>1</sup>

<sup>1</sup>Health Effects Laboratory Division, NIOSH, CDC, Morgantown, WV, USA

<sup>2</sup>Health Canada, Tunney's Pasture 0803A, Ottawa, Ontario K1A 0L2 Canada

### ABSTRACT

This research attempts to develop a molecular bioassay that uses a Polymerase Chain Reaction (PCR) technique for detecting fungi commonly found in an indoor environment. During sample preparation, the method of bead-beating was identified as the most effective way for spore breakage and fungal DNA release. In the process of assay development, a fungal primer set, FF2/FR1, was designed and tested with DNA from human, rats, mice, bacteria, pollens and six fungal species. The results based on the crude extracts indicated that the primer set successfully amplified the fungal DNA without any cross-amplification with non-fungal DNA. In addition, higher amplification efficiencies were achieved using 20% nutrient media, rather than water, as the process solution. This PCR assay has a sensitivity of detecting low levels of fungi (2 fungal spores per reaction).

### INDEX TERMS

Fungi, Indoor air quality, Polymerase chain reaction (PCR), Exposure assessment, Allergies and asthma

### INTRODUCTION

Biological hazards in an indoor environment are diverse. Reported problems have ranged from allergic and infectious diseases to nonspecific "sick building symptoms" such as headaches, fatigue, and chest tightness. According to the annual census of NIOSH health hazard evaluations, the proportion of workers' requests related to indoor air quality in non-industrial environments has increased over the years, from 2% in 1980 up to 65% in recent years, and the costs are estimated at tens of billions of dollars per year (NORA, 1996). The conventional approach for assessing worker exposure to bioaerosols is to take samples and then perform laboratory analyses, which could be directly counting the organisms, indirectly counting colony-forming units, or using an immunoassay to detect the presence of specific biological agents. Although these methods may provide reasonably adequate assessment of bioaerosol concentration, the process of assessing exposure has been hampered by the lack of measurement techniques with near real-time, on-site, and high-sensitivity features. The present study is to establish and examine the applicability of a convenient, rapid, sensitive, and fungi-specific molecular detection system that uses polymerase chain reaction (PCR) to amplify DNA from the common indoor fungi without any cross-amplification with DNA from non-fungal organisms. This system would be useful in quickly screening low levels of airborne fungi in an environment and allow industrial hygienists or health professionals to rapidly determine the seriousness of fungal contamination and make investigation plans. In addition, this method has the advantage of monitoring for culturable fungal spores as well as non-culturable fungal substances that may not be viable but could be toxic or allergic.

---

\* Contact author email: [bdc4@cdc.gov](mailto:bdc4@cdc.gov)

## MATERIALS AND METHODS

### PCR

PCR is a rapid and simple technique for specific amplification of a particular segment of DNA. Normally the amplification occurs as long as the segment is present in the sample regardless of the culturability or viability of the microorganisms. As in other cells, ribosomal RNA (rRNA) genes in a microbial cell are both highly conserved and present in multiple copies, thus making them popular targets for gene amplification and molecular analysis (Lee and Taylor, 1990; Einsele, Hebart, Roller, *et al.*, 1997). PCR techniques based on rRNA genes have been widely used in developing sensitive systems for the detection of microorganisms (White, Bruns, Lee, *et al.*, 1990).

### Test fungi selection and PCR primer design

Six fungal species, including *Alternaria chlamydospora*, *Aspergillus flavus*, *Candida famata*, *Cladosporium fermentans*, *Penicillium chrysogenum*, and *Stachybotrys chartarum*, were selected to represent the molds (except *Ca. famata*) commonly found in a problematic indoor environment (Godish, 1995). A PCR method without any cross-amplification with DNA from other common indoor organisms (i.e., bacteria, pollen, mite, algae, rat, mouse and human) was developed. The fungal forward-primer FF2 (5'-GGTTCATTTTTGTTGGTTTCTA-3'), and reverse-primer FR1 (5'-CTCTCAATCTGTCAATCCTTATT-3') were designed from the 18S rRNA gene by selecting sequences homologous among the fungi, but not homologous with the other species. Although mite and algae were examined in their sequences during the primer design, they were not included in the actual experimental tests, because their sequences contain >10 mismatched bases with the designed primers and were thought not to be conducive to cross-amplification. Since fungal genetic characteristics are closer to plants than to bacteria, six types of plant pollen from cottonwood, red oak, ragweed, sagebrush, bentgrass, and Bermuda grass were selected to test the specificity of the fungal primers and only *Escherichia coli* was selected to represent the prokaryote in this test.

### Spore breakage and DNA release

Fungal cultures were grown and spores were harvested. Several methods, including freeze-thaw, French press, freeze press, and bead-beating, were examined for the release of fungal DNA. The bead-beating method, by vigorously colliding cells with the 0.1-mm zirconia/silica beads, reached nearly 100% spore breakage (Biospec products, Bartlesville, OK) and, thus, was selected for breaking up fungal spores, bacteria, and pollens. Percentage of cell breakage was calculated by microscopically counting the cells with a haemocytometer before and after the treatment. A vortex was also used for bead-beating with a efficiency of 85-100%. Because it is smaller and lighter, it could be an alternative when using the assay in the field DNA from rat liver, BALB/c-3T3 cells (mouse) and human peripheral blood lymphocytes were extracted using the standard phenol/chloroform method (Sambrook, Fritsch, and Maniatis, 1989).

Fungal spores collected from the culture plates were counted under a microscope and then diluted serially from  $10^6$  to  $10^2$  spores/ml with 20% nutrient media (nutrient broth - beef extract, peptone, and salt). A 0.1 ml of sample was directly added into a 0.5 ml micro-tube containing approximately the same volume of beads and homogenized for 3 min, then heated in a boiling water bath for 5-10 min to inactivate released nucleases. The samples were cooled to room temperature and spun for 2-5 s in a micro-centrifuge to discard cell debris. The supernatants were directly used as templates for PCR.

Pollen and bacterial DNA were also released with a bead-beating technique using the same preparation steps as described above.

In parallel to the above study design, a protocol using a different preparation procedure (homogenization first rather than dilution first) and a different process solution (pure water rather than 20% nutrient media) were also followed to investigate the effects of PCR amplification efficiency on the procedure and the process solution.

### PCR amplification and gel electrophoresis


Various supernatants (2 µl each) or DNA (200 ng) from human, rats or BALB/c-3T3 cells were amplified with each primer (10 pmol), Taq DNA polymerase (1.25 U), dNTPs (200 µM), and MgCl<sub>2</sub> (1.5 mM) in 50 µl of reaction buffer using a Thermal Cycler (Model 480, Perkin-Elmer Cetus, Foster City, CA). The conditions for PCR were as follows: initial denaturation of DNA at 95°C for 3 min and then 35 cycles of three-step PCR amplifications consisting of denaturation at 94°C for 1 min, primer reannealing at 52°C for 1 min, and extension at 72°C for 2 min. The samples were subjected to an additional extension at 72°C for 10 min at the end of the amplification cycles. Ten microliters of PCR products, mixed with loading buffer, were loaded on a 2.0% agarose : Nusieve GTG agarose gel (1:1) and electrophoresed with 1x TAE buffer at 8 V/cm. The gels were stained with ethidium bromide (0.5 µg/ml) in TAE for 10 min and visualized under UV light using an Eagle Eye II image system and photographed.

To investigate effects of fungal mixtures on the assay performance, various fungal spores were mixed in equal ratios to give similar total spore count ( $7.5 \times 10^4$  spores/ml) as that of individual spores and bead-beaten to release mixtures of spore DNA equivalents. These samples were prepared, amplified, and analyzed as described above.

## RESULTS

### Fungal specificity of primer pairs

To determine whether cross-amplification would occur, fungal, plant, human, rat, mouse, and bacterial DNA were broadly tested using the FF2/FR1 primer set. Figure 1 shows that this primer pair could successfully amplify the DNA of all fungal samples and yield an amplicon of 425 bp without showing any cross-amplification with non-fungal DNA. The amplified products were directly sequenced and the results indicated that the fragments from fungal samples indeed matched fungal DNA sequences.



**Figure 1.** PCR products amplified by primers FF2/FR1 from various sources of DNA

### Effects of procedure in spore preparation

In this study, the dilution-first and homogenization-first procedures were compared using 20% nutrient media. The amplified bands were observed at as low as 2 spore DNA equivalents/reaction using the dilution-first procedure, while the homogenization-first procedure could show amplified bands even at 0.2 spore DNA equivalent/reaction with 20% nutrient media.

### Effects of process solution

The amplification efficiency in water or 20% nutrient media was measured using the homogenization-first procedure to determine whether PCR would be hindered when DNA from disrupted spores was amplified directly. After  $10^6$  fungal spores/ml were disrupted and diluted tenfold serially, PCR amplification was performed to determine the efficiency of the assay system. Positive PCR bands were shown clearly at  $2 \times 10^3$  to  $0.2$  fungal spore DNA/reaction (equivalent to  $10^6$  to  $10^2$  spores/ml in concentration) and the amplification efficiency increased with the concentration (Table 1), except for *Alternaria* and *Stachybotrys* in which the efficiencies decreased at concentrations higher than  $2 \times 10^3$  spore DNA/reaction. Overall, a higher amplification efficiency was observed when 20% nutrient media was used to replace water as the cell breakage and dilution solution for all samples.


**Table 1.** Intensity of PCR Amplification in Different Breakage Solutions

| Fungal spore                   | Spore DNA/reaction in medium |     |      |       |       | Spore DNA/reaction in water |    |     |      |      |
|--------------------------------|------------------------------|-----|------|-------|-------|-----------------------------|----|-----|------|------|
|                                | 0.2                          | 2   | 20   | 200   | 2000  | 0.2                         | 2  | 20  | 200  | 2000 |
| <i>Alternaria chamydopora</i>  | ++                           | +++ | ++++ | +++++ | +++   | +                           | ++ | +++ | ++++ | ++   |
| <i>Aspergillus flavus</i>      | +                            | ++  | +++  | ++++  | +++++ | -                           | +  | ++  | +++  | ++++ |
| <i>Candida famata</i>          | +                            | ++  | +++  | ++++  | +++++ | -                           | +  | ++  | +++  | ++++ |
| <i>Cladosporium fermentans</i> | +                            | ++  | +++  | ++++  | +++++ | -                           | +  | ++  | +++  | ++++ |
| <i>Penicillium chrysogenum</i> | +                            | ++  | +++  | ++++  | +++++ | -                           | +  | ++  | +++  | ++++ |
| <i>Stachybotrys chartarum</i>  | +                            | ++  | +++  | ++++  | +++   | -                           | +  | ++  | +++  | +++  |

-, no amplification; +, very weak signal; ++, weak signal; +++, moderate signal; +++++, strong signal; ++++++, very strong signal

### Effects of mixtures

PCR amplification from crude extract of mixtures of spores produced an amplification product of the expected size similar to the PCR product from individual spores (Figure 2). The PCR products were semi-quantitated by analysis of gel image using Image Quant 3.3 and molecular weight markers for calibration. The amplified products were in the range of 70.61 to 93.77 ng of DNA/band, with an average of  $81.90 \pm 7.27$  (s.d) corresponding to approximately 150 spore DNA equivalents/reaction in PCR assay. Although this is a preliminary study on 3 different fungi mixed in equal ratios, the results were reproducible with amplified products within the same range.



**Figure 2.** PCR products amplified by primers FF2/FR1 from various mixtures of spore DNA

## DISCUSSION

A microbiological assay system has been established in our laboratory, which has the potential to become a screening tool to detect total airborne fungi in the indoor environment. This system includes a bead-beating process for breaking up the spores in the sample and a fungi-specific primer pair (FF2/FR1) to amplify the fungal DNA retrieved from spore breakage.

For environmental monitoring, it is important to exclude false positives that could result from the lack of specificity of the assay system. Real-life samples from the air, water or soil usually contain some degree of biological contaminants associated with human, rat, mouse, mite, algae, pollen or bacterial cells which may have potential for cross-amplification with fungi when applying PCR analysis. Most PCR assay systems reported in the literature have focused primarily on clinically relevant fungi and, thus, considered mainly cross-amplification with human and bacterial DNA (Makimura, Murayama and Yamaguchi, 1994; Einsele, Hebart, Roller, *et al.*, 1997), as well as plant DNA, (Kappe, Fauser, Okeke, *et al.*, 1996; Kappe, Okeke, Fauser, *et al.*, 1998) but not other biological contaminants. Our previous study on examining other "universal" fungal primers (Zhou, Wong, Ong, *et al.*, 2000) demonstrated that the primer pair, FF2/FR1 had no cross-amplification with rat and mouse DNA, and has more than 10 mismatched bases with the sequences of mite and algae. In addition, this primer set may be a suitable fungi-specific primer set for screening fungal presence in a setting other than a problematic indoor workplace, e.g., a hospital environment as determined by either complete homology or 1 base mismatch with 18 medically relevant fungi.

In this study, the supernatants were directly used because it would make the assay protocol more convenient and practical if cell-disrupted crude extract from the samples could be used directly for PCR without DNA purification. However, fungi supernatant may contain contaminants that could affect PCR amplification. When the nutrient media was used as the processing solution, proteins or other substances in the nutrient media might have increased PCR efficiency by preventing released fungal DNA from nonspecifically sticking to the beads as well as combining with inhibition-causing substances to help stabilize the Taq DNA polymerase. Practically speaking, a procedure of serial dilution of the crude DNA release after breakage might be needed for samples of high fungal concentrations (e.g., in an agricultural or other industrial environments) to attenuate the inhibition effect. For the samples collected from a nonindustrial indoor environment, the spore counts without dilution would be similar to the situation used in the dilution-first procedures and the sensitivity of 2 fungal spores/reaction would be a more proper representative of this assay system. The value is similar to that reported by Haugland and Heckman (1999) in which they describe a TaqMan<sup>TM</sup> system able to measure as low as 2 conidia/reaction of *S. chartarum*.

The simple three-fungi mixture study on PCR assay showed amplification products within a close range. However, the study was performed for mixtures in equal ratios. It is well documented that PCR bias due to many factors exists (Farrelly, Rainey, and Stackebrandt, 1995; Suzuki and Giovannoni, 1996; Polz and Cavanaugh, 1998). These include genome size, rDNA operon numbers, PCR selection, PCR drift, template saturation, etc. For most fungi, the detailed study of the overall genome is not known. It will be important to explore PCR bias from mixed template PCR reactions to interpret PCR-based results in a quantitative measure.

In summary, the designed primer set, FF2/FR1, has been tested to be fungus-specific and this PCR assay has a sensitivity of detecting low levels (2 spores) of fungi in a reasonable time period (5-6 h). This technique may be useful in estimating fungal bio-mass in an

environmental sample in general and have particular value in investigations of indoor air quality where exposure levels are low and adverse health effects are due to allergic and toxic reactions. It is, however, essential to test spiked environmental samples in determining the actual limits of detection as well as the effects of environmental inhibitors to the PCR prior to any actual field applications.

## ACKNOWLEDGMENTS

The authors are indebted to Drs. S. Soderholm and D. Lewis at NIOSH for their advice and review of this paper. The research is partly supported by the HUD Healthy Homes Program under Interagency Agreement IAA No. HUD I-OPC-21537.

## REFERENCES

- Einsele, H., Hebart, H., Roller, G. *et al.* 1997. Detection and identification of fungal pathogens in blood by using molecular probes. *J. Clin. Microbiol.* **35**, 1353-60.
- Farrelly, V, Rainey, F.A, and Stackebrandt, E. 1995. Effect of genome size and *rrn* gene copy number on PCR amplification of 16S rRNA genes from a mixture of bacterial species. *Applied and Environmental Microbiology*, **61**, 2798-2801.
- Godish, T. 1995. *Sick Building*. Pp. 171-203. Boca Raton: Lewis Publishers (CRC Press).
- Haugland, R.A. and Heckman, J. L. 1999. Quantitative measurement of *Stachybotrys chartarum* conidia using real time detection of PCR products with the TaqMan™ fluorogenic probe system. *Molecular and Cellular Probes* **13**, 329-40.
- Kappe, R., Fauser, C, Okeke, C. N., *et al.* 1996. Universal fungus-specific primer systems and group-specific hybridization oligonucleotides for 18S rRNA. *Mycoses* **39**, 25-30.
- Kappe, R., Okeke, C. N., Fauser, C., *et al.* 1998. Molecular probes for the detection of pathogenic fungi in the presence of human tissue. *J. Med. Microbiol.* **47**, 811-20.
- Lee, S. B. and Taylor, J. W. 1990. Isolation of DNA from fungal mycelia and single spores PCR protocols. In *PCR protocols: A guide to methods and applications* (Innis, M. A., Gefand, D. H., Sninsky, J. J. & White, T. J., eds.) Pp. 282-287. San Diego: Academic Press.
- Makimura, K., Murayama, S. Y. and Yamaguchi, H. 1994. Detection of a wide range of medically important fungi by the PCR. *J. Med. Microbiol.* **40**, 358-64.
- NORA. 1996. Indoor Environment, *National Occupational Research Agenda*, DHHS (NIOSH) Publication No. 96-115, pp 28-29.
- Polz, M.F and Cavanaugh, C.M 1998. Bias in template-to-product ratios in multitemplate PCR. *Applied and Environmental Microbiology*, **64**, 3724-3730.
- Sambrook, J., Fritsch, E. F. and Maniatis, T. 1989. *Molecular cloning: a laboratory manual*. Pp. 9.13-9.23. Cold Spring Harbor: Cold Spring Harbor Laboratory Press.
- Suzuki, M.T, and Giovannoni, S.J. 1996. Bias caused by template annealing in the amplification of mixtures of 16S rRNA genes by PCR. *Applied and Environmental Microbiology*, **62**, 625-630.
- White, T. J., Bruns, T., Lee, S., *et al.* 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In *PCR protocols: A guide to methods and applications* (Innis, M. A., Gefand, D. H., Sninsky, J. J. & White, T. J., eds.) Pp. 315-322. San Diego: Academic Press.
- Zhou, G, Wong, W. Z., Ong, T., *et al.* 2000. Development of a fungus-specific PCR assay for detecting low-level fungi in an indoor environment. *Molecular and Cellular Probes* **14**, 339-348.

# ***Indoor Air 2002***

Proceedings:  
9th International Conference on  
Indoor Air Quality and Climate

Monterey, California, June 30-July 5, 2002

Conference Proceedings Editor:  
*Hal Levin*

Editorial Assistants:  
*Gina Bendy, Joyce Cordell*

**Volumes I - V**

**ISBN 0-9721832-0-5**

published by  
Indoor Air 2002, 2548 Empire Grade, Santa Cruz, CA 95060  
email: [postmaster@indoorair2002.org](mailto:postmaster@indoorair2002.org)  
web: [www.indoorair2002.org](http://www.indoorair2002.org)