

Hyperosmolar Solution Effects in Guinea Pig Airways. IV. Lipopolysaccharide-Induced Alterations in Airway Reactivity and Epithelial Bioelectric Responses to Methacholine and Hyperosmolarity

Richard A. Johnston, Michael R. Van Scott, Choudari Kommineni, Lyndell L. Millecchia, Juanita Dortch-Carnes, and Jeffrey S. Fedan

Department of Pharmacology and Toxicology, Robert C. Byrd Health Sciences Center of West Virginia University, Morgantown, West Virginia (R.A.J., J.S.F.); Pathology and Physiology Research Branch, Health Effects Laboratory Division, National Institute for Occupational Safety and Health, Morgantown, West Virginia (R.A.J., C.K., L.L.M., J.D.-C., J.S.F.); and Department of Physiology, The Brody School of Medicine at East Carolina University, Greenville, North Carolina (M.R.V.S.)

Received March 14, 2003; accepted October 8, 2003

ABSTRACT

We investigated the *in vivo* and *in vitro* effects of lipopolysaccharide (LPS) treatment (4 mg/kg *i.p.*) on guinea pig airway smooth muscle reactivity and epithelial bioelectric responses to methacholine (MCh) and hyperosmolarity. Hyperosmolar challenge of the epithelium releases epithelium-derived relaxing factor (EpDRF). Using a two-chamber, whole body plethysmograph 18 h post-treatment, animals treated with LPS were hyporeactive to inhaled MCh aerosol. This could involve an increase in the release and/or actions of EpDRF, because LPS treatment enhanced EpDRF-induced smooth muscle relaxation *in vitro* in the isolated perfused trachea apparatus. In isolated perfused tracheas the basal transepithelial potential difference (V_t) was increased after LPS treatment. The increase in V_t was inhibited by amiloride and indomethacin. Concentration-re-

sponse curves for changes in V_t in response to serosally and mucosally applied MCh were biphasic (hyperpolarization, $<3 \times 10^{-7}$ M; depolarization, $>3 \times 10^{-7}$ M); MCh was more potent when applied serosally. The hyperpolarization response to MCh, but not the depolarization response, was potentiated after LPS treatment. In both treatment groups, mucosally applied hyperosmolar solution (using added NaCl) depolarized the epithelium; this response was greater in tracheas from LPS-treated animals. The results of this study indicate that airway hyporeactivity *in vivo* after LPS treatment is accompanied by an increase in the release and/or actions of EpDRF *in vitro*. These changes may involve LPS-induced bioelectric alterations in the epithelium.

Asthma is a chronic disease that is characterized by reversible airway obstruction, airway inflammation, and airway hyperresponsiveness. Lipopolysaccharide (LPS) or endotoxin, the major component of the outer membrane of gram negative bacteria, poses problems for asthmatic patients. For example, inhaled endotoxin causes a slight decrease in the forced expiratory volume in 1 s and an increase in histamine

responsiveness in asthmatic patients, but not in normal subjects (Michel et al., 1989, 1996). Several reports have demonstrated that LPS administration leads to the development of airway hyperreactivity and inflammation in mice (Held and Uhlig, 2000), rats (Pauwels et al., 1990), and guinea pigs (Toward and Broadley, 2000).

Airway reactivity is under the control of numerous physiological mechanisms, including epithelial-derived excitatory and inhibitory substances (Fedan et al., 1988; Goldie and Hay, 1997; Folkerts and Nijkamp, 1998). One such inhibitory substance that has been shown to modulate airway reactivity is the non-nitric oxide nonprostanoid epithelium-derived relaxing factor (EpDRF), which is released in response to hy-

This work was supported, in part, by National Institutes of Health Grant 5-T32-GM07039 (to R.A.J.). Mention of brand name does not constitute product endorsement. This article is the fourth one of a series of four companion articles that report the effects of hyperosmolar solutions in guinea pig airways (Fedan et al., 2003a,b; Wu et al., 2003).

Article, publication date, and citation information can be found at <http://jpet.aspetjournals.org>.
DOI: 10.1124/jpet.103.051672.

ABBREVIATIONS: LPS, lipopolysaccharide; EpDRF, epithelium-derived relaxing factor; MCh, methacholine; SRaw, specific airway resistance; V_t , transepithelial potential difference; L-NAME, *N*^G-nitro-L-arginine methyl ester; D-M, D-mannitol; NO, nitric oxide; MKHS, modified Krebs-Henseleit solution; CI, confidence interval; SNP, sodium nitroprusside; PC₂₀₀, the provocative methacholine concentration producing a 2-fold increase in specific airway resistance above the value after saline administration.

perosmolarity¹ at the mucosal or serosal surface of airway epithelial cells (Munakata et al., 1988, 1990; Fedan et al., 1999, 2003a). EpDRF release during hyperosmolar challenge occurs in response to the incremental increase in osmolarity (Fedan et al., 2003a) and is associated with epithelial bioelectric events (Dortch-Carnes et al., 1999; Wu et al., 2003). Evidence has been obtained to suggest that hyperosmolar challenge of the epithelium releases carbon monoxide (Fedan et al., 2003b). Once released, EpDRF initiates airway smooth muscle relaxation via an unknown mechanism.

There have been several reports indicating the importance of EpDRF in modulating airway reactivity. For example, in the guinea pig isolated perfused trachea, reactivity to methacholine (MCh) is decreased substantially when the release of EpDRF is stimulated by challenging the epithelium with hyperosmolar solution (Fedan et al., 1999). In addition, EpDRF-induced smooth muscle relaxation is significantly attenuated in isolated perfused tracheas from guinea pigs exposed to ozone, which is accompanied by *in vitro* and *in vivo* hyperreactivity to MCh (Fedan et al., 2000). Finally, preliminary evidence indicates that sensitization and challenge of guinea pigs with ovalbumin enhances the release of EpDRF and causes *in vitro* airway hyporeactivity to MCh, at a time when the animals exhibit hyperreactivity to inhaled MCh (Warner et al., 1996).

In addition, there is evidence that indicates that the synthesis, release, and/or effects of EpDRF are functionally linked to the electrical activity of epithelial cells. Dortch-Carnes et al. (1999) demonstrated that a decrease in the transepithelial potential difference (V_t) preceded EpDRF-induced smooth muscle relaxation elicited by elevating serosal or mucosal osmolarity with either ionic or nonionic osmolytes. Fedan et al. (1999) demonstrated that amiloride-sensitive Na^+ channels and 4,4'-diisothiocyano-2,2'-stilbene disulfonate-sensitive Cl^- channels are involved in EpDRF-induced smooth muscle relaxation, and Tamaoki et al. (1997) published evidence for involvement of Ca^{2+} -activated K^+ channels in the synthesis and/or release of EpDRF in human bronchial strips.

LPS has been shown to alter airway reactivity in human and animal subjects. Therefore, the purpose of this study was to examine the possible relationship between alterations in airway reactivity induced by LPS and alterations in the smooth muscle effects of EpDRF and the epithelial bioelectric events associated with its release.

Materials and Methods

Animals. These studies were conducted in facilities accredited fully by the Association for the Assessment and Accreditation of Laboratory Animal Care International and were approved by the institutional Animal Care and Use Committee. The animals were anesthetized with sodium pentobarbital (65 mg/kg *i.p.*) and sacrificed by thoracotomy and bleeding before removing the trachea. Other details of animal use have been given (Fedan et al., 2003a).

In Vivo Saline and LPS Treatment. Guinea pigs were injected *i.p.* with 4 mg/kg LPS from *Salmonella typhimurium* or a volume

equivalent of saline (control). Eighteen hours post-treatment, *in vivo* or *in vitro* experiments were conducted.

Two-Chamber, Whole-Body Plethysmograph. This method has been described in detail previously (Lawrence et al., 1997). Briefly, a two-chamber, whole-body plethysmograph (Buxco Electronics, Inc., Sharon, CT) was used to measure specific airway resistance (S_{Raw}) before and after delivery of MCh aerosol to measure airway reactivity in saline- and LPS-treated animals. Each chamber of the plethysmograph was connected to a differential pressure transducer and then to a noninvasive LS-20 airway mechanics analyzer (Buxco Electronics, Inc.) connected to a computer for logging acquired data at 6-s intervals.

The animal was acclimated in the plethysmograph while breathing air for 2 h on the day before the experiment. Eighteen hours after saline- or LPS treatment, the animal was placed in the plethysmograph and S_{Raw} measurements were recorded for 4 min after an initial 16-min acclimation period in air. Dose-response curves for responses to aerosolized MCh were then generated.

In Vivo MCh Dose-Response Curves. The animal was exposed to aerosolized saline as a vehicle control. The saline was delivered for 3 min, followed by a 10-min recording period. Dose-response curves were generated by administering MCh aerosol (Ultra Neb 99 nebulizer; Devilbiss Co., Somerset, PA) in half-log concentrations ranging from 0.03 to 3.16 mg/ml. Each MCh aerosol was delivered for 3 min, followed by a 10-min recording period. The peak S_{Raw} value was taken as the response to that particular concentration of MCh. Before the next higher dose of MCh was administered, S_{Raw} was allowed to return to baseline. Airway reactivity to MCh was determined by calculating the MCh PC₂₀₀, the provocative MCh concentration producing a 2-fold increase in S_{Raw} above the value after saline administration. The PC₂₀₀ was calculated by linear interpolation of graphed data.

In Vivo Administration of N^G-Nitro-L-arginine Methyl Ester (L-NAME). Guinea pigs were exposed to L-NAME aerosol (1.2 mM) for 5 min before the delivery of each MCh concentration. Two MCh dose-response curves were generated in both saline- and LPS-treated animals, the first in the absence and the second after treatment with L-NAME. This protocol is similar to that used by Nijkamp et al. (1993).

Histological Analysis of Guinea Pig Airways. After anesthesia and exsanguination, trachea and lungs from saline- and LPS-treated animals were inflated with 15 to 20 ml of 10% buffered formalin phosphate and removed en bloc. Sections of the trachea as well as the azygous, left, right apical, right cardiac, and right diaphragmatic lobes of the lung were then embedded with paraplast-plus paraffin before being cut into 5- μm slices with a microtome. Individual sections were stained according to Harris' hematoxylin and eosin procedure and prepared for reading. The slides were examined in a "blind" manner under the light microscope by a veterinary pathologist.

Modified Isolated Perfused Trachea Apparatus for Simultaneous Measurement of Transepithelial Potential Difference and Mechanical Responses. The isolated perfused trachea preparation was used to measure responses of the smooth muscle elicited by challenge of the epithelium with hyperosmolar solution, and other drug effects. This preparation permits separate application of agents to the mucosal (intraluminal) or serosal (extraluminal) surfaces of the trachea. The method has been described previously (Fedan and Frazer, 1992; Fedan et al., 2003a), along with modifications (Dortch-Carnes et al., 1999) for measuring V_t simultaneously with smooth muscle contractile and relaxant responses.

MCh and Elevated Osmolarity Concentration-Response Curves. MCh concentration-response curves were generated by adding MCh in stepwise-increasing, cumulative concentrations to the extraluminal and intraluminal baths. Both extraluminal and intraluminal MCh concentration-response curves were obtained from each trachea, the extraluminal curve first, followed 90 min later by the intraluminal curve, with washes every 15 min.

¹ Hypertonic solutions are those that cause cell shrinkage. Hyperosmolar solutions have osmolarity greater than that of the physiological extracellular solution. For simplicity, in this report we do not draw distinctions between the two terms when describing general phenomena.

Concentration-response curves for the effects of hyperosmolarity were generated by adding NaCl, D-mannitol (D-M), urea, or KCl in stepwise-increasing cumulative concentrations to the extraluminal and/or intraluminal baths in the presence or absence of extraluminal MCh (3×10^{-7} M), as described under *Results*.

Tracheal Epithelium Removal. In experiments that required removal of the tracheal epithelium, a trimmed 6-cm segment of pipe cleaner was slowly inserted into the tracheal lumen and then withdrawn while rotating slowly (Fedan and Frazer, 1992).

Inhibitors. L-NAME [nitric-oxide (NO) synthase inhibitor; 10^{-4} M] and indomethacin (cyclooxygenase inhibitor; 3×10^{-6} M) were used to determine whether hyperosmolarity-induced smooth muscle relaxation involved an NO or prostanoid component. The inhibitors were added to the extraluminal and intraluminal baths 30 min before inducing tone with extraluminally added 3×10^{-7} M MCh. The relaxation responses in the presence of the inhibitors were compared with relaxation responses in preparations where the inhibitor was absent.

Effect of Agents on the Basal V_t . The following agents were evaluated for their effects on epithelial bioelectric properties: indomethacin (3×10^{-6} M), L-NAME (10^{-4} M), and the epithelial Na^+ channel blocker amiloride (3×10^{-5} M). Indomethacin and L-NAME were added simultaneously to the extraluminal and intraluminal baths, whereas amiloride was only added to the intraluminal bath.

Solutions and Reagents. The modified Krebs-Henseleit solution (MKHS) contained 113.0 mM NaCl, 4.8 mM KCl, 2.5 mM CaCl_2 , 1.2 mM KH_2PO_4 , 1.2 mM MgSO_4 , 25.0 mM NaHCO_3 , and 5.7 mM glucose. The MKHS was continuously gassed with 95% O_2 , 5% CO_2 and maintained at 37°C and pH 7.4. The osmolarity of the MKHS was 281 ± 5 mosM.

All drugs, chemicals, and agents were from Sigma-Aldrich (St. Louis, MO) and dissolved in saline unless otherwise noted. Amiloride was dissolved in distilled water; indomethacin was dissolved in 100 mM sodium carbonate.

Statistical Analysis of Results. For in vivo airway reactivity studies, SRaw and MCh PC_{200} values among saline- and LPS-treated animals were compared using the paired or nonpaired Student's *t* test or the Mann-Whitney rank sum test, as appropriate. For in vitro airway reactivity studies, geometric EC_{50} values were derived from least-squares analysis of a sigmoidal curve fit. When comparing the EC_{50} values, normally distributed $-\log\text{EC}_{50}$ values were used. The EC_{50} values, maximum responses, and the intraluminal/extraluminal maximum response ratios were analyzed using the paired or nonpaired Student's *t* test or one-way repeated measures analysis of variance, as appropriate. When examining the effect of agents on the basal V_t , the data were analyzed statistically using either the paired or nonpaired Student's *t* test or the Mann-Whitney rank sum test, as appropriate. The results, except where noted, were expressed as the mean \pm S.E.; *n* is the number of separate experiments. $p < 0.05$ was considered significant.

Results

Effect of Saline and LPS Treatments on Basal Pulmonary Function and in Vivo Airway Reactivity to MCh. Basal respiratory rate was slightly, but significantly, decreased by saline treatment (data not shown). However, there were no other differences between the pre- or post-treatment breathing frequency and SRaw values within each treatment group. LPS treatment induced a rightward shift in the MCh dose-response curve (Fig. 1). The MCh PC_{200} (milligrams of MCh per milliliter) values [95% confidence interval (CI) in parentheses] were saline, 0.08 (0.05–0.14); and LPS-treated, 0.26 (0.08–0.79) ($p < 0.05$).

Effect of L-NAME on in Vivo Airway Reactivity to MCh in Saline- and LPS-Treated Animals. Because LPS




Fig. 1. Effect of LPS treatment on in vivo airway reactivity to inhaled MCh aerosol. Reactivity to MCh was measured 18 h after saline or LPS treatment. Saline- and LPS-treated, $n = 6$.

is known to induce NO synthase activity in the lung (Salter et al., 1991) and because NO is capable of modulating airway smooth muscle reactivity to agonists (Fedan et al., 1995; Folkerts and Nijkamp, 1998), it was hypothesized that NO could account for the in vivo airway hyporeactivity to MCh after LPS treatment. Thus, the effect of an aerosol of L-NAME, an NO synthase inhibitor, on in vivo airway reactivity to MCh was determined (Fig. 2). MCh dose-response curves obtained in the absence of L-NAME indicated that the LPS-treated animals were hyporeactive to MCh (Table 1). Administration of L-NAME before MCh had no effect upon airway reactivity in either the saline- or LPS-treated animals (Table 1).

Histological Examination of Guinea Pig Airways after LPS Treatment. Because L-NAME failed to provide any clues to the mechanisms underlying LPS-induced in vivo airway hyporeactivity to MCh, we examined the effects of LPS treatment on the structure of various regions of the guinea pig respiratory tract. This was done to determine whether LPS treatment altered the structure of the airways in such a way as to limit access of MCh to the airway smooth muscle (i.e., excessive mucus secretion, epithelial and basement membrane hyperplasia). Histological sections of the distal trachea as well as the azygous, left, right, apical, right cardiac, and right diaphragmatic lobes of the lung from saline- and LPS-treated animals were examined. The sections




Fig. 2. Lack of effect of L-NAME on in vivo airway reactivity to inhaled MCh aerosol in saline- (A) and LPS-treated (B) animals. Each animal served as its own control, and reactivity to MCh was measured 18 h after saline or LPS treatment, before and after delivery of aerosolized L-NAME (1.2 mM). Saline- and LPS-treated, $n = 4$.

TABLE 1
Effect of L-NAME on airway reactivity to inhaled MCh in saline- and LPS-treated guinea pigs

Treatment (n)	MCh PC ₂₀₀ (mg/ml) (95% CI)	
	-L-NAME	+L-NAME
Saline (4)	0.07 (0.06-0.10)	0.01 (0.00-0.94)
LPS (4)	0.78 (0.31-0.97) ^a	0.40 (0.11-1.45)

^a Significantly greater than saline (-L-NAME).

from both treatment groups were indistinguishable from one another, at least at the level of the light microscope. Representative sections of the distal trachea and left lobe of the lung from saline- and LPS-treated animals are shown in Fig. 3. The bronchial and tracheal epithelium as well as the alveolar sacs remained structurally intact within each treatment group. Finally, in both treatment groups, a few inflammatory cells, namely, eosinophils and neutrophils, could be seen infiltrating the airways in some regions of the respiratory tract that were examined (data not shown).

Effect of LPS Treatment on in Vitro Reactivity to MCh in Epithelium-Intact and -Denuded Isolated Perfused Trachea. Because LPS treatment caused in vivo airway hyporeactivity to inhaled MCh, we investigated whether this effect manifested itself in vitro in epithelium-intact and -denuded isolated perfused trachea.

In epithelium-intact tracheas, there was no difference in reactivity to extraluminally and intraluminally applied MCh

between the two treatment groups. The tracheas from both treatment groups were more sensitive to extraluminally applied MCh (Fig. 4A; Table 2). In addition, there was no difference in the intraluminal/extraluminal maximum response ratio (Fedan and Frazer, 1992) between the saline- and LPS-treated groups.

In epithelium-denuded tracheas, the differences between extraluminal and intraluminal reactivity to MCh were abolished, and the intraluminal/extraluminal maximum response ratio approached unity (Fig. 4B; Table 3). In addition, there was no difference in reactivity to extraluminally and intraluminally applied MCh in epithelium-denuded tracheas from saline- and LPS-treated animals.

Effect of Hyperosmolar Solution-Induced Smooth Muscle Relaxation in Isolated Perfused Trachea. Because LPS treatment caused in vivo airway hyporeactivity to MCh, it was hypothesized that EpDRF-mediated relaxation responses to hyperosmolar challenge would be potentiated. Intraluminal osmolarity was increased using NaCl, D-M, urea, or KCl as osmolytes after contraction of the smooth muscle with extraluminally applied MCh (3×10^{-7} M). When NaCl and D-M were used to increase osmolarity, relaxation responses of LPS-treated animals were potentiated in tracheas from LPS-treated animals (Fig. 5; Table 4). However, LPS treatment had no effect on responses to KCl ($n = 4$; data not shown), and it reduced significantly the potency of urea ($n = 6$; data not shown).

Effect of L-NAME and Indomethacin on Hyperosmolar Solution-Induced Smooth Muscle Relaxation in Isolated Perfused Trachea from Saline and LPS-Treated Guinea Pigs. Because LPS is known to up-regulate mRNA of both inducible NO synthase and cyclooxygenase (Okamoto et al., 1998), the effect of L-NAME and indomethacin on relaxant responses to hyperosmolar challenge was examined. This was done to rule out the possible contribution of NO and prostaglandins to the potentiation of EpDRF-induced smooth muscle relaxation after LPS treatment because both are capable of relaxing smooth muscle (Folkerts and Nijkamp, 1998). Neither L-NAME nor indomethacin affected responses to hyperosmolarity (NaCl




Fig. 3. Representative Harris' hematoxylin-eosin stained histological sections obtained 18 h after saline treatment (A, C, and E) and LPS treatment (B, D, and F). A and B are sections of distal trachea, C and D are sections of bronchi, and E and F are sections of the alveolar regions from saline- and LPS-treated animals, respectively. Sections of the bronchi and alveolar region were taken from the left lobe of the lung. Scale bars (50, 200, and 100 μ m) found in B, D, and F also apply to A, C, and E, respectively. These results are representative of eight different experiments from each treatment group, using animals treated specifically for histological analysis.




Fig. 4. Lack of effect of LPS treatment on tracheal smooth muscle reactivity to extraluminally and intraluminally applied MCh in epithelium-intact (A) and -denuded (B) isolated perfused trachea. Extraluminal and intraluminal MCh concentration-response curves were obtained from tracheas removed from saline- and LPS-treated animals 18 h post-treatment. Saline- and LPS-treated, $n = 6$, for epithelium-intact, and $n = 4$ for epithelium-denuded.

TABLE 2

Effect of LPS treatment on in vitro airway reactivity to MCh in epithelium-intact perfused trachea
The tracheas were removed from the animals 18 h post-treatment.

Treatment (n)	EC ₅₀ (M) (95% CI)		Maximum Response ΔP (cm H ₂ O)		IL/EL Maximum Response Ratio ^a
	EL	IL	EL	IL	
Saline (6)	1.9 × 10 ⁻⁷ (1.4–2.5)	3.7 × 10 ⁻⁴ (1.5–9.1) ^b	10.7 ± 1.5	8.4 ± 1.1 ^c	0.78 ± 0.03
LPS (6)	2.1 × 10 ⁻⁷ (1.7–2.4)	3.2 × 10 ⁻⁴ (1.8–5.8) ^d	6.6 ± 0.9	4.7 ± 0.9 ^e	0.70 ± 0.12

EL, extraluminal; IL, intraluminal.

^a The IL/EL maximum response ratio is used to gauge the modulatory effect of the epithelium on the IL maximum contractile response. In the absence of the epithelium, the ratio is unity, whereas it becomes less than unity in the presence of the epithelium. As the modulatory effect of the epithelium becomes more inhibitory, the ratio becomes smaller (Fedan and Frazer, 1992; Fedan et al., 2000).

^b Significantly greater than saline (EL).

^c Significantly less than saline (EL).

^d Significantly greater than LPS (EL).

^e Significantly less than LPS (EL).

TABLE 3

Effect of LPS treatment on in vitro reactivity to MCh in epithelium-denuded perfused trachea
The tracheas were removed from the animals 18 h post-treatment.

Treatment (n)	EC ₅₀ (M) (95% CI)		Maximum Response ΔP (cm H ₂ O)		IL/EL Maximum Response Ratio ^a
	EL	IL	EL	IL	
Saline (4)	2.1 × 10 ⁻⁶ (0.8–5.2)	7.7 × 10 ⁻⁶ (2.0–30.0)	3.8 ± 0.7	3.5 ± 0.6	0.97 ± 0.12
LPS (4)	1.1 × 10 ⁻⁶ (0.5–2.2)	1.5 × 10 ⁻⁶ (0.8–2.8)	3.9 ± 0.4	3.5 ± 0.3	1.00 ± 0.06

EL, extraluminal; IL, intraluminal.

^a See Fig. 2 legend.




Fig. 5. Effect of LPS treatment on EpDRF-induced smooth muscle relaxation elicited with intraluminal MKHS made hyperosmolar with added NaCl (A) or D-M (B). The concentration-response curves were obtained from tracheas removed from saline- and LPS-treated animals 18 h post-treatment. The tracheal smooth muscle was contracted with extraluminally applied MCh (3×10^{-7} M) before NaCl or D-M were added to the intraluminal bath in stepwise-increasing, cumulative concentrations. Saline- and LPS-treated, $n = 8$ for NaCl and $n = 6$ for D-M. *, $p < 0.05$ compared with saline.

added) in isolated perfused tracheas from saline- and LPS-treated guinea pigs (Figs. 6 and 7). These findings buttress the view that EpDRF is neither a prostanoid nor NO in tracheas removed from either control or LPS-treated guinea pigs.

Effect of LPS Treatment on Basal V_t . In tracheas from LPS-treated animals, the basal V_t was increased by 80% compared with tracheas from saline-treated animals. The basal V_t values were -14.6 ± 1.2 and -28.3 ± 1.9 mV for saline- and LPS-treated animals, respectively ($p < 0.05$).

Effect of LPS in Vitro on Basal V_t in Tracheas from Naive Animals. In view of the hyperpolarization observed in tracheas from LPS-treated animals, it was of interest to investigate whether LPS applied in vitro could affect V_t across tracheas from naive animals. Using a concentration

TABLE 4

Effect of LPS treatment on relaxation responses of perfused trachea to hyperosmolarity
The tracheas were removed from the animals 18 h post-treatment.

Treatment (n)	EC ₅₀ (mosM) ^a (95% CI)	Maximum Response (Percentage of MCh) ^b
NaCl		
Saline (8)	11.4 (8.2–15.8)	78.2 ± 6.2
LPS (8)	10.4 (8.4–12.8)	113.3 ± 9.5 ^c
D-Mannitol		
Saline (6)	15.2 (8.2–28.2)	86.1 ± 4.9
LPS (6)	15.9 (11.1–22.7)	109.7 ± 8.5 ^c

^a Values refer to milliosmoles of osmolyte added to the intraluminal MKHS.

^b Relaxation responses in centimeters of H₂O were normalized as a percentage of the extraluminal MCh (3×10^{-7} M)-induced contraction.

^c Significantly greater than saline-treated controls.

(10 μg/ml) that was shown previously to inhibit MCh-induced airway smooth muscle contraction in vitro (Fedan et al., 1995), LPS was added simultaneously to the solutions bathing the extraluminal and intraluminal surfaces of tracheas isolated from nontreated animals. Within minutes, LPS began to hyperpolarize the epithelium (data not shown). The time required for stabilization to occur varied from trachea to trachea but ranged between 1 to 3 h. Over similar intervals, control tracheas run in parallel exhibited a small hyperpolarization, which did not achieve statistical significance.

Effect of Indomethacin, L-NAME, Atropine, and SNP on Basal V_t . To determine whether the increase in the basal V_t after LPS treatment could involve prostanoids or NO, the effects of indomethacin (3×10^{-6} M) and L-NAME (10^{-4} M) were examined. Prostaglandins and NO have been shown to stimulate Cl⁻ secretion (Tamaoki et al., 1992; Tamai and Gaginella, 1993). Indomethacin decreased V_t in both treatment groups (Fig. 8), but the effect was significantly greater in the LPS-treated group: $+5.5 \pm 0.7$ mV for the LPS-treated group compared with $+3.1 \pm 0.5$ mV for the saline-treated group ($p < 0.05$).

L-NAME slightly hyperpolarized the epithelium in both




Fig. 6. Lack of effect of extraluminally and intraluminally applied L-NAME (10^{-4} M) on EpDRF-induced smooth relaxation elicited with intraluminal MKHS made hyperosmolar with added NaCl. The concentration-response curves were obtained from tracheas removed from saline- (A) and LPS-treated (B) animals 18 h post-treatment. The tracheal smooth muscle was contracted with extraluminally applied MCh (3×10^{-7} M) before NaCl was added to the intraluminal bath in stepwise-increasing, cumulative concentrations. Saline- and LPS-treated, $n = 4$.




Fig. 7. Lack of effect of extraluminally and intraluminally applied indomethacin (3×10^{-6} M) on EpDRF-induced smooth relaxation elicited with intraluminal MKHS made hyperosmolar with added NaCl. The concentration-response curves were obtained from tracheas removed from saline- (A) and LPS-treated (B) animals 18 h post-treatment. The tracheal smooth muscle was contracted with extraluminally applied MCh (3×10^{-7} M) before NaCl was added to the intraluminal bath in stepwise-increasing, cumulative concentrations. Saline- and LPS-treated, $n = 4$.

treatment groups; there was no effect of LPS on the magnitude of the hyperpolarization ($n = 4$; data not shown). SNP (10^{-6} M) caused a very small depolarization ($n = 4$; data not shown). These results suggest that V_t is under a slight influence of NO. However, the effect of L-NAME could have resulted from muscarinic receptor blockade (Buxton et al., 1993). Therefore, the effect of atropine (10^{-6} M) was examined to compare the effect of muscarinic receptor blockade with that of L-NAME. Atropine caused depolarization or hyperpolarization responses in different preparations. On average atropine increased V_t by 0.9 ± 0.5 mV in tracheas from saline-treated animals and decreased V_t by 0.3 ± 0.6 mV in tracheas from LPS-treated animals ($p > 0.05$; data not shown). Therefore, atropine did not mimic L-NAME, but the results suggest that basal V_t is under a modest cholinergic influence.

Effect of LPS Treatment on Bioelectric Reactivity to MCh. In tracheas from both saline- and LPS-treated ani-




Fig. 8. Effect of indomethacin on the basal V_t of isolated perfused trachea from saline- and LPS-treated animals. A, millivolt change in the basal V_t evoked by the simultaneous addition of indomethacin (3×10^{-6} M) to the intraluminal and extraluminal baths. B, V_t values before and after the addition of indomethacin. Saline- and LPS-treated, $n = 5$. *, $p < 0.05$ compared with saline-treated controls. †, $p < 0.05$ compared with saline-treated controls (-indomethacin). ††, $p < 0.05$ compared with saline-treated controls (-indomethacin). †††, $p < 0.05$ compared with LPS-treated (-indomethacin).

mals, extraluminally applied MCh produced a biphasic concentration-response curve (Fig. 9A; Table 5). At MCh concentrations less than 10^{-6} M, V_t increased, whereas at MCh concentrations greater than 10^{-6} M, V_t decreased. LPS treatment decreased the MCh EC_{50} value for hyperpolarization and increased the maximum hyperpolarization. In contrast, the MCh EC_{50} value and maximum response for depolarization were not affected. The intraluminal application of MCh also generated biphasic concentration-response curves in both treatment groups (hyperpolarizing at MCh concentrations less than 10^{-3} M and depolarizing at MCh concentrations greater than 10^{-3} M; Fig. 9B; Table 5). In the case of intraluminally applied MCh, the maximum hyperpolarization response was greater in the LPS-treated group with respect to the saline-treated controls, whereas there was no difference in the depolarization responses among the two treatment groups.

Effect of LPS Treatment on the V_t Responses to Hyperosmolarity. Smooth muscle relaxation elicited by hyperosmolarity has been reported to be preceded by depolarization of the epithelium (Dortch-Carnes et al., 1999), indicating




Fig. 9. Effect of LPS treatment on epithelial reactivity to extraluminally (A) and intraluminally (B) applied MCh. Positive values refer to depolarization, whereas negative values refer to hyperpolarization. Saline-treated, $n = 7$ and LPS-treated, $n = 6$. *, $p < 0.05$ compared with saline-treated controls.

TABLE 5
Effect of LPS treatment on epithelial bioelectric reactivity to MCh

Treatment (n)	EC ₅₀ (M) (95% CI)		Maximum Response ΔV_t (mV) ^a	
	EL	IL	EL	IL
Hyperpolarization (n)				
Saline (7)	6.2×10^{-8} (2.9–13.0)	9.8×10^{-6} (2.1–45.0) ^b	-0.9 ± 0.3	-1.6 ± 0.3
LPS (6)	4.2×10^{-9} (1.6–11.0) ^c	2.7×10^{-6} (1.8–5.9) ^d	-3.0 ± 0.7^b	-5.3 ± 1.6^e
Depolarization				
Saline (7)	2.6×10^{-6} (1.5–4.3)	7.6×10^{-2} (2.3–23.0) ^b	2.3 ± 0.6	4.0 ± 1.1
LPS (6)	1.9×10^{-6} (1.0–3.7)	2.8×10^{-2} (0.7–10.0) ^d	4.8 ± 1.6	5.0 ± 1.1

EL, extraluminal; IL, intraluminal.

^a Values refer to the change in V_t .

^b Significantly greater than saline (EL).

^c Significantly less than saline (EL).

^d Significantly greater than LPS (EL).

^e Significantly greater than saline (IL).

that the two events may be functionally linked. Because LPS treatment potentiated smooth muscle relaxation in response to hyperosmolar challenge of the epithelium, we sought to determine whether LPS treatment also affected epithelial bioelectric responses to hyperosmolar solution.

NaCl concentration-response curves for bioelectric responses were generated in the presence of extraluminally applied 3×10^{-7} M MCh (Fig. 10), to mimic the conditions utilized for studying relaxation responses. In tracheas from LPS-treated animals, the intraluminal addition of NaCl led to a significantly greater depolarization response compared with saline-treated controls; however, the epithelium eventually depolarized to the same level in each treatment group (Fig. 10; Table 6).

The concentration-response relationships for hyperosmolarity-induced epithelial depolarization and smooth muscle relaxation are compared in Fig. 11. In both treatment groups, the EC₅₀ value for depolarization was significantly greater than the EC₅₀ value for EpDRF-induced smooth muscle relaxation (Table 6).

Effect of Amiloride on Basal V_t . It has been demonstrated previously that relaxation and bioelectric responses of the perfused trachea to hyperosmolar solutions involve epithelial Na⁺ transport (Fedan et al., 1999; Wu et al., 2003). We, therefore, considered the possibility that the bioelectric changes resulting from LPS treatment could involve a change in Na⁺ transport (Fig. 12). In tracheas from both

saline- and LPS-treated animals amiloride (3×10^{-5} M) applied to the intraluminal bath decreased basal V_t . The depolarization was larger in the LPS-treated group. In the presence of amiloride, there was no longer any difference in the V_t values between the saline- and LPS-treated groups.

Discussion

Several general conclusions can be drawn from this study. First, systemic, noninhalational administration of LPS induces in vivo airway hyporeactivity to inhaled MCh in the absence of an inflammatory response. Second, LPS treatment potentiates EpDRF-induced airway smooth muscle relaxation in response to hyperosmolar solution in an osmolyte-specific manner and is associated with potentiated epithelial depolarization responses. Finally, epithelial depolarization does not always accompany EpDRF-induced smooth muscle relaxation, as judged by the disparity in the potency of NaCl as a relaxant and depolarizing osmolyte. This could indicate, at least in the case of NaCl, either that multiple mechanisms underlie epithelium-dependent relaxation, and only one of these is linked to depolarization of the epithelium, or that EpDRF release and epithelial depolarization are coincident but unrelated phenomena.

The LPS treatment protocol (4 mg/kg i.p.; 18 h post-treatment) used in this study caused in vivo airway hyporeactivity to inhaled MCh, whereas no differences in in vitro reactivity to MCh existed in the isolated perfused tracheas taken from saline- and LPS-treated animals. There have been several reports concerning the effect of LPS treatment on in vitro and in vivo guinea pig airway reactivity to contractile agents. LPS has been reported to induce hyperreactivity (Folkerts et al., 1989; Yamawaki et al., 1990; Van Oosterhout et al., 1991), hyporeactivity (Folkerts et al., 1988), and no change in reactivity of airway smooth muscle (Fedan et al., 1995). In the present study, there were no differences in airway reactivity to MCh in epithelium-intact and -denuded isolated perfused tracheas. This is consistent with a previous study from our laboratory (Fedan et al., 1995), even though the 18 h post-treatment time point was not examined. In vivo, LPS-treated guinea pigs were hyporeactive to the effects of inhaled MCh. This is not consistent with the earlier published report from our laboratory (Fedan et al., 1995), but again, the 18 h post-treatment time point may account for the differences.

Because it is well known that LPS up-regulates inducible NO synthase, it was plausible that NO may have contributed




Fig. 10. Effect of LPS treatment on epithelial bioelectric reactivity to intraluminal hyperosmolar NaCl in the presence of extraluminally applied MCh (3×10^{-7} M). A, millivolt change in V_t evoked by the addition of intraluminal NaCl. B, data from the same experiments described in A, but the plot shows the actual V_t values after the addition of intraluminal NaCl. Saline- and LPS-treated, $n = 4$, $p < 0.05$ compared with saline-treated controls.

TABLE 6
Effect of LPS treatment on mechanical and bioelectric reactivity to intraluminal hyperosmolar NaCl

Treatment (n)	EC ₅₀ (mosM) ^f (95% CI)		Maximum Response	
	Relaxation (Percentage of MCh) ^a	Depolarization ΔV _t (mV) ^b	Relaxation (Percentage of MCh)	Depolarization ΔV _t (mV)
Saline (7)	11.4 (8.2–15.8)	121.6 (75.0–197.2) ^d	78.2 ± 6.2	15.3 ± 0.5
LPS (6)	10.4 (8.4–12.8)	183.6 (120.2–280.4) ^e	113.3 ± 9.5 ^f	38.0 ± 4.1 ^f

^a Relaxation responses (centimeters of H₂O) were normalized as a percentage of the extraluminal MCh-induced contraction; saline- and LPS-treated, n = 8.
^b Values refer to the change of V_t after the administration of extraluminal MCh; saline- and LPS-treated, n = 4.
^c Values refer to milliosmoles of NaCl added to the intraluminal MKHS.
^d Significantly greater than saline (relaxation).
^e Significantly greater than LPS (relaxation).
^f Significantly greater than saline-treated controls.




Fig. 11. Effect of LPS treatment on the relationship between smooth muscle relaxation and epithelial depolarization elicited by hyperosmolar NaCl. The preparations were contracted with extraluminally applied MCh before NaCl was added cumulatively to the intraluminal bath. Saline- and LPS-treated for relaxation, n = 8 and saline- and LPS-treated for depolarization, n = 4.




Fig. 12. Effect of amiloride on the basal V_t of isolated perfused trachea from saline- and LPS-treated animals. A, change in the basal V_t evoked by the addition of amiloride to the intraluminal bath. B, V_t before and after the addition of amiloride. Saline- and LPS-treated, n = 5. *, p < 0.05 compared with saline-treated controls. †, p < 0.05 compared with saline-treated controls (-amiloride). ††, p < 0.05 compared with saline-treated controls (-amiloride). †††, p < 0.05 compared with LPS-treated (-amiloride).

to in vivo airway hyporeactivity to inhaled MCh; however, there was no effect of L-NAME on responses to MCh in saline- or LPS-treated, conscious animals. Thus, NO is not responsible for hyporeactivity to inhaled MCh after LPS administration. In contrast, Nijkamp et al. (1993) found in intubated anesthetized animals that L-NAME increased airway resistance.

Folkerts et al. (1988) reported the development of airway

hyperreactivity to histamine and inflammation of the airways 24 h after inhalation exposure to LPS. In the report of in vivo airway hyporeactivity to histamine after LPS treatment (Folkerts et al., 1988), the airways of the guinea pigs became infiltrated with large numbers of neutrophils and monocytes. In contrast, there were no signs of pulmonary inflammation or changes in the physical structure of the airways 18 h after LPS treatment in the present study. The lack of pulmonary inflammation in our LPS-treated guinea pigs suggests that the route of LPS administration is important to the development of inflammation and that inflammation is not a prerequisite for LPS-induced changes in the EpDRF system.

Paradoxically, airway hyporeactivity to MCh in vivo was not manifest in vitro in isolated perfused tracheas from LPS-treated animals. There could be several reasons for this. First, the mechanism underlying the in vivo airway hyporeactivity may not involve the trachea, but could involve intrapulmonary bronchi and bronchioles. Second, the other mechanisms responsible for the development of in vivo airway hyporeactivity could be lost in vitro, such as neural pathways involved in the control of reactivity.

Although there was no effect of LPS on airway smooth muscle reactivity to MCh in the perfused trachea, LPS did alter the basal bioelectric properties of the epithelium as well as its reactivity to MCh, i.e., LPS significantly hyperpolarized the trachea. LPS increased the conductance of Ca²⁺-activated K⁺ channels in vascular smooth muscle cells (Hoang et al., 1997). In our system, LPS treatment could have initiated epithelial cell Ca²⁺ influx and activation of K⁺ channels leading to hyperpolarization, especially because Ca²⁺-activated K⁺ channels have been shown to be involved in the synthesis and/or release of EpDRF in response to MCh in bronchial strips (Tamaoki et al., 1997). However, we observed no effect of iberiotoxin or glibenclamide on basal V_t or hyperosmolarity induced relaxation responses (Fedan et al., 2003b; Wu et al., 2003).

Biphasic concentration-response curves were obtained in response to extraluminal and intraluminal MCh in both treatment groups. The greater potency of extraluminal MCh compared with intraluminal MCh suggests the existence of multiple muscarinic receptors on the basolateral epithelial membrane. Our results are consistent with the electrophysiological effects produced by MCh in guinea pig nasal epithelium (Suzuki et al., 1999), in which MCh produced triphasic short-circuit current responses: a transient increase followed by a small decrease and then, finally, a sustained increase. The authors concluded that these changes were brought about by alterations in Cl⁻ secretion and amiloride-sensitive Na⁺ transport (Suzuki et al., 1999). MCh is capable of acti-

vating and/or inhibiting ion transport after its interaction with more than one muscarinic receptor subtype (Somlyo and Somlyo, 1994; Janssen et al., 1998; Roux et al., 1998). The activation of more than one channel was most likely responsible for the biphasic MCh concentration-response curve, but it is not known at present which muscarinic receptor subtypes were initiating these responses, i.e., M_1 , M_2 , and M_3 muscarinic receptors have been localized in the guinea pig lung (Mak and Barnes, 1990). A high-affinity receptor seems to be associated with hyperpolarization, whereas a low-affinity receptor may be linked to depolarization. The effect of LPS treatment, i.e., potentiation of hyperpolarization responses, seems to be mediated by the high-affinity hyperpolarizing receptor.

To explore retained changes in the airway wall in vitro that might have affected responses to MCh in vivo, we examined whether hyperosmolar solution-induced relaxation mechanism is altered after LPS treatment and observed that relaxation was potentiated. The observation that EpDRF-induced relaxation was potentiated after LPS treatment at the time that the airways were hyporeactive to inhaled MCh supports the general view that EpDRF modulates airway reactivity. This is the third instance in which alterations in the actions of EpDRF have been observed to accompany changes in in vivo and/or in vitro airway reactivity. For example, exposure to ozone causes a decrease in the actions of EpDRF, which is associated with in vivo and in vitro airway hyperreactivity to MCh (Fedan et al., 2000). In addition, sensitization and challenge of guinea pigs with ovalbumin potentiated EpDRF-induced relaxation, which is accompanied by in vitro airway hyporeactivity to MCh (Warner et al., 1996).

We observed that LPS treatment potentiated hyperosmolarity-induced smooth muscle relaxation and depolarization of the epithelium. These results support the hypothesis that the relaxation and bioelectric events are functionally linked. This hypothesis originated in the observation that depolarization of the epithelium in response to elevation of the intraluminal or extraluminal osmolarity preceded the relaxation response (Dortch-Carnes et al., 1999). However, in the present study, in both treatment groups, the smooth muscle relaxed at lower hyperosmolar concentrations than were required to elicit depolarization. Relaxation occurred when as little as 0.8 mosM of NaCl was added to the intraluminal bath; however, depolarization did not occur until the osmolarity was raised by 8.4 mosM. If epithelial depolarization in response to hyperosmolarity signaled the release of EpDRF (Dortch-Carnes et al., 1999), the epithelium would have depolarized with the same osmolar concentration dependence as that for relaxation. It is necessary to reformulate the hypothesis linking bioelectric and mechanical events as cause and effect phenomena, at least insofar as NaCl is concerned. At low hyperosmolar concentrations, EpDRF release occurs independently of epithelial depolarization, whereas at higher elevations of osmolarity the epithelium becomes depolarized, which may signal the release of greater quantities of EpDRF. The meaning of these two pathways in terms of ion transport is not currently understood. The effects of low level hyperosmolarity could be mediated by electrically neutral ion transport, whereas at the higher osmotic particle concentrations permeation through ion channels also would be engaged. Transepithelial depolarization might be a coincident phenomenon associated with EpDRF release stim-

ulated by a different, nonelectrogenic ion transport mechanism that is more sensitive to extracellular osmolarity changes. For example, exposure of cells to hyperosmolar solution causes cell alkalization (Miyata et al., 2000). The Na^+ channel blocker amiloride and the Cl^- channel blockers 4,4'-diisothiocyano-2,2'-stilbene disulfonate and 5-nitro-2-(3-phenylpropylamino)benzoic acid both inhibit relaxation responses to hyperosmolarity, and both agents are capable of inhibiting the Na^+, H^+ exchanger and the Cl^-, HCO_3^- exchanger (Kreydiyyeh, 2000; Lamb et al., 2000), respectively. These pharmacological inhibitors could be predicted to influence intracellular pH.

Finally, we investigated mechanism(s) that could have participated in the hyperpolarization of the epithelium after LPS treatment. Because LPS is known to induce NO synthase and cyclooxygenase (Okamoto et al., 1998), the effect of inhibitors of these enzymes on V_t was examined. Although NO has been shown to influence epithelial ion transport (Tamai and Gagnella, 1993), the small electrophysiological effects of L-NAME and SNP suggest that NO plays little, if any, role in regulating ion transport in guinea pig tracheal epithelium. On the other hand, indomethacin decreased markedly V_t in both treatment groups, but significantly more so in the LPS-treated group. This depolarizing effect is consistent with an inhibition of prostanoid-stimulated Cl^- secretion (Tamaoki et al., 1992). The fact that indomethacin decreased V_t to a greater extent in the LPS-treated group suggests that the role of prostanoids is heightened after LPS treatment. The mechanism whereby this occurs cannot be determined from our experiments, but it could involve up-regulation of cyclooxygenase, or changes in prostaglandin receptors or postreceptor signaling pathways. Nevertheless, the results suggest that prostanoid-stimulated Cl^- secretion may contribute to the hyperpolarization of the epithelium after LPS treatment. Inasmuch as the hyperpolarization caused by LPS was abolished by amiloride and inhibited by indomethacin, it is tempting to speculate that a functional relationship between prostanoids and Na^+ channels exists, which is altered by LPS.

Amiloride applied to the intraluminal surface of the trachea abolished the effect of LPS treatment on basal V_t . This suggests that the hyperpolarization of the epithelium after LPS treatment was due primarily to an increase in Na^+ transport. An increase in Na^+ removal from the lumen, via these channels, would generate a more negative V_t . Amiloride has been shown to inhibit EpDRF-mediated relaxation responses to hyperosmolar solution (Fedan et al., 1999). An increase in the activity of amiloride-sensitive Na^+ channels may therefore be linked to potentiated responses to hyperosmolar challenge, the increase in basal V_t , and the potentiated V_t response to hyperosmolarity.

In conclusion, LPS treatment induces in vivo airway hyporeactivity to MCh that may be due, at least in part, to an increase in the actions of EpDRF. In addition, our results demonstrate that LPS treatment induces appreciable alterations in basal V_t of tracheal epithelium as well as its reactivity to MCh. These changes affect the airway smooth muscle indirectly in that the EpDRF-mediated relaxation response to hyperosmolarity is also potentiated. It is conceivable that such alterations in the airways could exist during gram negative bacterial infections of the airways.

Acknowledgments

We thank Deborah C. Sbarra, Nicole Diotte, Janet A. Dowdy, and Scott G. Olson for technical assistance with the in vivo or in vitro airway reactivity studies, and Dean Newcomer and Patsy Willard for preparing the histological sections.

References

- Buxton IL, Cheek DJ, Eckman D, Westfall DP, Sanders KM, and Keef KD (1993) N^G -Nitro L-arginine methyl ester and other alkyl esters of arginine are muscarinic receptor antagonists. *Circ Res* 72:387-395.
- Dortch-Carnes J, Van Scott MR, and Fedan JS (1999) Changes in smooth muscle tone during osmotic challenge in relation to epithelial bioelectric events in guinea pig isolated trachea. *J Pharmacol Exp Ther* 289:911-917.
- Fedan JS, Dowdy JA, Johnston RA, and Van Scott MR (2003a) Hyperosmolar solution effects in guinea-pig airways. I. Mechanical responses to relative changes in osmolarity. *J Pharmacol Exp Ther* 308:10-18.
- Fedan JS, Dowdy JA, Van Scott MR, Wu DX-Y, and Johnston RA (2003b) Hyperosmolar solution effects in guinea-pig airways. III. Studies on the identity of epithelium-derived relaxing factor in isolated perfused trachea using pharmacological agents. *J Pharmacol Exp Ther* 308:30-36.
- Fedan JS and Frazer DG (1992) Influence of epithelium on the reactivity of guinea pig isolated perfused trachea to bronchoactive drugs. *J Pharmacol Exp Ther* 262:741-750.
- Fedan JS, Hay DWP, Farmer SG, and Raeburn D (1988) Epithelial cells: modulation of airway smooth muscle reactivity. In *Asthma: Basic Mechanisms and Clinical Management* (Barnes PJ, Rodger IW, and Thomson NC eds) pp 143-162. Academic Press, New York.
- Fedan JS, Millecchia LL, Johnston RA, Rengasamy A, Hubbs A, Dey RD, Yuan LX, Watson D, Goldsmith WT, Reynolds JS, et al. (2000) Effect of ozone treatment on airway reactivity and epithelium-derived relaxing factor in guinea pigs. *J Pharmacol Exp Ther* 293:724-734.
- Fedan JS, Warner TE, Yuan LX, Robinson V, and Frazer DG (1995) Nitric oxide synthase inhibitor and lipopolysaccharide effects on reactivity of guinea pig airways. *J Pharmacol Exp Ther* 272:1141-1150.
- Fedan JS, Yuan LX, Chang VC, Viola JO, Cutler D, and Pettit LL (1999) Osmotic regulation of airway reactivity by epithelium. *J Pharmacol Exp Ther* 289:901-910.
- Folkerts G, Engels F, and Nijkamp FP (1989) Endotoxin-induced hyperreactivity of the guinea-pig isolated trachea coincides with decreased prostaglandin E₂ production by the epithelial layer. *Br J Pharmacol* 96:388-394.
- Folkerts G, Henricks PA, Slootweg PJ, and Nijkamp FP (1988) Endotoxin-induced inflammation and injury of the guinea pig respiratory airways cause bronchial hyperreactivity. *Am Rev Respir Dis* 137:1441-1448.
- Folkerts G and Nijkamp FP (1998) Airway epithelium: more than just a barrier! *Trends Pharmacol Sci* 19:334-341.
- Goldie RG and Hay DWP (1997) Epithelium-dependent responsiveness of airway smooth muscle: the role of epithelium-derived relaxant factors. In *Asthma* (Barnes PJ, Grunstein MM, Leff AR, and Woolcock AJ eds) pp 901-915. Lippincott-Raven, Philadelphia.
- Held HD and Uhlig S (2000) Mechanisms of endotoxin-induced airway and pulmonary vascular hyperreactivity in mice. *Am J Respir Crit Care Med* 162:1547-1552.
- Hoang LM, Chen C, and Mathers DA (1997) Lipopolysaccharide rapidly activates K^+ channels at the intracellular membrane face of rat cerebral artery smooth muscle cells. *Neurosci Lett* 231:25-28.
- Janssen LJ, Daniel EE, and Rodger IW (1998) Airway smooth muscle cells: structure and function. In *Asthma: Basic Mechanisms and Clinical Management* (Barnes PJ, Rodger IW, and Thomson NC eds) pp 90-112. Elsevier, San Diego.
- Kreydiyyeh SI (2000) Cyclic AMP and furosemide stimulate the Na^+-K^+ ATPase in isolated rat jejunal cells. *Pharmacol Res* 41:179-185.
- Lamb FS, Kouy NW, and Lewis SJ (2000) Role of Cl^- channels in alpha-adrenoceptor-mediated vasoconstriction in the anesthetized rat. *Eur J Pharmacol* 401:403-412.
- Lawrence TE, Millecchia LL, Frazer DG, and Fedan JS (1997) Pulmonary dendritic cell distribution and prevalence in guinea pig airways: effect of ovalbumin sensitization and challenge. *J Pharmacol Exp Ther* 282:995-1004.
- Mak JC and Barnes PJ (1990) Autoradiographic visualization of muscarinic receptor subtypes in human and guinea pig lung. *Am Rev Respir Dis* 141:1559-1568.
- Michel O, Duchateau J, and Sergysels R (1989) Effect of inhaled endotoxin on bronchial reactivity in asthmatic and normal subjects. *J Appl Physiol* 66:1059-1064.
- Michel O, Kips J, Duchateau J, Vertongen F, Robert L, Collet H, Pauwels R, and Sergysels R (1996) Severity of asthma is related to endotoxin in house dust. *Am J Respir Crit Care Med* 154:1641-1646.
- Miyata Y, Muto S, Yanagiba S, and Asano Y (2000) Extracellular Cl^- modulates shrinkage-induced activation of Na^+/H^+ exchanger in rat mesangial cells. *Am J Physiol* 278:C1218-C1229.
- Munakata M, Huang I, Mitzner W, and Menkes H (1989) Protective role of epithelium in the guinea pig airway. *J Appl Physiol* 66:1547-1552.
- Munakata M, Masaki Y, Sakuma I, Ukita H, Otsuka Y, Homma Y, and Kawakami Y (1990) Pharmacological differentiation of epithelium-derived relaxing factor from nitric oxide. *J Appl Physiol* 69:665-670.
- Munakata M, Mitzner W, and Menkes H (1988) Osmotic stimuli induce epithelium-dependent relaxation in the guinea pig trachea. *J Appl Physiol* 64:466-471.
- Nijkamp FP, van der Linde HJ, and Folkerts G (1993) Nitric oxide synthesis inhibitors induce airway hyperresponsiveness in the guinea pig in vivo and in vitro. Role of the epithelium. *Am Rev Respir Dis* 148:727-734.
- Okamoto H, Ito O, Roman RJ, and Hudetz AG (1998) Role of inducible nitric oxide synthase and cyclooxygenase-2 in endotoxin-induced cerebral hyperemia. *Stroke* 29:1209-1218.
- Pauwels RA, Kips JC, Peleman RA, and Van Der Straeten ME (1990) The effect of endotoxin inhalation on airway responsiveness and cellular influx in rats. *Am Rev Respir Dis* 141:540-545.
- Roux E, Molimard M, Savineau JP, and Marthan R (1998) Muscarinic stimulation of airway smooth muscle cells. *Gen Pharmacol* 31:349-356.
- Salter M, Knowles RG, and Moncada S (1991) Widespread tissue distribution, species distribution and changes in activity of Ca^{2+} -dependent and Ca^{2+} -independent nitric oxide synthases. *FEBS Lett* 291:145-149.
- Somlyo AP and Somlyo AV (1994) Signal transduction and regulation in smooth muscle. *Nature (Lond)* 372:231-236.
- Suzuki K, Kawahara K, Terada N, Nomura T, Konno A, and Fukuda Y (1999) Effect of ion transport inhibitors and methacholine on short-circuit current of isolated guinea pig nasal epithelium. *Jpn J Physiol* 49:99-106.
- Tamai H and Gaginella TS (1993) Direct evidence for nitric oxide stimulation of electrolyte secretion in the rat colon. *Free Radic Res Commun* 19:229-239.
- Tamaoki J, Kanemura T, Sakai N, Isono K, Chiyotani A, Yamauchi F, Takizawa T, and Konno K (1992) Prostaglandin D₂ increases Cl^- secretion across canine tracheal epithelium through cyclo-oxygenase stimulation and cAMP production. *Acta Physiol Scand* 145:1-6.
- Tamaoki J, Tagaya E, Isono K, Kondo M, and Konno K (1997) Role of Ca^{2+} -activated K^+ channel in epithelium-dependent relaxation of human bronchial smooth muscle. *Br J Pharmacol* 121:794-798.
- Toward TJ and Bradley KJ (2000) Airway reactivity, inflammatory cell influx and nitric oxide in guinea-pig airways after lipopolysaccharide inhalation. *Br J Pharmacol* 131:271-281.
- Van Oosterhout AJ, Woutersen-Van Nijnanten FM, and Nijkamp FP (1991) Effect of cyclosporin-A treatment on endotoxin-induced airway hyperreactivity in vivo and in vitro in guinea-pigs. *Naunyn-Schmiedeberg's Arch Pharmacol* 343:542-545.
- Warner T, Frazer DG, and Fedan J (1996) Hyporeactivity to methacholine in vitro accompanying hyperreactivity to methacholine in vivo in a guinea-pig model of asthma (Abstract). *Am J Respir Crit Care Med* 153:A627.
- Wu DX-Y, Johnston RA, Rengasamy A, Van Scott MR, and Fedan JS (2003) Hyperosmolar solution effects in guinea-pig airways. II. Epithelial bioelectric responses to relative changes in osmolarity. *J Pharmacol Exp Ther* 308:xxx-xxx.
- Yamawaki I, Tamaoki J, Kanemura T, Horii S, and Takizawa T (1990) Effects of lipopolysaccharide from *Pseudomonas aeruginosa* on airway smooth muscle functions in guinea pigs. *Respiration* 57:268-274.

Address correspondence to: Dr. Jeffrey S. Fedan, Pathology and Physiology Research Branch, National Institute for Occupational Safety and Health, 1095 Willowdale Rd., Morgantown, WV 26505-2888. E-mail: jsf2@cdc.gov