

Male infertility and environmental exposure to lead and cadmium

Susan Benoff^{1,2,3,*}, Asha Jacob^{1,2} and Ian R. Hurley¹

¹Division of Human Reproduction, Department of Obstetrics and Gynecology, North Shore University Hospital-New York University School of Medicine, Manhasset, New York, ²Department of Obstetrics and Gynecology, New York University School of Medicine, New York, New York, and ³Department of Cell Biology, New York University School of Medicine, New York, New York, USA

Humans are exposed occupationally and environmentally to metal aerosols including lead (Pb²⁺) and cadmium (Cd²⁺). These toxicants accumulate in male reproductive organs. Epidemiological studies have been equivocal about effects of Pb²⁺ and Cd²⁺ on hormone concentrations, male fertility and sperm parameters. Comparison of Pb²⁺ and Cd²⁺ concentrations in fertile and infertile men are problematic. Problem areas include failure to control confounding variables, but genetic polymorphisms as in somatic diseases may modulate Pb²⁺ and Cd²⁺ damage. Multiple calcium (Ca²⁺) and potassium (K⁺) channel isoforms have been identified in human testes and spermatozoa. These Ca²⁺ and K⁺ channels are involved in early events of acrosome reactions. Ca²⁺ channel are susceptible to Cd²⁺ poisoning and K⁺ channels to Pb²⁺. These channels offer entry paths for metallic toxicants into mature spermatozoa. Ion channel polymorphisms may cause differential sensitivities to Cd²⁺ and Pb²⁺, explaining in part prospective blinded studies showing high Cd²⁺ in varicocele-related human infertility and high Pb²⁺ in unexplained infertility. In both forms of male infertility the ability to undergo an acrosome reaction decreases. Reverse transcriptase–polymerase chain reaction assays for Ca²⁺ and K⁺ channel isoforms may identify susceptibility subgroups with lower resistance to environmental exposures.

Key words: acrosome reaction/cadmium/ion channels/lead/male infertility

TABLE OF CONTENTS

Introduction	107
Effect of occupational exposure to metal ions	108
Sources of Pb ²⁺ and Cd ²⁺ found in the environment	109
Examination of the effects of environmental exposure to metal ions	110
Metal ion concentrations in fertile and infertile men	111
Confounding variables	112
Genetic determinants of susceptibility	112
Ion channels and response to environmental metal exposures	113
Potential models for metal effects mediated by ion channels	113
Conclusions	116
Acknowledgements	117
References	117

Introduction

In recent years, there has been an increasing interest in the contribution of occupational and environmental exposures to toxic pollutants to declining sperm concentrations and human male infertility (Carlsen *et al.*, 1992; Auger *et al.*, 1995; Adamopoulos

et al., 1996; Becker and Berhane, 1997). We recognize that at least one of these studies (Carlsen *et al.*, 1992) has been criticized because it fails to consider that sperm counts clearly differ by geographical location (Fisch and Goluboff, 1996; Paulsen *et al.*, 1996). Nonetheless, geographic variations in semen quality (Fisch and Goluboff, 1996; Becker and Behane, 1997) may still be influenced by environmental factors (Fisch and Goluboff, 1996) which exhibit considerable variation between climatic seasons (Sram *et al.*, 1996). More importantly, environmental factors differ between areas (Friberg and Vahter, 1983; Svensson *et al.*, 1987; Buchancova *et al.*, 1994; Sram *et al.*, 1996), with higher amounts of pollutants closer to sources of industrialization (Benin *et al.*, 1999).

Unfortunately, relatively few studies have systematically addressed the impact of environmental exposures on human health in general (Buchancova *et al.*, 1994; Staessen *et al.*, 1996) or on human reproductive function (El-Zohairy *et al.*, 1996; Sram *et al.*, 1996). What limited data that does exist suggests that exposure to high values of airborne fine particles is associated with decreased semen quality, i.e. increased aberrant morphology and motility. Heavy and transition metal ions, e.g. cadmium (Cd²⁺), cobalt (Co²⁺), copper (Cu²⁺), iron (Fe^{2+/3+}), lead (Pb²⁺), mercury (Hg²⁺), magnesium (Mg²⁺), manganese (Mn²⁺), nickel (Ni²⁺)

*To whom correspondence should be addressed at: North Shore University Hospital, 300 Community Drive, Boas-Marks Biomedical Science Research Center, Room 125, Manhasset, New York 11030, USA. Telephone: 001 516 562 1050; Fax: 001 516 562 1022; E-mail: sbenoff@nshs.edu

and selenium ($\text{Se}^{2+}/4+$), are common components of such airborne particulates that are within the respirable range (Ragan and Mast, 1990). Pb^{2+} and Cd^{2+} preferentially accumulate in male reproductive organs (Danielsson *et al.*, 1984; Oldereid *et al.*, 1993; Jackson *et al.*, 1995). An increase in Pb^{2+} and Cd^{2+} often occur simultaneously (Stachel *et al.*, 1989). This review focuses on the effects of Pb^{2+} and Cd^{2+} . Our studies, which are summarized herein, indicate that Pb^{2+} and Cd^{2+} exposures are of particular importance to the production of the infertile state.

Effect of occupational exposure to metal ions

The correlation of trace metal exposures with human sperm production and function has been studied mainly in men occupationally exposed to high concentrations of airborne metal particulates (Danielsson *et al.*, 1984; Assennato *et al.*, 1986; Stachel *et al.*, 1989; Gennart *et al.*, 1992b; Bonde, 1993). The earliest studies were triggered by reports that semen values of another class of environmental toxicants, halogen-containing hydrocarbon pesticides, were associated with decreases in sperm counts and motility, increased percentages of spermatozoa with morphological abnormalities and male infertility (Wong *et al.*, 1979; Dougherty *et al.*, 1986; Ratcliffe *et al.*, 1987; Wagner *et al.*, 1990).

The possible xenobiotic exposures associated with industrial metal aerosols include metals such as aluminum (Al^{3+}), Cd^{2+} , $\text{Fe}^{2+}/^{3+}$, chromium (Cr^{2+}), Ni^{2+} , and Pb^{2+} , as well as nitrous gases and ozone (Stern *et al.*, 1986; Mortensen, 1988). Metal workers (e.g. welders, smelters, men engaged in the manufacture of alkaline batteries) comprise 0.5–2.0% of the general working population (Stern *et al.*, 1986). There is epidemiological evidence that exposure to industrial metal aerosols may be detrimental to the male reproductive system; fertility of men employed in the metal industry is apparently reduced when compared with men in other types of work. These statements are based upon studies reporting decreased family size, delay in time to conception and reduced sperm quality in metal workers (Bonde, 1993; Spinelli *et al.*, 1997). These problems have been correlated with both exposure dose and length of exposure (Bonde, 1990a), and are not reversible by short periods of non-exposure (Bonde, 1990b). However, despite these findings, attempts to correlate occupational exposure to Pb^{2+} and Cd^{2+} with semen quality and/or infertility have produced conflicting results.

Blood plasma and seminal plasma Pb^{2+} concentrations are elevated in exposed workers (Stachel *et al.*, 1989; Telisman *et al.*, 1990; Ng *et al.*, 1991; Gennart *et al.*, 1992a; Keck *et al.*, 1995; Aribarg and Sukcharoen, 1996; Robins *et al.*, 1997; see Table I). A positive relationship was detected between blood and seminal plasma Pb^{2+} concentrations (Aribarg and Sukhareon, 1996). In some studies, evidence is presented suggesting that men with a long duration of Pb^{2+} exposure in the workplace may have reduced fertility (Lancranjan *et al.*, 1975; Gennart *et al.*, 1992b; Lin *et al.*, 1996). However, in other studies, Pb^{2+} exposure has not been reported to decrease fertility (Coste *et al.*, 1991) or only exhibits a weak association with infertility (Robins *et al.*, 1997). Dose–response relationships and/or causal mechanisms remain to be defined (Winder, 1993). Seminal plasma Pb^{2+} concentrations and fertility status have not been correlated (Keck *et al.*, 1995).

Endocrine dysfunction has (Ferioli *et al.*, 1989; MacGregor and Mason, 1990; Ng *et al.*, 1991) and has not (Lancranjan *et al.*, 1975; Assennato *et al.*, 1986) been observed in men occupationally exposed to Pb^{2+} . Where occurring, Pb^{2+} -associated changes in hormone concentrations were thought to be mediated at the hypothalamic–pituitary level (Gustafson *et al.*, 1989; Ng *et al.*, 1991). These data are consistent with findings in animal models, where the primary toxic action of Pb^{2+} on male reproduction appears to be disruption of the hypothalamus–pituitary–testicular axis (Sokol *et al.*, 1985, 1994; Sokol, 1987; Kempinas *et al.*, 1994).

The majority of reports indicate that workers exposed to Pb^{2+} exhibit decreased sperm density and a high rate of teratozoospermia (Lancranjan *et al.*, 1975; Hu *et al.*, 1992; Lerda, 1992; Xuezhi *et al.*, 1992; Robins *et al.*, 1997) and may also present with a decreased sperm count (Assennato *et al.*, 1986). An inverse correlation between blood Pb^{2+} concentrations and decreased semen quality has been observed (Lancranjan *et al.*, 1975; Telisman *et al.*, 1990). The study by Telisman *et al.* (1990) also suggested that reproductive toxicity may occur at blood Pb^{2+} concentrations below the permissible exposure limit of 40 $\mu\text{g}/\text{dl}$. Thus, current occupational standards may not be sufficiently stringent (Winder, 1993).

In some studies, however, no relationship was detected between seminal plasma Pb^{2+} concentrations and semen profiles (Aribarg and Sukcharoen, 1996; Hovatta *et al.*, 1998). There are three possible explanations for this failure. Firstly, exposure levels were apparently very low as blood Pb^{2+} concentrations were generally <40 $\mu\text{g}/\text{dl}$, the recommended limit of exposure for workers in an occupational setting as defined by the World Health Organization (WHO, 1980). Secondly, in some cases the men studied had only short-term exposure to Pb^{2+} (Aribarg and Sukhareon, 1996). As in animal models (Kempinas *et al.*, 1994), duration of Pb^{2+} exposure appears to be an important factor when searching for a correlation between Pb^{2+} and changes in semen parameters (Lerda, 1992). Thirdly, measurements of metal ion concentrations in seminal plasma may be of less significance in the examination of environmental effects than direct measurements of metal ion concentrations in spermatozoa themselves. As an example of the latter, Hovatta *et al.* (1998) reported that sperm motility was inversely correlated with aluminium in spermatozoa but not in seminal plasma.

Whether Cd^{2+} concentrations in blood and seminal plasma are elevated following occupational exposures is still not clear (Table I). Early studies indicated that, based on autopsy findings, spermatogenesis was reduced in workers exposed to Cd^{2+} (Smith *et al.*, 1960) and that the family size was smaller than average (Favino *et al.*, 1968). More recent studies did not support these findings. Cd^{2+} exposures have not been associated with a significant reduction in semen quality (Tielmans *et al.*, 1999) or with a decreased probability of a live birth (Gennart *et al.*, 1992b). In contrast, occupational exposure to Cd^{2+} has been reported to increase the risk for prostate cancer (Elghany *et al.*, 1990). This is consistent with observations indicating that Cd^{2+} concentrations are increased in prostate neoplasms, compared with normal tissue (Ogunlewe and Osegbe, 1989; Brys *et al.*, 1997) and studies *in vitro* and in animal models have demonstrated a relationship

Table I. Effects of occupational exposures on the levels of Pb²⁺ and Cd²⁺ in blood and seminal plasma

	Unexposed	Exposed	Source
Pb ²⁺ (blood)	13.1 ± 3.5 µg/dl		Plechaty <i>et al.</i> (1977)
	23 ± 14 µg/dl	29–100.5 µg/dl*	Lancrajan <i>et al.</i> (1975)
	20.8 ± 2.8 µg/dl	50.5 ± 9.6 µg/dl*	Feroli <i>et al.</i> (1989)
	6.7 – 20.8 µg/dl	11.7 – 104.0 µg/dl*	Telisman <i>et al.</i> (1990)
	8.3 ± 2.8 µg/dl	35.2 ± 13.2 µg/dl*	Ng <i>et al.</i> (1991)
	20.9 ± 11.1 µg/dl	51.0 ± 8.0 µg/dl*	Gennart <i>et al.</i> (1992a)
	10.4 ± 3.3 µg/dl	46.3 ± 11.1 µg/dl*	Gennart <i>et al.</i> (1992b)
	23.5 ± 1.4 µg/100 g	86.6 ± 0.6 µg/100 g*	Lerda (1992)
	23.6 ± 47.1 µg/dl	47.1 ± 31.0 µg/dl*	Xuezhi <i>et al.</i> (1992)
	16.9 ± 2.6 µg/dl	26.7 ± 2.1 µg/dl*	El-Zohairy <i>et al.</i> (1996)
Pb ²⁺ (semen)		(Fertile)	
		37.0 ± 4.9 µg/dl*	
		(Infertile)	
	27.9 ± 2.5 µg/dl	45.0 ± 9.4 µg/dl*	Aribarg and Sukcharoen (1996)
	5.9 ± 2.7 µg/dl		Plechaty <i>et al.</i> (1977)
	16.5 ± 17.7 µg/dl	66.5 ± 83.6 µg/dl*	Xuezhi <i>et al.</i> (1992)
	11.0 ± 1.9 µg/dl	21.9 ± 2.0 µg/dl*	El-Zohairy <i>et al.</i> (1996)
Cd ²⁺ (blood)		(Fertile)	
		25.8 ± 3.2 µg/dl*	
		(Infertile)	
Cd ²⁺ (semen)	18.6 ± 3.3 µg/dl	38.3 ± 10.5 µg/dl*	Aribarg and Sukcharoen (1996)
	0.22 ± 0.21 µg/dl	0.16 ± 0.14 µg/dl	Xuezhi <i>et al.</i> (1992)
	0.04 ± 0.03 µg/dl	0.05 ± 0.05 µg/dl	Xuezhi <i>et al.</i> (1992)
	0.38 ± 0.64 µg/l	3.28 ± 0.40 µg/l*	Keck <i>et al.</i> (1995)

*Statistically significant increase, $P < 0.01$ – 0.0001 .

between Cd²⁺ and malignant transformation of prostate epithelial cells (Voeller *et al.*, 1991; Waalkes and Rehms, 1994).

The controversy concerning whether or not the fertility potential of Pb²⁺- and/or Cd²⁺-exposed men is decreased may not be resolved in the near future as the developmental risks, e.g. abortions, stillbirths, postnatal deaths, congenital malformations (Sallmen *et al.*, 1992; Xuezhi *et al.*, 1992), from occupational Pb²⁺ exposures appears to be of greater concern to employers than male reproductive effects (Paul and Kurtz, 1994). This appear also true of investigators studying the effects of environmental exposures (Sram *et al.*, 1996).

Sources of Pb²⁺ and Cd²⁺ found in the environment

Environmental discharges of Pb²⁺ are a major concern as Pb²⁺ wastes are tracked as hazardous materials. Pb²⁺ is probably the most common neurotoxin and suspected reproductive toxicant present in the environment at high concentrations as the result of the use of petroleum products (Hutton and Symon, 1986) and

demand for lead-acid batteries, the latter requiring mining, smelting and recycling of Pb²⁺ (Lave *et al.*, 1995). Combustion of fossil fuels (petroleum, coal) and municipal refuse also contribute to airborne Cd²⁺ pollution (Hutton and Symon, 1986; Ragan and Mast, 1990). As for Pb²⁺, regional concentrations of Cd²⁺ may be higher near industrial operations, such as smelters and battery and paint factories (Benin *et al.*, 1999). In addition, the public may be unwittingly exposed to Pb²⁺ or Cd²⁺ via contaminated food or paper (Wu *et al.*, 1995) and cosmetics and herbal folk remedies (Wong and Koh, 1986; Lockitch, 1993). The major risk factors for Pb²⁺ toxicity include age, nutrition (deficiencies in Ca²⁺, Fe²⁺/³⁺ and Zn²⁺), housing and socioeconomic status and, importantly, smoking (Goyer, 1993).

Cigarette smoking is an important confounding variable when considering the effects of both environmental Pb²⁺ and Cd²⁺ exposures on human health (Sram *et al.*, 1996). Cigarette smoke is a major source of airborne environmental Pb²⁺ and Cd²⁺ exposures. A single cigarette contains 0.6–2.0 µg Pb²⁺ (Chiba and Masironi, 1991) and 1–4.5 µg Cd²⁺ (Chia *et al.*, 1994b;

Saldivar *et al.*, 1991) and at least one tenth of the metal content of a cigarette is inhaled (Elinder *et al.*, 1985).

Examination of the effects of environmental exposure to metal ions

Agents which are reproductive toxicants in the workplace are found in human semen, e.g. Cd²⁺, dibromochloropropane, lithium, and pesticides (Gagnon, 1988). However, the role of increased seminal plasma trace metal concentrations in regulating reproductive function in the occupationally unexposed male is poorly understood. Geographic differences in the amount of naturally occurring Cd²⁺ have been correlated with incidence rates of prostate cancer (Garcia Sanchez *et al.*, 1992; Angwafo, 1998). In contrast, it has been difficult to identify an association between environmental metal exposures and male infertility. Since the desire for reproduction occurs only intermittently, environmental exposures that interfere with reproduction may go unnoticed (Mattison *et al.*, 1990).

Tobacco smoke is the best studied environmental exposure. Cigarette smoking impairs both female and male fecundity (Zenses, 1995; Hughes and Brennan, 1996; Curtis *et al.*, 1997). The decrease in fecundity in men who smoke cigarettes may be the result of reductions in semen volume, sperm density, sperm motility, normal sperm morphology and/or sperm viability (Saaranen *et al.*, 1987b; Radcliffe *et al.*, 1992; Chia *et al.*, 1994a,b; Vine *et al.*, 1994; Omu *et al.*, 1995; Sofikitis *et al.*, 1995; Vine, 1996; Merino *et al.*, 1998; Rubes *et al.*, 1998). An increase in sperm aneuploidy, e.g. Y disomy, has recently been noted (Rubes *et al.*, 1998). The latter is consistent with an increase in proportion of female offspring in men exposed to Pb²⁺ (Dickinson and Parker, 1994). The outcome of sperm function assays was significantly lower in men who smoke, compared with non-smoking controls (Sofikitis *et al.*, 1995). Cigarette smoke may also potentiate the effects of other suspected reproductive toxicants, e.g. caffeine (Bolumar *et al.*, 1997).

Although blood concentrations of Cd²⁺ and Pb²⁺ may be increased in smokers (Brockhaus *et al.*, 1983; Friberg and Vahter,

Table II. Comparison of Pb²⁺ and Cd²⁺ concentrations in blood and seminal plasma in men who smoke cigarettes and non-smoking control subjects

	Non-smokers	Smokers	Source
Pb ²⁺ (blood)	92.0 µg/l	99.2 µg/l	Brockhaus <i>et al.</i> (1983)
		7.09 µg/l	Chia <i>et al.</i> (1994b)
Pb ²⁺ (semen)		12.9 µg/l	Chia <i>et al.</i> (1994b)
	0.24–0.38 µg/l	0.24–0.34 µg/l	Oldereid <i>et al.</i> (1994)
		(<1 pack/day)	
		0.22–0.39 µg/l	
		(>1 pack/day)	
Cd ²⁺ (blood)	0.44 µg/l	1.85 µg/l*	Brockhaus <i>et al.</i> (1983)
	0.63 ± 0.44 µg/l	1.66 ± 0.93 µg/l	Moreau <i>et al.</i> (1983)
	0.24 ± 0.09 µg/l	0.31 ± 0.06 µg/l*	Saaranen <i>et al.</i> (1989)
		(<1 pack/day)	
		0.39 ± 0.05 µg/l*	
		(>1 pack/day)	
	0.84 µg/l	1.55 µg/l*	Chia <i>et al.</i> (1994b)
Cd ²⁺ (semen)	0.19 ± 0.21 µg/l	0.23 ± 0.17 µg/l	Saaranen <i>et al.</i> (1989)
		(<1 pack/day)	
		0.40 ± 0.40 µg/l	
		(>1 pack/day)	
	0.54 µg/l	0.86 µg/l	Chia <i>et al.</i> (1994b)
	0.14–0.32 µg/l	0.19–0.38 µg/l	Oldereid <i>et al.</i> (1994)
		(<1 pack/day)	
		0.29–0.49 µg/l*	
		(>1 pack/day)	

*Statistically significant increase, $P < 0.05$ – 0.001 .

Table III. Concentrations of Pb2+ and Cd2+ reported in blood and semen from fertile and infertile men

	Fertile	Infertile	Source
Pb2+ (blood)		6.5 ± 5.4 µg/l	Chia <i>et al.</i> (1992)
	16.9 ± 2.6 µg/dl	29.3 ± 3.8 µg/dl*	El-Zohairy <i>et al.</i> (1996)
Pb2+ (semen)	7.4 ± 0.7 µg/l		Pleban and Mei (1983)
	255 ± 123 µg/l	243 ± 179 µg/l	Umeyama <i>et al.</i> (1986)
	1.7 ± 1.0 µg/l	3.6 ± 3.2 µg/l*	Saaranen <i>et al.</i> (1987a)
	8 ± 1.1 µg/l	1.2 ± 1.2 µg/l	Abou-Shakra <i>et al.</i> (1989)
		(Oligospermic)	
		6 ± 4 µg/l	
		(Severely oligospermic)	
		1.0 ± 1.3 µg/l	
		(Azoospermic)	
	5.61 ± 0.53 µg/l	11.18 ± 0.62 µg/l*	Jockenhovel <i>et al.</i> (1990)
	7.7 ± 5.6 µg/l		Noack-Fuller <i>et al.</i> (1993)
	11.0 ± 1.9 µg/dl	19.6 ± 2.4 µg/dl*	El-Zohairy <i>et al.</i> (1996)
Cd2+ (blood)		1.35 ± 1.0 µg/l	Chia <i>et al.</i> (1992)
Cd2+ (semen)	0.48 ± 0.04 µg/l		Pleban and Mei (1983)
	5 ± 6 µg/l	13 ± 12 µg/l*	Umeyama <i>et al.</i> (1986)
	0.34 ± 0.16 µg/l		Noack-Fuller <i>et al.</i> (1993)
	0.38 ± 0.64 µg/l	0.43 ± 0.69 µg/l	Keck <i>et al.</i> (1995)

*Statistically significant increase, $P < 0.05$ – 0.005 .

1983; Moreau *et al.*, 1983; Xu *et al.*, 1993; see Table II), it is more likely that the negative effects of smoking on semen quality are a result of the observed trend for Cd2+ in tobacco smoke to accumulate in the genital tract of smokers (Saaranen *et al.*, 1989; Pacifici *et al.*, 1993; Oldereid *et al.*, 1994; Keck *et al.*, 1995; Omu *et al.*, 1995; see Table II). In contrast, seminal plasma Pb2+ concentrations appear unaffected by smoking status (see Table II).

Not only are smokers at risk. A positive correlation has been observed between the degree of passive exposure to cigarette smoke and the concentration of components of cigarette smoke in seminal plasma (Pacifici *et al.*, 1995).

Subfertile or infertile men may be more sensitive to the deleterious effects of cigarette smoke than fertile men (Zenses, 1995). Risk factors for unexplained dyspermia in infertile men include smoking (Parazzini *et al.*, 1993). Total sperm count and sperm motility are lower in oligozoospermic men who smoke than in oligozoospermic non-smokers (Rantala and Koskimies, 1987; Vine *et al.*, 1996). An acrosome reaction insufficiency was first suggested from the significant decrease in penetration of zona-free hamster oocytes by spermatozoa from infertile men who smoked cigarettes, compared with infertile men who were non-smokers (Close *et al.*, 1990). A reduced inducibility of the acrosome reaction in subfertile men who smoke has since been confirmed (El Mulla *et al.*, 1995). The latter is consistent with reports of increased 'round-headed' spermatozoa, which lack an acrosome

(Zamboni, 1987; Escalier, 1990), in the semen of smokers (Rubes *et al.*, 1998). Hormonal alterations may occur (Gerhard *et al.*, 1992). Anti-sperm antibody production is increased in smokers (Omu *et al.*, 1997). An improvement in semen quality has been noted when subfertile men quit smoking (Vine, 1996). The concept of infertility patients as forming 'susceptibility subgroups' is discussed in greater detail later in this review (see section on genetic determinants of susceptibility below).

Metal ion concentrations in fertile and infertile men

It has long been suggested that at least one half of the cases of human male infertility of unknown aetiology may be attributable to various environmental and occupational exposures (Steen and Pangkahila, 1984a,b; Gagnon, 1988). Therefore, infertility patients have often been used in attempts to investigate the role of environmental metals. Data from such studies may be skewed as a result of selection bias. Not all couples experiencing infertility seek medical intervention (Olsen *et al.*, 1998).

Semen from men presenting for fertility evaluations generally contain only low concentrations of Pb2+ and Cd2+ (Noack-Fuller *et al.*, 1993). Nevertheless, a considerable body of evidence indicates that decreased concentrations of Zn2+ in seminal plasma and semen occur in some forms of infertility (Marmar *et al.*, 1975; Umeyama *et al.*, 1986; Saaranen *et al.*, 1987a; Hurley *et al.*, 1997;

S.Benoff, A.Hershlag and I.R.Hurley, unpublished observations). In contrast, the majority of reports suggest that environmental exposures to Pb²⁺ do not significantly contribute to male infertility. In at least half the published studies, seminal plasma Pb²⁺ values have been reported to be similar in fertile and infertile men (Umeyama *et al.*, 1986; Abou-Shakra *et al.*, 1989; see Table III).

Many studies have failed to find any significant differences in the seminal plasma Cd²⁺ concentrations of fertile and infertile men (Keck *et al.*, 1995). In contrast, in other studies, seminal plasma Cd²⁺ concentrations were found to be increased in infertile, compared with fertile, men (Umeyama *et al.*, 1986; Saaranen *et al.*, 1987a; Jockenhovel *et al.*, 1990). In addition, increases in blood plasma Cd²⁺ concentrations in infertile men have been associated with teratozoospermia (Chia *et al.*, 1992) and an inverse correlation between blood plasma Cd²⁺/seminal plasma Cd²⁺ and sperm density and semen volume has been reported (Xu *et al.*, 1993). These conflicting findings are summarized in Table III.

Confounding variables

Confounding variables are likely to be responsible for many of the apparent contradictions between similar studies cited in the preceding section. Groups of environmental toxicologists have been convinced of this, and have proposed guidelines for controlling such variables (Tas *et al.*, 1996; Wyrobek *et al.*, 1997). Central to these strategies is use of multiple study endpoints (Perreault *et al.*, 1989), including biomarkers (Smith and Suk, 1994), to assess exposures to other toxicants which may produce or mask conditions induced by the chemical(s) under study. The extent of knowledge of a particular chemical's toxicology and its underlying molecular biology insofar as it relates to the outcome measure under study will determine the comprehensiveness of the list of confounding variables and endpoints.

Depending upon outcome measure, endpoints may include questionnaires to gather medical and reproductive histories, information on lifestyle factors, e.g. smoking, herbal therapy (Ondrizek *et al.*, 1999) and exposures to unrelated toxicants. Physical examinations of subjects are useful in revealing physical conditions affecting male fertility which may not be evident from questionnaires, including varicocele and small testicular size. A standardized classical semen analysis, computerized sperm morphometry and motility together with blood hormone concentrations are recommended both to characterize the effects of a particular exposure and to identify sub-populations, which presumably differ with respect to some confounding variable. Choice of control groups are also a major issue, e.g. see Favino *et al.* (1968) who used Pb²⁺-exposed men as controls for the effects of Cd²⁺ on reproductive hormones.

The state of knowledge of heavy metal confounders is relatively satisfactory. Concentrations of various metal ions have been assayed in the same tissues in animal models. This has revealed that transport and equilibrium concentrations of heavy metals can be altered by concentrations of other metals administered in the course of both acute and chronic exposure studies. For example, similarities of the ionic size and normal ionic charge between Cd²⁺, Ca²⁺ and Zn²⁺ with Pb²⁺ mean that both Cd²⁺ and Pb²⁺

can pass the gut/blood barrier by binding to Ca²⁺ and Zn²⁺ transport sites on the surface of intestinal villae (Goyer, 1997). High amounts of Ca²⁺ in diet are protective against both Cd²⁺ and Pb²⁺ poisoning. However, gaps remain. Measurements of effects of Pb²⁺ upon sperm fertility outcome measures, such as progesterone-induced acrosome loss (see section on ion channels and response to environmental metal exposures below), therefore require assessment of the inter-reacting metal confounders, Cd²⁺, Zn²⁺ and Ca²⁺ in the same samples. These measurements must be both accurate and sensitive, as different methods appear to produce discordant results (e.g. colorimetric versus graphite furnace atomic absorption spectroscopy; Fuentes *et al.*, 1981 and Jeyendran *et al.*, 1989 versus Canale *et al.*, 1986, Xu *et al.*, 1993 and Hurley *et al.*, 1997). Other confounding variables which affect the availability of metals may be addressed using conventional assays. These include transport and storage proteins, and components of diet. For example, high phosphate in the diet potentiates Pb²⁺ poisoning, by forming insoluble complexes with dietary Ca²⁺.

There is also a need for reliable molecular biomarkers to assess individual susceptibilities to heavy metal toxicity (Todd *et al.*, 1996). Individual variability could explain why, though blood Pb²⁺ concentrations can clearly be increased by environmental exposures (Sram *et al.*, 1996), population studies of effects of chronic Pb²⁺ exposure upon fertility-related outcome measures have not produced consensus. The goal therefore is to develop a group of markers similar to the genetically determined polymorphisms in enzymes susceptible to Pb²⁺ inhibition, which have been suggested to be useful for predicting renal toxicity (Doss *et al.*, 1982; Todd *et al.*, 1996). We suggest, however, that the most important confounding variable is the choice of the population(s) to be studied.

Genetic determinants of susceptibility

There is an increasing awareness in environmental toxicology for factors that predispose individuals to be sensitive to toxicant exposure (Anderson *et al.*, 1994). One of the primary factors leading to the existence of this susceptibility is genetic polymorphisms.

Somatic diseases

Susceptibility subgroups associated with genetic polymorphisms have been identified in a variety of somatic cell diseases, including sickle cell anaemia (Lu and Steinberg, 1996), tobacco-related lung cancer and gastric cancer (Hirvonen, 1995), breast or ovarian cancer (Suk *et al.*, 1996) and prostate cancer (Kantoff *et al.*, 1998). These polymorphisms may take the form of unstable simple DNA repeats (Panzer *et al.*, 1995) or functional enzyme polymorphisms with marked inter-individual variation (Caraco, 1998) that mediate oxidative metabolism and detoxification reactions, e.g. cytochrome P-450 (CYP), monamine oxidase, superoxide dismutase and glutathione S-transferase (GST) (Hirvonen, 1995; Raunio *et al.*, 1995; Watanabe, 1998).

Infertility

A large portion of male infertility is thought to have an underlying genetic basis, e.g. gonadotrophin-releasing hormone (GnRH) deficiency, spermatogenic failure and both obstructive (e.g.

congenital absence of the vas deferens) and non-obstructive azoospermia (e.g. associated with Y deletions) (Mak and Jarvi, 1996). An association between expression of certain human leukocyte antigen antigens (e.g. HLA-A28 and Bw40) and azoospermia has also been noted (Bisson *et al.*, 1976; Miura *et al.*, 1998). Studies on genetic factors regulating fertility/fecundity and response to environmental agents have been at best extremely limited. Recent studies suggest that inter-individual differences in expression of isoforms of certain enzymes underlie the response to airborne environmental toxicants (e.g. GST and *N*-acetyltransferase versus effects of polycyclic aromatic hydrocarbons; Sram *et al.*, 1996) and may contribute to the production of endometriosis (Baranova *et al.*, 1997) or an increase in spontaneous abortions (Wang *et al.*, 1998). In contrast, we do not know why occupational exposure to suspected reproductive toxicants appears to have a greater effect on male fecundity than do female occupational exposures (Spinelli *et al.*, 1997), or why men are more sensitive than women to environmental exposure to metals (Staessen *et al.*, 1991; Dickman *et al.*, 1998) or organic toxicants (Welch *et al.*, 1988; Schrader, 1997).

Clearly, there is much to learn about the factors which regulate susceptibility. We believe that analysis of infertility patients offers a unique opportunity to address these factors. Our data suggest that infertility patients may be predisposed to and may respond more strongly to environmental metal exposures than the population in general.

Ion channels and response to environmental metal exposures

Acrosome exocytosis induced by sperm–zona binding and required for sperm penetration through the zona pellucida involves sequential activation of sperm head delayed (outward) rectifier voltage-gated potassium (K^+) channel (VGKC) and L-type voltage-dependent Ca^{2+} (VDCC) ion channels (L-VDCC) (Benoff, 1999). Distinct binding sites for Pb^{2+} , Cd^{2+} and other metal ions appear to regulate ion permeation (Leinders *et al.*, 1992). Each type of channel is composed of multiple subunits. When expressed *in vitro*, α subunits possess all characteristics necessary to form channel pores (Catterall, 1995; Benoff, 1998).

The α subunit of the typical VGKC consists of six putative transmembrane segments designated S1–S6 (Catterall, 1995). Transmembrane segment S4 is the voltage sensor. The regions between transmembrane segments S5 and S6 contain the ion conducting pore and binding sites for channel blockers and metals ions (Shieh and Kirsch, 1994; Catterall, 1995; Christie, 1995; Doyle *et al.*, 1998). Different isoforms result from sequence changes in the ion conduction pore (Grissmer *et al.*, 1994). Thus, genetic polymorphisms in these ion channels can regulate resistance or susceptibility to environmental metal exposures.

The α -1 subunits of VDCC are related to those of VGKC, being derived from a common ancestor (Catterall, 1995). VDCC α -1 subunits are composed of four repeats (domains I–IV) of the basic structural unit of the VGKC α subunit (S1–S6) (Catterall, 1995; MacKinnon, 1995). Multiple L-VDCC α -1 isoforms (e.g. α -1A, α -1B, α -1C, α -1D, α -1E) are generated by transcription of different genes and by alternate splicing of the primary transcript from each gene (Benoff, 1998). The α -1C subunit forms the pore

of the L-VDCC expressed in mammalian testis and spermatozoa (Benoff, 1998,1999; Goodwin *et al.*, 1999a,b). This subunit contains the binding sites for a variety of pharmacological agents, including dihydropyridines which are prescribed for hypertension control and which can produce a reversible infertile state (Benoff *et al.*, 1994; Hershlag *et al.*, 1995). Ion selectivity (e.g. Ca^{2+} versus Na^+) is conferred by single amino acid changes in the extracellular mouth of the pore and/or the pore wall itself (Heinemann *et al.*, 1992). By analogy with Na^+ channels, sensitivity or resistance to metals ions will also be conferred by single amino acid substitutions (i.e. Backx *et al.*, 1992; Satin *et al.*, 1992).

Genetic polymorphisms in these ion channels expressed in somatic cells have been associated with the generation of diseases. For example, a study of CCG repeat polymorphisms in human brain cDNA (targeted to develop markers for neuro-psychiatric disorders including autism) amplified a partial clone of BCNG-1 (Kleiderlein *et al.*, 1998). BCNG-1 has been identified as encoding a voltage-gated potassium (K^+) channel (VGKC) with properties indistinguishable from brain pacemaker channels (Santoro *et al.*, 1998). If BCNG-1 shows the sensitivity to Pb^{2+} of other voltage-gated K^+ channels (Kiss and Osipenko, 1994), its polymorphic expression may provide a mechanism for the variability confounding neuro–psychiatric effects of paediatric Pb^{2+} exposure (Needleman, 1988).

With regard to VDCC, alternate splicing in the carboxy terminus of the α -1A subunit is linked to a severe form of human ataxia (Zhuchenko *et al.*, 1997). Sequence changes, including one in transmembrane segment IIIS2, of the α -1A result in familial hemiplegic migraine and episodic ataxia type-2 where patients exhibit migraine-like symptoms (Ophoff *et al.*, 1996). These findings are of particular importance given that we have identified multiple splice variants of the human α -1C subunit that are expressed in testis and spermatozoa. The alternatively spliced regions include the amino terminus and transmembrane segments IS6, IIIS2 and IVS3 (see section on varicocele-associated infertility below).

Potential models for metal effects mediated by ion channels

We now present evidence that the interaction between environmental exposure to Cd^{2+} and L-VDCC results in infertility in the presence of a varicocele. We also report that many cases of ‘unexplained’ infertility can be attributed to the interaction of VGKC and environmental Pb^{2+} exposures.

Varicocele-associated infertility

Varicocele-associated infertility is widespread: ~40% of males from infertile couples present with varicocele (Greenberg, 1977; Goldstein *et al.*, 1992; WHO, 1992; Schlesinger *et al.* 1994). The mechanism(s) underlying varicocele-associated infertility, however, are poorly characterized. The most widely accepted explanation for the pathophysiology of varicocele in male infertility is abnormally elevated testicular temperature due to impaired heat transfer by the scrotum and/or changes in testicular blood flow (Comhaire, 1991). Although men with varicocele exhibit higher mean scrotal temperatures, there is a large overlap with the range of scrotal temperatures in fertile men (Mieusset and

Bujan, 1995). More importantly, only 13% of men with varicocele are infertile (Kursh, 1987; Sylora and Pryor, 1994), and, although varicocele repair has been documented to reduce testicular temperature (Agger, 1971; Yamaguchi *et al.*, 1989), only 1/3 of infertile men with varicocele will experience a return of fecundity following varicocele correction (Schlesinger *et al.*, 1994). These findings suggest that varicocele may not be a primary cause of infertility and that it is the interaction of varicocele with other as yet unidentified factors that produces the infertile state (Peng *et al.*, 1990; Gentile and Cockett, 1991; Mieusset and Bujan, 1995).

We suggest that one such factor is the concentration of Cd²⁺ in the male reproductive tract (Benoff, 1997; Benoff *et al.*, 1997, 1998a). In animal varicocele models, exposure to components of cigarette smoke increases impairment of spermatogenesis. Although there is some controversy as to whether smoking reduces semen quality (Goverde *et al.*, 1995; Hughes and Brennan, 1996) or reduces fertility (Vine, 1996) in all men of reproductive age, it is generally agreed that men with marginal semen quality may be made infertile by smoking (see discussion in section on examination of the effects of environmental exposure to metal ions above). Oligozoospermia among men with varicocele who also smoke is: (i) 10-fold more frequent than among other men with varicocele and (ii) five-fold more frequent than among smokers without varicocele (Klaiber *et al.*, 1987).

Our data indicate that the concentration of Cd²⁺ in seminal plasma from infertile men with varicocele is markedly increased, compared with fertile men with varicocele (Figure 1) or men with other forms of infertility (e.g. see Figure 2). Recent findings indicate that Cd²⁺ concentrations are similarly increased in the testis of infertile men with varicocele (Figure 1; Benoff *et al.*, 1999b). After varicocele repair, pregnancy by coitus was observed only in those men whose Cd²⁺ concentrations were at the lower end of the range (Benoff *et al.*, 1998a). These findings suggest that measurements of seminal plasma Cd²⁺ concentrations may serve as a biomarker for the outcome of varicocele repair.

Men with secondary infertility and varicocele present with seminal plasma Cd²⁺ concentrations intermediate between those of fertile men and men with primary varicocele-associated infertility (Benoff *et al.*, 1999b), suggesting that the deleterious effects of Cd²⁺ are time and concentration dependent. Blood plasma Cd²⁺ concentrations did not, however, differ significantly between infertile and fertile men with varicocele (Fig. 1). These data emphasize that the choice of variable to be analysed influences whether or not an association is detected between an environmental exposure and production of the infertile state.

It is known that infertile men with varicocele are typically oligozoospermic. The sperm populations from these men have increased tapering head forms ('stress pattern'), reduced zona pellucida binding ability and an acrosome reaction insufficiency (Rogers *et al.*, 1985; Naftulin *et al.*, 1991; Vigil *et al.*, 1994; Benoff *et al.*, 1995a, 1997; El Mulla *et al.*, 1995; Benoff, 1997). These defects can clearly result from an interaction between the observed increase in testicular Cd²⁺ and actin. Cd²⁺ exposure results in depolymerization of actin filaments and a net decrease in actin monomer protein (Wang *et al.*, 1996). In somatic cells, actin loss increases apoptosis (Russo *et al.*, 1982; Tsukidate *et al.*,




Figure 1. Comparison of the cadmium (Cd²⁺) content in blood plasma (BP), seminal plasma (SP) and testis biopsies (TB) of infertile men with varicocele (VAI) and men without varicocele (controls). Cd²⁺ concentrations were determined by graphite furnace atomic absorption spectroscopy (Benoff, 1997; Benoff *et al.*, 1997, 1998a, 1999b; Hurley *et al.*, 1997). The bars in the figure indicate the BP and SP Cd²⁺ content in µg/l and in testis biopsies (obtained by percutaneous needle aspiration; Marmar, 1998) in ng/mg dry weight. The concentrations of Cd²⁺ in blood plasma from male partners without varicocele from couples seeking fertility evaluation ($n = 66$) and from VAI males ($n = 14$) were similar. In contrast, SP Cd²⁺ concentrations were significantly elevated in men with VAI ($n = 14$), compared with fertile men with varicocele ($n = 4$) (a, $P < 0.0001$; t -test). Similarly, TB Cd²⁺ levels were higher in men with VAI [$n = 8$; Johnsen score (Johnsen, 1970) = 8–9] than in TB from men with non-obstructive azoospermia ($n = 4$; Johnsen score = 6) (b, $P < 0.03$; t -test).

1993) (which could lead to oligozoospermia in infertile men with varicocele) and decreases the ability to undergo exocytosis (Muallem *et al.*, 1995) (which requires both actin polymerization and depolymerization, as in the acrosome reaction; Rogers *et al.*, 1989; Spungin *et al.*, 1995; Benoff *et al.*, 1996; Benoff, 1999).

Preliminary examination of testis biopsies from infertile men with varicocele suggests that this indeed is the case. Actin immunoreactivity was decreased and apoptosis, which normally occurs at low concentrations during spermatogenesis (Furuchi *et al.*, 1996; Tesarik *et al.*, 1998) was increased, compared with biopsies from men without varicocele (Benoff *et al.*, 1999b). These data are consistent with findings in animal models wherein in-vivo administration of Cd²⁺ induced apoptosis in testicular tissue (Xu *et al.*, 1996; Yan *et al.*, 1997).

In-vitro modelling experiments support our in-vivo findings (Benoff, 1997; Benoff *et al.*, 1997). Cd²⁺ exposure changed fertile donor sperm protein expression (e.g. loss of sperm head actin, reduced percentages of spermatozoa expressing surface mannose and non-nuclear progesterone receptors) and function (e.g. acrosome reaction insufficiency) to mimic varicocele in a dose-response fashion, albeit at Cd²⁺ concentrations >30-fold higher than in seminal plasma from infertile men with varicocele. After fertile donor spermatozoa were briefly exposed to higher temperatures, lower Cd²⁺ concentrations produced more drastic sperm function changes, indicating that temperature and Cd²⁺ act synergistically, as in cryptorchid animal models (Chatterjee and Ray, 1972; Fende and Niewenhuys, 1977).

Increased testicular Cd²⁺ concentrations are probably derived from the increased transvascular fluid exchange which occurs with varicocele (Shafik and Bedeir, 1980; Sweeney *et al.*, 1995).

This means that there is the potential for more of the Cd²⁺ in serum to enter the testis. As there is no active pump to remove this Cd²⁺ (Gunn *et al.*, 1961), testicular Cd²⁺ concentrations would be elevated over time. Results from autometallography (Benoff, 1998,1999; Benoff *et al.*, 1999b) indicate that this Cd²⁺ can enter spermatogenic cells via L-VDCC (Kiss and Osipenko, 1994; Atar *et al.*, 1995), which are expressed in all cells within the germinal epithelium (Goodwin *et al.*, 1998). The multiple isoforms of the testis-specific L-VDCC which have been identified result from alternate splicing of a primary transcript from a single gene (Benoff, 1998). Preliminary findings suggest that each man expresses only one L-VDCC splice variant in his ejaculated spermatozoa (Goodwin *et al.*, 1999a,b). The Cd²⁺ sensitivity or resistance of L-VDCC is conferred by the amino acid sequence within a short segment of the protein (domain I, transmembrane segment S6; e.g. Backx *et al.*, 1992). This region is alternatively spliced in the testis-specific L-VDCC and a series of amino acid substitutions within this region have also been observed (Benoff, 1999). Thus, our working hypothesis is that infertility in these men results from the interaction of three factors: (i) the presence of a varicocele; (ii) an increase in testicular Cd²⁺ concentrations; and (iii) the expression of particular L-VDCC isoforms.

Lead and unexplained infertility

We have been prospectively evaluating men from infertile couples undergoing their first IVF cycle in order to identify biomarkers which predict fertilization outcome and which are not related to sperm count, motility, viability or morphology (Benoff *et al.*, 1999a). An unexpected finding was that >40% of these males, who were not exposed to Pb²⁺ in their places of work and who did not smoke cigarettes, exhibited blood and seminal plasma Pb²⁺ concentrations that were above the permissible limit in men occupationally exposed to Pb²⁺ (Benoff *et al.*, 1998b). Both blood and seminal plasma Pb²⁺ concentrations in these men were inversely correlated with the rate of fertilization of metaphase II oocytes in IVF (Benoff *et al.*, 1998b; Benoff, 1999; see Figure 2). The negative effect of increased Pb²⁺ on IVF outcome was attributable to altered sperm function, i.e. increased spontaneous acrosome loss and a decreased ability to undergo an acrosome reaction induced by exposure of capacitated spermatozoa to progesterone or model zona ligands containing mannose. The progesterone-stimulated acrosome reaction was the parameter most severely affected by Pb²⁺. Mannose receptors and non-nuclear progesterone receptors are co-expressed on a subpopulation of motile human spermatozoa (Benoff *et al.*, 1995c) and the response to progesterone determines whether or not human spermatozoa undergo an acrosome reaction after zona binding and fertilize human oocytes (Jacob *et al.*, 1998c).

Results from cDNA cloning and in-vitro modelling studies employing K⁺ channel inhibitors provided evidence that the effects of progesterone on the human sperm acrosome reaction were mediated by a delayed rectifier VGKC (Jacob *et al.*, 1997a,b, 1998a,b) and that this VGKC is upstream of the L-VDCC which regulates the sustained Ca²⁺ influx required for the AR (Benoff, 1999). K⁺ channels in somatic tissues can be inhibited by Pb²⁺ (Kiss and Osipenko, 1994) and environmental Pb²⁺ exposures have been reported to alter K⁺ transport by red blood cells (Hajem *et al.*, 1990).




Figure 2. Comparison of the concentrations of zinc (Zn²⁺), cadmium (Cd²⁺) and lead (Pb²⁺) in seminal plasma from men exhibiting high rates of fertilization following conventional IVF and men exhibiting reduced or failed fertilization. In a prospective analysis, we evaluated the concentrations of metals in seminal plasma (SP) from men from consecutive couples undergoing their first cycle of conventional IVF (*n* = 96). The number of spermatozoa used in each inseminate was normalized on the basis of acrosomal morphology, so that at least 25 000 spermatozoa with an oval head with a well-defined acrosome covering >50% of the head were inseminated per oocyte cultured in 1 ml of medium (Benoff *et al.*, 1999a). The mean fertilization for this study population was 85.2 ± 11.1% (Jacob *et al.*, 1998c; Benoff *et al.*, 1999a). Cases of 'reduced' fertilization were defined as exhibiting fertilization rates >2 SD below the mean, e.g. <63%. Cases of 'high' fertilization were defined as exhibiting fertilization rates >63%. SP metal concentrations were determined by graphite furnace atomic absorption spectroscopy (Benoff, 1997; Benoff *et al.*, 1997, 1998b; Hurley *et al.*, 1997; S.Benoff and I.R.Hurley, unpublished). SP Zn²⁺ concentrations are presented as millimolar concentrations while SP Cd²⁺ and SP Pb²⁺ concentrations are reported as µg/l. Although SP Cd²⁺ did not differ between the two groups, SP Zn²⁺ was markedly decreased (a, *P* < 0.0001; *t*-test) and SP Pb²⁺ was significantly increased (b, *P* < 0.0002; *t*-test) in cases of reduced fertilization, compared with cases with high fertilization rates.

Sperm surface VGKC were mapped using a biotinylated charybdotoxin (CBTx) probe (Figure 3a,b). Of spermatozoa from fertile donors ~38% exhibited head-directed CBTx binding. Sites of exogenous Pb²⁺ entry into the human sperm head were similarly localized by autometallography (Figure 3c,d). The same percentages of spermatozoa that bound CBTx also bound Pb²⁺, providing strong evidence that a VGKC capable of transporting Pb²⁺ is expressed on the human sperm head. Preliminary results from in-vitro modelling studies confirmed these findings (Figure 4). CBTx strongly inhibited the increase in spontaneous acrosome loss which normally results from Pb²⁺ exposure. These data are consistent with findings in animal models that in-vivo Pb²⁺ exposure causes premature acrosome reactions (Johansson, 1989) and provide strong support for our hypothesis that Pb²⁺ action to reduce human sperm fertilizing potential is mediated by a sperm head VGKC.

Analysis of the effects of the K⁺ channel inhibitors CBTx, tetraethylammonium ion (TEA), and 4-aminopyridine (4AP) on progesterone-stimulated acrosome loss by fertile donor spermatozoa provided evidence for inter- male variability in VGKC isoform expression (Benoff, 1999). The existence of multiple VGKC isoforms would help to explain why two out of 53 men who exhibited high rates of fertilization had seminal plasma Pb²⁺ concentrations that were >10 µg/dl (the upper limit of the range of blood Pb²⁺ concentrations in unexposed men) and three




Figure 3. Voltage-gated potassium channels and sites of exogenous lead entry co-localize on the human sperm head surface. Specimens were viewed at $\times 400$ original magnification with an Olympus microscope (Olympus Corp., Lake Success, NY, USA) and photographed at $\times 600$ on 35 mm/400 ASA black and white film (Eastman Kodak Co, Rochester, NY, USA) with automatic exposure for phase-contrast images (a, c, d) and 50 s exposure for the epifluorescence image (b). (a, b) A minimally biotinylated charybdotoxin probe was prepared by modifying the protocols supplied with commercial reagents (Pierce, Rockford, IL; Bio Rad Laboratories, Hercules, CA, USA) (A.Jacob, I.R. Hurley and S.Benoff, unpublished observations). Motile spermatozoa from fertile donors were surface labelled with the biotinylated charybdotoxin and sperm-bound charybdotoxin was visualized by indirect immunocytochemistry using polyclonal antibodies to biotin prepared in rabbit (Enzo Diagnostics, New York, NY, USA). (a) Paired phase-contrast and (b) epifluorescence images are shown. The arrow in the figure points to charybdotoxin binding over the entire sperm head surface. Scale bar = 12.5 μm . (c, d) Motile spermatozoa from fertile donors were incubated overnight in capacitation media supplemented with 2.5 $\mu\text{mol/l}$ lead (Pb^{2+}). Sperm-bound Pb^{2+} as localized by autometallography (Stoltenberg *et al.*, 1997a,b). Pb^{2+} bound over the entire sperm head is visualized as black silver deposits (d, arrow) after conversion of Pb^{2+} to Pb sulphide and physical development with silver lactate. Black silver deposits are not detected in control aliquots (c) exposed to Pb^{2+} but not treated with sodium sulphide prior to physical development. Scale bar = 12.5 μm .

out of 21 men who exhibited reduced or failed fertilization had Pb^{2+} concentrations $<10 \mu\text{g/dl}$. Further studies are now in progress to identify the specific structural variations in the human sperm VGKC α subunit which regulate sensitivity to Pb^{2+} .

Conclusions

We are left with a puzzle. The clinical data presented above show that two groups of infertile men have abnormally high metal concentrations in seminal plasma, without any obvious recent high exposure to these toxicants. The in-vitro studies and the molecular genetics provide, in outline, a plausible mechanism showing how these toxicants could produce the characteristic clinical symptoms of each type of infertility. However, this does not explain the aetiology of high metal concentrations. Both Cd^{2+} (characteristic of the varicocele-related infertile group) and Pb^{2+} (high spontaneous acrosome reaction group) are metals with a low clearance rate. It is possible that members of both groups had some unrecorded high exposure to these metals in the past, and their blood and seminal plasma concentrations are equilibrium concentrations reflective of normal clearance. This is hard to credit, as an exposure large enough to give the concentrations we observed after an uncertain number of years of the initial exposure is likely to have exhibited clinically significant symptoms at the time of that exposure. Another possibility is that the individuals with high metals concentrations have some other condition which




Figure 4. Opposing effects of lead (Pb^{2+}) and charybdotoxin (CBTx) on the spontaneous acrosome reaction. Motile sperm populations from fertile donors ($n = 5$) were divided and exposed to 50 nmol/l CBTx or 2.5 $\mu\text{mol/l}$ Pb^{2+} , alone or in combination, and compared with aliquots not exposed to either agent (control). Spontaneous acrosome loss was assessed following reaction with rhodamine-labelled *Pisum sativum* agglutinin, which binds to acrosome content (Benoff *et al.*, 1995b). Pb^{2+} exposure significantly increased acrosome loss ($P < 0.007$). CBTx decreased both the level of control acrosome loss ($P < 0.005$) and that observed following Pb^{2+} exposure ($P < 0.006$). The percentages of spermatozoa undergoing an acrosome reaction in the presence of both Pb^{2+} and CBTx were similar to that in the presence of CBTx alone (not significant).

predisposes them to accumulate high amounts of toxicant from exposures no higher than those most of us encounter. This

condition may be dietary, related to intake of some other metals or minerals which alter the bio-availability of toxicants. Alternatively, these conditions may be genetic, involving isoforms of proteins involved in transport, membrane passage or storage of metals. If genetic predisposition does indeed create susceptibility subgroups with significantly lower tolerance of toxicant exposures, it may be necessary to reconsider the standards embodied in current legislation.

Acknowledgements

The authors thank Avner Hershlag, Gerald M. Scholl and Terry Paine, for contribution of clinical IVF results, Joel L. Marmar and Bruce R. Gilbert, for contribution of specimens from infertile men with varicocele, Sally D. Perreault and Mary Ann Butler, for stimulating discussions and assistance in identifying pertinent background literature, Meredin Stoltenberg, for detailed protocols for autometallography, Barbara Napolitano, for statistical consultations, George W. Cooper for critical reading of the manuscript, and Stephanie Canaras, for expert technical assistance. This work was supported by Grant No. ES 06100 to S.B. with a research supplement for Underrepresented Minority Individuals in Postdoctoral Training to A.J. from the National Institute of Environmental Health Sciences, National Institutes of Health, Bethesda, Maryland and by Grant No. OH 03584 to A.J. from the National Institute for Occupational Safety and Health of the Centers for Disease Control and Prevention, Cincinnati, OH, USA.

References

- Abou-Shakra, F.R., Ward, N.I. and Everard, D.M. (1989) The role of trace elements in male infertility. *Fertil. Steril.*, **52**, 307–310.
- Adamopoulos, D.A., Pappa, A., Nicopoulou, S. *et al.* (1996) Seminal volume and total sperm number trends in men attending subfertility clinics in the Greater Athens area during the period 1977–1993. *Hum. Reprod.*, **9**, 1936–1941.
- Agger P. (1971) Scrotal and testicular temperature: its relation to sperm count before and after operation for varicocele. *Fertil. Steril.*, **22**, 286–296.
- Anderson, S., Sadinski, W., Shugart, L. *et al.* (1994) Genetic and molecular ecotoxicology: a research framework. *Environ. Health Perspect.*, **102** (Suppl. 12), 3–8.
- Angwafo, F.f. (1998) Migration and prostate cancer: an international perspective. *J. Natl Med. Assoc.*, **90** (Suppl. 11), S720–S723.
- Aribarg, A. and Sukcharoen, N. (1996) Effects of occupational lead exposure on spermatogenesis. *J. Med. Assoc. Thailand*, **79**, 91–97.
- Assennato, G., Paci, G., Baser, M.E. *et al.* (1986) Sperm count suppression without endocrine dysfunction in lead-exposed men. *Arch. Environ. Health*, **41**, 387–390.
- Atar, D., Backx, P.H., Appel, M.M. *et al.* (1995) Excitation–transcription coupling mediated by zinc influx through voltage-dependent calcium channels. *J. Biol. Chem.*, **270**, 2473–2477.
- Auger, J., Kunstmann, J.M., Czyglik, F. *et al.* (1995) Decline in semen quality among fertile men in Paris during the past 20 years. *N. Engl J. Med.*, **332**, 281–285.
- Backx, P.H., Yue, D.T., Lawrence, J.H. *et al.* (1992) Molecular localization of an ion-binding site within the pore of mammalian sodium channels. *Science*, **257**, 248–251.
- Baranova, H., Bothorishvilli, R., Canis, M. *et al.* (1997) Glutathione S-transferase M1 gene polymorphism and susceptibility to endometriosis in a French population. *Mol. Hum. Reprod.*, **3**, 775–780.
- Becker, S. and Berhane, K. (1997) A meta-analysis of 61 sperm count studies revisited. *Fertil. Steril.*, **67**, 1103–1108.
- Benin, A.L., Sargent, J.D., Dalton, M. *et al.* (1999) High concentrations of heavy metals in neighborhoods near ore smelters in Northern Mexico. *Environ. Health Perspect.*, **107**, 279–284.
- Benoff, S. (1997) Environmental toxins and varicocele. *Assist. Reprod. Technol. Androl.*, **9**, 261–284.
- Benoff, S. (1998) Voltage-dependent calcium channels in spermatozoa. *Frontiers in Bioscience*, **3**, d1220–d1240.
- Benoff, S. (1999) Receptors and channels regulating acrosome reactions. *Hum. Fertil.*, **2**, 42–55.
- Benoff, S., Cooper, G.W., Hurley, I. *et al.* (1994) The effect of calcium ion channel blockers on sperm fertilization potential. *Fertil. Steril.*, **62**, 606–617.
- Benoff, S., Barcia, M., Hurley, I.R. *et al.* (1995a) Varicocele-related infertility: altered expression of sperm receptor proteins with myosin motors. *J. Soc. Gynecol. Invest.*, **2**, 376.
- Benoff, S., Hurley, I.R., Mandel, F.S. *et al.* (1995b) Reproductive hazards assay: the effect of acute exposure to metal ions on human sperm mannose receptor expression. [Abstr. no. O-001] In *Scientific Program and Abstracts of the 51st Annual Meeting of the American Society for Reproductive Medicine*, p. S1.
- Benoff, S., Rushbrook, J.I., Hurley, I.R. *et al.* (1995c) Co-expression of mannose-ligand and non-nuclear progesterone receptors identifies an acrosome reaction-inducible subpopulation in fertile donor sperm. *Am. J. Reprod. Immunol.*, **34**, 100–115.
- Benoff, S., Barcia, M., Hurley, I.R. *et al.* (1996) Classification of male factor infertility relevant to in-vitro fertilization strategies using mannose ligands, acrosome status and anti-cytoskeletal antibodies. *Hum. Reprod.*, **11**, 1905–1918.
- Benoff, S., Hurley, I.R., Barcia, M. *et al.* (1997) A potential role for cadmium in the etiology of varicocele-associated infertility. *Fertil. Steril.*, **67**, 336–347.
- Benoff, S., Hurley, I.R., Cooper, G.W. *et al.* (1998a) Human sperm mannose receptor expression (MR) and seminal plasma (SP) metal levels: potential biomarkers for the outcome of varicocele repair. [Abstr. no. O-193] In *Scientific Program and Abstracts of the 54th Annual Meeting of the American Society for Reproductive Medicine*, p. S72.
- Benoff, S., Jacob, A., Mandel, F.S. *et al.* (1998b) Environmental lead (Pb2+) exposures, the acrosome reaction (AR) and human male infertility. [Abstr. no. 7] In *Program and Abstracts of the 23rd Annual Meeting of the American Society of Andrology*, p. 26.
- Benoff, S., Cooper, G.W., Paine, T. *et al.* (1999a) Numerical dose-compensated in vitro fertilization inseminations yield high fertilization and pregnancy rates. *Fertil. Steril.*, **71**, 1019–1028.
- Benoff, S., Hurley, I.R., Jacob, A. *et al.* (1999b) Increased testicular cadmium (Cd2+) levels contribute to infertility with varicocele. [Abstr. no. O-64] In *Scientific Program and Abstracts of the 55th Annual Meeting of the American Society for Reproductive Medicine*, p. S25.
- Bisson, J.P., David, G., Kolevski, P. *et al.* (1976) *HLA and Disease*. INSERM, Paris, pp.1–242.
- Bolumar, F., Olsen, J., Rebagliato, M. *et al.* (1997) Caffeine intake and delayed conception: a European multicenter study on infertility and subfecundity. *Am. J. Epidemiol.*, **145**, 324–334.
- Bonde, J.P. (1990a) Semen quality and sex hormones among mild steel and stainless steel welders: a cross sectional study. *Br. J. Indust. Med.*, **47**, 508–515.
- Bonde, J.P. (1990b) Semen quality in welders before and after three weeks of non-exposure. *Br. J. Indust. Med.*, **47**, 515–518.
- Bonde, J.E. (1993) The risk of male subfecundity attributable to welding of metals. *Int. J. Androl.*, **16** (Suppl 1), 1–23.
- Brockhaus, A., Freier, I., Ewers, U. *et al.* (1983) Levels of cadmium and lead in blood in relation to smoking, sex, occupation, and other factors in an adult population of the FRG. *Int. Arch. Occup. Environ. Health*, **52**, 167–175.
- Brys, M., Nawrocka, A.D., Miekos, E. *et al.* (1997) Zinc and cadmium analysis in human prostate neoplasms. *Biol. Trace Elem. Res.*, **59**, 145–152.
- Buchancova, J., Knizkova, M., Hyllova, D. *et al.* (1994) Content of the selected trace elements (Al, As, Cd, Cu, Fe, Hg, Mn, Ni, Pb, Zn) in blood, urine and hair of blood donors without occupational exposure to these metals. *Cent. Eur. J. Public Health*, **2**, 82–87.
- Canale, D., Bartelloni, M., Negroni, A. *et al.* (1986) Zinc in human semen. *Int. J. Androl.*, **9**, 477–480.
- Caraco, Y. (1998) Genetic determinants of drug responsiveness and drug interactions. *Ther. Drug Monit.*, **20**, 517–524.
- Carlsen, E., Giwercman, A., Keiding, N. *et al.* (1992) Evidence for decreasing quality of semen during the past 50 years. *Br. Med. J.*, **305**, 609–613.
- Catterall, W.A. (1995) Structure and function of voltage-gated ion channels. *Ann. Rev. Biochem.*, **64**, 493–531.

- Chatterjee, A. and Ray, P. (1972) An early differential effect of cadmium on the scrotal and contralateral cryptorchid testes of the rat. *J. Reprod. Fertil.*, **30**, 297–300.
- Chia, S.-E., Ong, C.N., Lee, S.T. *et al.* (1992) Blood concentrations of lead, cadmium, mercury, zinc, and copper and human semen parameters. *Arch. Androl.*, **29**, 177–183.
- Chia, S.E., Ong, C.N. and Tsakok, F.M. (1994a) Effects of cigarette smoking on human semen quality. *Arch. Androl.*, **33**, 163–168.
- Chia, S.E., Xu, B., Ong, C.N. *et al.* (1994b) Effect of cadmium and cigarette smoking on human semen quality. *Int. J. Fertil. Menopausal Stud.*, **39**, 292–298.
- Chiba, M. and Masironi, R. (1991) *Toxic and Trace Elements in Tobacco*. World Health Organization Report, Geneva, Switzerland.
- Christie, M.J. (1995) Molecular and functional diversity of K⁺ channels. *Clin. Exp. Pharmacol. Physiol.*, **22**, 944–951.
- Close, C.E., Roberts, P.L. and Berger, R.E. (1990) Cigarettes, alcohol and marijuana are related to pyospermia in infertile men. *J. Urol.*, **144**, 900–903.
- Comhaire, F. (1991) The pathogenesis of epididymo–testicular dysfunction in varicocele: factors other than temperature. *Adv. Exp. Biol. Med.*, **286**, 281–285.
- Coste, J., Mandereau, L., Pessione, F. *et al.* (1991) Lead-exposed workman and fertility: a cohort study on 354 subjects. *Eur. J. Epidemiol.*, **7**, 154–158.
- Curtis, K.M., Savitz, D.A. and Arbuckle, T.E. (1997) Effects of cigarette smoking, caffeine consumption, and alcohol intake on fecundability. *Am. J. Epidemiol.*, **146**, 32–41.
- Danielsson, B.R.G., Dencker, L., Lindgren, A. *et al.* (1984) Accumulation of toxic metals in male reproductive organs. *Arch. Toxicol.*, (Suppl. 7), 177–180.
- Dickinson, H. and Parker, L. (1994) Do alcohol and lead change the sex ratio. *J. Theor. Biol.*, **169**, 313–315.
- Dickman, M.D., Leung, C.K.M. and Leong, M.K.H. (1998) Hong Kong male subfertility links to mercury in human hair and fish. *Sci. Total Environ.*, **214**, 165–174.
- Doss, M., Becker, U., Sixel, F. *et al.* (1982) Persistent protoporphyria in hereditary porphobilinogen synthase (delta-aminolevulinic acid dehydratase) deficiency under low lead exposure. *Klin. Wochenschr.*, **60**, 599–606.
- Dougherty, R.C., Whitaker, M.J., Tany, S.Y. *et al.* (1986) Sperm density and toxic substances: a potential key to environmental health hazards. In McKinney, J.D. (ed.), *The Chemistry of Environmental Agents as Potential Human Hazards*. Ann Arbor Science Publications, Ann Arbor, MI, USA, pp. 263–77.
- Doyle, D.A., Cabral, J.M., Pfuetzner, R.A. *et al.* (1998) The structure of the potassium channel: molecular basis of K⁺ conduction and selectivity. *Science*, **280**, 69–77.
- Elghany, N.A., Schumacher, M.C., Slatery, M.L. *et al.* (1990) Occupation, cadmium exposure, and prostate cancer. *Epidemiology*, **1**, 107–115.
- Elinder, C.G., Kjellstrom, T., Hogstedt, C. *et al.* (1985) Cancer mortality of cadmium workers. *Br. J. Ind. Med.*, **42**, 651–655.
- El Mulla, K.F., Kohn, F.M., Beheiry, A.H. *et al.* (1995) The effect of smoking and varicocele on human sperm acrosin activity and acrosome reaction. *Hum. Reprod.*, **10**, 3190–3194.
- El-Zohairy, E.A., Youssef, A.F., Abul-Nasr, S.M. *et al.* (1996) Reproductive hazards of lead exposure among urban Egyptian men. *Reprod. Toxicol.*, **10**, 145–151.
- Escalier, D. (1990) Failure of differentiation of the nuclear–perinuclear complex in the round-headed human spermatozoa. *Int. J. Dev. Biol.*, **34**, 287–297.
- Favino, A., Candura, F., Chiappino, G. *et al.* (1968) Study on the androgen function of men exposed to cadmium. *Med. Lav.*, **59**, 105–110.
- Fende, P.L. and Niewenhuis, R. (1977) An electronic microscopic study of the effects of cadmium chloride on cryptorchid testes of the rat. *Biol. Reprod.*, **16**, 298–305.
- Feroli, A., Apostoli, P. and Romeo, L. (1989) Alteration of steroid hormone sulfation and D-glucuronic acid excretion in lead workers. *Biol. Trace Elem. Res.*, **21**, 289–294.
- Fisch, H. and Goluboff, E.T. (1996) Geographic variations in sperm counts: a potential cause of bias in studies of semen quality. *Fertil. Steril.*, **65**, 1044–1046.
- Friberg, L. and Vahter, M. (1983) Assessment of exposure to lead and cadmium, through biological monitoring: results of a UNEP/WHO global study. *Environ. Res.*, **30**, 95–128.
- Fuentes, J., Miro, J. and Riera, J. (1981) Simple colorimetric method for seminal plasma zinc assay. *Andrologia*, **14**, 322–327.
- Furuchi, T., Masuko, K., Nishimune, Y. *et al.* (1996) Inhibition of testicular germ cell apoptosis and differentiation in mice misexpressing Bcl-2 in spermatogonia. *Development*, **122**, 1703–1709.
- Gagnon, C. (1988) The role of environmental toxins in unexplained male infertility. *Sem. Reprod. Endocrinol.*, **6**, 369–376.
- Garcia Sanchez, A., Antona, J.F. and Urrutia, M. (1992) Geochemical prospection of cadmium in a high incidence area of prostate cancer, Sierra de Gata, Salamanca, Spain. *Sci. Total Environ.*, **116**, 243–251.
- Gennart, J.-P., Bernard, A. and Lauwerys, R. (1992a) Assessment of thyroid, testes, kidney and autonomic nervous system function in lead-exposed workers. *Int. Arch. Occup. Environ. Health*, **64**, 49–57.
- Gennart, J.P., Buchet, J.P., Roels, H. *et al.* (1992b) Fertility of male workers exposed to cadmium, lead or manganese. *Am. J. Epidemiol.*, **135**, 1208–1219.
- Gentile, D.P. and Cockett, A.T.K. (1991) Varicocele: effect on fertility. *Adv. Exp. Biol. Med.*, **286**, 289–293.
- Gerhard, I., Lenhard, K., Eggert-Kruse, W. *et al.* (1992) Clinical data which influence semen parameters in infertile men. *Hum. Reprod.*, **7**, 830–837.
- Goldstein, M., Gilbert, B., Dicker, A. *et al.* (1992) Microsurgical inguinal varicocele with delivery of the testis: an artery and lymphatic sparing technique. *J. Urol.*, **148**, 1808–1811.
- Goodwin, L.O., Leeds, N.B., Hurley, I.R. *et al.* (1998) Alternative splicing of exons of the alpha subunit of the rat testis L-type voltage-dependent calcium channel generates germ line-specific dihydropyridine binding sites. *Mol. Hum. Reprod.*, **4**, 215–226.
- Goodwin, L.O., Karabinus, D.S., Pergolizzi, R.G. *et al.* (1999a) L-type voltage-dependent calcium (Ca²⁺) channel (VDCC) alpha subunit transcripts from ejaculated sperm: inter-male variability in isoform expression. [Abstr. no. 10] In *Program and Abstracts of the 24th Annual Meeting of the American Society of Andrology*, p. 28.
- Goodwin, L.O., Leeds, N.B., Karabinus, D.S. *et al.* (1999b) Variability in L-type voltage dependent calcium channel (L-VDCC) sequence in human testis and sperm. [Abstr. no. O-62] In *Scientific Program and Abstracts of the 55th Annual Meeting of the American Society for Reproductive Medicine*, p. S24.
- Goverde, H.J., Dekker, H.S., Janssen, H.J. *et al.* (1995) Semen quality and frequency of smoking and alcohol consumption—an explorative study. *Int. J. Fertil. Menopausal Stud.*, **40**, 135–138.
- Goyer, R.A. (1993) Lead toxicity: current concepts. *Environ. Health Perspect.*, **100**, 177–187.
- Goyer, R.A. (1997) Toxic and essential metal interactions. *Ann. Rev. Nutr.*, **17**, 37–50.
- Greenberg, S.H. (1977) Varicocele and male fertility. *Fertil. Steril.*, **28**, 699–706.
- Grissmer, S., Nguyen, A.N., Aiyar, J. *et al.* (1994) Pharmacological characterization of five cloned voltage-gated K⁺ channels, Types Kv1.1, 1.2, 1.3, 1.5, and 3.1, stably expressed in mammalian cell lines. *Mol. Pharmacol.*, **45**, 1227–1234.
- Gunn, S.A., Gould, T.C. and Anderson, W.A.D. (1961) Competition for zinc in rat testis and dorsolateral prostate. *Acta Endocrinol.*, **37**, 24–30.
- Gustafson, A., Hedner, P., Schutz, A. *et al.* (1989) Occupational lead exposure and pituitary function. *Int. Arch. Occup. Environ. Health*, **61**, 277–281.
- Hajem, S., Moreau, T., Hannaert, P. *et al.* (1990) Influence of environmental lead on membrane ion transport in a French urban male population. *Environ. Res.*, **53**, 105–118.
- Heinemann, S.H., Teriau, H., Stuhmer, W. *et al.* (1992) Calcium channel characteristics conferred on the sodium channel by single mutations. *Nature*, **356**, 441–443.
- Herslag, A., Cooper, G.W. and Benoff, S. (1995) Pregnancy following discontinuation of a calcium channel blocker in the male partner. *Hum. Reprod.*, **10**, 599–606.
- Hirvonen, A. (1995) Genetic factors in individual responses to environmental exposures. *J. Occup. Environ. Med.*, **37**, 37–43.
- Hovatta, O., Venalainen, E.-R., Kuusimäki, L. *et al.* (1998) Aluminum, lead and cadmium concentrations on seminal plasma and spermatozoa, and semen quality in Finnish men. *Hum. Reprod.*, **13**, 115–119.
- Hu, W.Y., Wu, S.H., Wang, L.L. *et al.* (1992) A toxicological and epidemiological study on reproductive functions of male workers exposed to lead. *J. Hyg. Epidemiol. Microbiol. Immunol.*, **36**, 25–30.
- Hughes, E.G. and Brennan, B.G. (1996) Does cigarette smoking impair natural or assisted fecundity? *Fertil. Steril.*, **66**, 679–689.
- Hurley, I.R., Herslag, A. and Benoff, S. (1997) Mannose receptors, zinc and fertilization success. *Arch. STD HIV*, **11**, 241–253.

- Hutton, M. and Symon, C. (1986) The quantities of cadmium, lead, mercury and arsenic entering the U.K. environment from human activities. *Sci. Total Environ.*, **57**, 129–150.
- Jackson, H., Bock, M., Jackson, N.C. *et al.* (1995) A turnover study in the male rat of plasma-bound ^{59}Fe , ^{114}In and ^{109}Cd with particular reference to the gonad. *Nuclear Med. Commun.*, **16**, 112–120.
- Jacob, A., Hurley, I.R., Goodwin, L.O. *et al.* (1997a) Differential expression of potassium (K^+) channels during spermatogenesis in the rat. [Abstr. no. P-187] In *Scientific Program and Abstracts of the 53rd Annual Meeting of the American Society for Reproductive Medicine*, p. S182.
- Jacob, A., Hurley, I.R., Mandel, F.S. *et al.* (1997b) Human sperm potassium (K^+) channels activated by progesterone (P) binding: role in induction of the acrosome reaction and male infertility. [Abstr. no. P-036] In *Scientific Program and Abstracts of the 53rd Annual Meeting of the American Society for Reproductive Medicine*, pp. S109–S110.
- Jacob, A., Hurley, I.R., Cooper, G.W. *et al.* (1998a) Human sperm surface progesterone (P) receptor (R) regulates voltage gated potassium channel (VGK+C). [Abstr. no. 68] In *Program and Abstracts of the 23rd Annual Meeting of the American Society of Andrology*, p. 41.
- Jacob, A., Hurley, I.R., Cooper, G.W. *et al.* (1998b) Lead (Pb^{2+})-induced reproductive toxicity: action via a sperm head potassium channel. [Abstr. no. O-029] In *Scientific Program and Abstracts of the 54th Annual Meeting of the American Society for Reproductive Medicine*, pp. S11–S12.
- Jacob, A., Hurley, I., Mandel, F.S. *et al.* (1998c) Human sperm non-nuclear progesterone receptor expression is a novel marker for fertilization outcome. *Mol. Hum. Reprod.*, **4**, 533–542.
- Jeyendran, R.S., Van der Ven, H.H., Rosecrans, R. *et al.* (1989) Chemical constituents of human seminal plasma: relationship to fertility. *Andrologia*, **21**, 423–428.
- Jockenhovel, F., Bals-Pratsch, M., Bertram, H.P. *et al.* (1990) Seminal lead and copper in fertile and infertile men. *Andrologia*, **22**, 503–511.
- Johansson, L. (1989) Premature acrosome reaction in spermatozoa from lead-exposed mice. *Toxicology*, **54**, 151–162.
- Johnsen, S.G. (1970) Testicular biopsy score count – a method for registration of spermatogenesis in human testis: normal values and results in 335 hypogonadal males. *Hormones*, **1**, 2–25.
- Kantoff, P., Giovannucci, E. and Brown, M. (1998) The androgen receptor CAG repeat polymorphism and its relationship to prostate cancer. *Biochim. Biophys. Acta*, **1378**, C1–C5.
- Keck, C., Bramkamp, G., Behre, H.M. *et al.* (1995) Lack of correlation between cadmium in seminal plasma and fertility status of nonexposed individuals and two cadmium-exposed patients. *Reprod. Toxicol.*, **9**, 35–40.
- Kempinas, W.G., Favaretto, A.L.V., Melo, V.R. *et al.* (1994) Time-dependent effects of lead on rat reproductive functions. *J. Appl. Toxicol.*, **14**, 427–433.
- Kiss, T. and Osipenko, O.N. (1994) Toxic effects of heavy metals on ionic channels. *Pharmacol. Rev.*, **46**, 245–267.
- Klaiber, E.L., Broverman, D.M., Pokoly, T.B. *et al.* (1987) Interrelationships of cigarette smoking, testicular varicoceles, and seminal fluid indexes. *Fertil. Steril.*, **47**, 481–486.
- Kleiderlein, J.J., Nisson, P.E., Jessee, J. *et al.* (1998) CCG repeats in cDNAs from human brain. *Hum. Genet.*, **103**, 663–673.
- Kursh, E.D. (1987) What is the incidence of varicocele in a fertile population? *Fertil. Steril.*, **48**, 510–511.
- Lancranjan, I., Popescu, H.I., Gavanescu, O. *et al.* (1975) Reproductive ability of workmen occupationally exposed to lead. *Arch. Environ. Health*, **30**, 396–401.
- Lave, L.B., Hendrickson, C.T. and McMichael, F.C. (1995) Environmental implications of electric cars. *Science*, **268**, 993–995.
- Leinders, T., van Kleef, G.D.M. and Vijverberg, H.P.M. (1992) Distinct metal ion binding sites on Ca^{2+} -activated K^+ channels in inside-out patches of human erythrocytes. *Biochim. Biophys. Acta*, **1112**, 75–82.
- Lerda, D. (1992) Study of sperm characteristics in persons occupationally exposed to lead. *Am. J. Ind. Med.*, **22**, 567–571.
- Lin, S., Hwang, S.-A., Marshall, E.G. *et al.* (1996) Fertility rates among lead workers and professional bus drivers: a comparative study. *Ann. Epidemiol.*, **6**, 201–208.
- Lockitch, G. (1993) Perspectives on lead toxicity. *Clin. Biochem.*, **26**, 371–381.
- Lu Z.H. and Steinberg, M.H. (1996) Fetal hemoglobin in sickle cell anemia: relation to regulatory sequences *cis* to the beta-globin gene. *Blood*, **87**, 1604–1611.
- MacGregor, A.J. and Mason, H.J. (1990) Chronic occupational lead exposure and testicular endocrine function. *Hum. Exp. Toxicol.*, **9**, 371–376.
- MacKinnon, R. (1995) Pore loops: an emerging theme in ion channel structure. *Neuron*, **14**, 889–892.
- Mak, V. and Jarvi, K.A. (1996) The genetics of male infertility. *J. Urol.*, **156**, 1245–1267.
- Marmar, J.L. (1998) The emergence of specialized procedures for the acquisition, processing and cryopreservation of epididymal and testicular sperm in connection with intracytoplasmic sperm injection. *J. Androl.*, **19**, 517–526.
- Marmar, J.L., Katz, S., Praiss, D.E. *et al.* (1975) Semen zinc levels in infertile and postvasectomy patients and patients with prostatitis. *Fertil. Steril.*, **26**, 1057–1063.
- Mattison, D.R., Plowchalk, D.R., Meadows, M.J. *et al.* (1990) Reproductive toxicity: male and female reproductive systems as targets for chemical injury. *Med. Clin. N. Am.*, **74**, 391–411.
- Merino, G., Lira, S.C. and Martinez-Chequer, J.C. (1998) Effects of cigarette smoking on semen characteristics of a population in Mexico. *Arch. Androl.*, **41**, 11–15.
- Mieusset, R. and Bujan, L. (1995) Testicular heating and its possible contributions to male infertility: a review. *Int. J. Androl.*, **18**, 169–184.
- Miura, H., Tsujimura, A., Nishimura, K. *et al.* (1998) Susceptibility to idiopathic azoospermia in Japanese men is linked to HLA class I antigen. *J. Urol.*, **159**, 1939–1941.
- Moreau, T., Lellouch, J., Juguet, B. *et al.* (1983) Blood cadmium levels in a general male population with special reference to smoking. *Arch. Environ. Health*, **38**, 163–167.
- Mortensen, J.T. (1988) Risk for reduced sperm quality among metal workers, with special reference to welders. *Br. J. Indust. Med.*, **14**, 27–30.
- Muallem, S., Kwaitowska, K., Xu, X. *et al.* (1995) Actin filament disassembly is a final trigger for exocytosis in nonexcitable cells. *J. Cell Biol.*, **128**, 589–598.
- Naftulin, B.N., Samuels, S.J., Hellstrom, W.J.G. *et al.* (1991) Semen quality in varicocele patients is characterized by tapered sperm cells. *Fertil. Steril.*, **56**, 149–151.
- Needleman, H.L. (1988) The persistent threat of lead: medical and sociological issues. *Curr. Probl. Pediatr.*, **18**, 697–744.
- Ng, T.P., Goh, H.H., Ong, H.Y. *et al.* (1991) Male endocrine functions in workers with moderate lead exposure. *Br. J. Ind. Med.*, **48**, 485–491.
- Noack-Fuller, G., De Beer, C. and Seibert, H. (1993) Cadmium, lead, selenium, and zinc in semen of occupationally unexposed men. *Andrologia*, **25**, 7–12.
- Ogunlewe, J.O. and Osegbe, D.N. (1989) Zinc and cadmium concentrations in indigenous blacks with normal, hypertrophic, and malignant prostate. *Cancer*, **63**, 1388–1392.
- Oldereid, N.B., Thomassen, Y., Attramadal, A. *et al.* (1993) Concentrations of lead, cadmium and zinc in the tissues of reproductive organs of men. *J. Reprod. Fertil.*, **99**, 421–425.
- Oldereid, N.B., Thomassen, Y. and Purvis, K. (1994) Seminal plasma lead, cadmium and zinc in relation to tobacco consumption. *Int. J. Androl.*, **17**, 24–28.
- Olsen, J., Basso, O., Spinelli, A. *et al.* (1998) Correlates of care seeking for infertility treatment in Europe. *Eur. J. Public Health*, **8**, 15–20.
- Omu, A.E., Dashti, H., Mohamed, A.T. *et al.* (1995) Significance of trace elements in seminal plasma of infertile men. *Nutrition*, **11** (Suppl. 5), 502–505.
- Omu, A.E., Maksheed, M., Mohammed, A.T. *et al.* (1997) Characteristics of men and women with circulating antisperm antibodies in a combined infertility clinic in Kuwait. *Arch. Androl.*, **39**, 55–64.
- Ondrizek, R.R., Chan, P.J., Patton, W.C. *et al.* (1999) An alternative medicine study on the penetration of zona-free hamster oocytes and the integrity of sperm deoxyribonucleic acid. *Fertil. Steril.*, **71**, 517–522.
- Ophoff, R.A., Terwindt, G.M., Vergouwe, M.N. *et al.* (1996) Familial hemiplegic migraine and episodic ataxia type-2 are caused by mutation in the Ca^{2+} gene *CACNL1A4*. *Cell*, **87**, 543–552.
- Pacifici, R., Altieri, I., Gandini, L. *et al.* (1993) Nicotine, cotinine, and trans-3-hydroxycotinine levels in seminal plasma of smokers: effects on sperm parameters. *Ther. Drug Monit.*, **15**, 358–363.
- Pacifici, R., Altieri, I., Gandini, L. *et al.* (1995) Environmental tobacco smoke: nicotine and cotinine concentration in semen. *Environ. Res.*, **68**, 69–72.
- Panzer, S., Kuhl, D.P. and Caskey, C.T. (1995) Unstable triplet repeat sequences: a source of cancer mutations? *Stem Cells*, **13**, 146–157.
- Parazzini, F., Marchini, M., Tozzi, L. *et al.* (1993) Risk factors for unexplained dyspermia in infertile men: a case-control study. *Arch. Androl.*, **31**, 105–113.

- Paul, M. and Kurtz, S. (1994) Analysis of reproductive health hazard information on material safety data sheets for lead and the ethylene glycol ethers. *Am. J. Ind. Med.*, **25**, 403–415.
- Paulsen, C.A., Berman, N.G. and Wang, C. (1996) Data from men in greater Seattle area reveals no downward trend in semen quality: further evidence that deterioration of semen quality is not geographically uniform. *Fertil. Steril.*, **65**, 1015–1020.
- Peng, B.C.H., Tomashefsky, P. and Nagler, H.M. (1990) The cofactor effect: varicocele and infertility. *Fertil. Steril.*, **54**, 143–148.
- Perrault, S.D., Linder, R.E., Strader, L.F. *et al.* (1989) The value of multiple endpoint data in male reproductive toxicology: revelations in the rat. *Preg. Clin. Biol. Res.*, **302**, 179–192.
- Pleban, P.A. and Mei, D.-S. (1983) Trace elements in human seminal plasma and spermatozoa. *Clin. Chim. Acta*, **133**, 43–50.
- Plechaty, M.M., Noll, B. and Sunderman, F.W. Jr (1977) Lead concentrations in semen of healthy men without occupational exposure to lead. *Ann. Clin. Lab. Sci.*, **7**, 515–518.
- Radcliffe, J.M., Gladen, B.C., Wilcox, A.J. *et al.* (1992) Does early exposure to maternal smoking affect future fertility in adult males? *Reprod. Toxicol.*, **6**, 297–307.
- Ragan, H.A. and Mast, T.J. (1990) Cadmium inhalation and male reproductive toxicity. *Rev. Environ. Contam. Toxicol.*, **114**, 1–22.
- Rantala, M.L. and Koskimies, A.I. (1987) Semen quality of infertile couples – comparison between smokers and non-smokers. *Andrologia*, **19**, 42–46.
- Ratcliffe, J.M., Schrader, S.M., Steenland, K. *et al.* (1987) Semen quality in papaya workers with long term exposure to ethylene dibromide. *Br. J. Ind. Med.*, **44**, 317–326.
- Raunio, H., Husgafvel-Pursiainen, K., Anttila, S. *et al.* (1995) Diagnosis of polymorphisms in carcinogen activating and inactivating enzymes and cancer susceptibility – a review. *Gene*, **159**, 113–121.
- Robins, T.G., Bornman, M.S., Ehrlich, R.I. *et al.* (1997) Semen quality and fertility of men employed in a South African lead acid battery plant. *Am. J. Ind. Med.*, **32**, 369–376.
- Rogers, B., Myatt, G., Soderdahl, D. *et al.* (1985) Monitoring of suspected infertile men by the sperm penetration assay. *Fertil. Steril.*, **44**, 800–805.
- Rogers, B.J., Bastias, C., Coulson, R.L. *et al.* (1989) Cytochalasin D inhibits penetration of hamster eggs by guinea pig and human spermatozoa. *J. Androl.*, **10**, 275–282.
- Rubes, J., Lowe, X., Moore, D. *et al.* (1998) Smoking cigarettes is associated with increased sperm disomy in teenage men. *Fertil. Steril.*, **70**, 715–723.
- Russo, M.A., Kane, A.B. and Farber, J.L. (1982) Ultrastructural pathology of phalloidin-intoxicated hepatocytes in the presence and absence of extracellular calcium. *Am. J. Pathol.*, **109**, 133–144.
- Saaranen, M., Suistomaa, U., Kantola, M. *et al.* (1987a) Lead, magnesium, selenium and zinc in human seminal fluid: comparison with semen parameters and fertility. *Hum. Reprod.*, **2**, 475–479.
- Saaranen, M., Suonio, S., Kauhanen, O. *et al.* (1987b) Cigarette smoking and semen quality of men of reproductive age. *Andrologia*, **19**, 670–676.
- Saaranen, M., Kantola, M., Saarikoski, S. *et al.* (1989) Human seminal plasma cadmium: comparison with fertility and smoking habits. *Andrologia*, **21**, 140–145.
- Saldivar, L., Luna, M., Reyes, E., Soto, R. *et al.* (1991) Cadmium determination in Mexican-produced tobacco. *Environ. Res.*, **55**, 91–96.
- Sallmen, M., Lindbohm, M.-L., Anttila, A. *et al.* (1992) Paternal occupational lead exposure and congenital malformations. *J. Epidemiol. Commun. Health*, **46**, 519–522.
- Santoro, B., Liu, D.T., Yao, H., Bartsch, D. and Kandel E.R. (1998) Identification of a gene encoding for a hyperpolarization-activated pacemaker channel of brain. *Cell*, **93**, 717–729.
- Satin, J., Kyle, J.W., Chen, M. *et al.* (1992) A mutant TTX-resistant cardiac sodium channels with TTX-sensitive properties. *Science*, **256**, 1202–1205.
- Schlesinger, M.H., Wilets, I.F. and Nagler, H.M. (1994) Treatment outcome after varicocelelectomy. *Urol. Clin. N. Am.*, **21**, 517–529.
- Schrader, S.M. (1997) Clinical and experimental endpoints used to detect impaired sperm function. In *Course IX, Effects of toxicant exposure on sperm and oocyte quality, 30th Annual Postgraduate Program, The American Society for Reproductive Medicine*; pp. 45–71.
- Shafik, A. and Bedeir, G.A.M. (1980) Venous tension patterns in cord veins. I. In normal and varicocele individuals. *J. Urol.*, **123**, 383–385.
- Shieh, C. and Kirsch, G.E. (1994) Mutational analysis of ion conduction and drug binding sites in the inner mouth of voltage-gated K⁺ channels. *Biophys. J.*, **67**, 2316–2325.
- Smith, J.P., Smith, J.C. and McGall, A.J. (1960) Chronic poisoning from cadmium fumes. *J. Pathol. Bacteriol.*, **80**, 287–296.
- Smith M.T. and Suk, W.A. (1994) Application of molecular biomarkers in epidemiology. *Environ. Health Perspect.*, **102** (Suppl. 1), 229–235.
- Sofikitis, N., Miyagawa, I., Dimitriadis, D. *et al.* (1995) Effects of smoking on testicular function, semen quality and sperm fertilizing capacity. *J. Urol.*, **154**, 1030–1034.
- Sokol, R.Z. (1987) Hormonal effects of lead acetate in the male rat: mechanism of action. *Biol. Reprod.*, **37**, 1135–1138.
- Sokol, R.Z., Madding, C.E. and Swerdloff, R.S. (1985) Lead toxicity and the hypothalamic–pituitary–testicular axis. *Biol. Reprod.*, **33**, 722–728.
- Sokol, R.Z., Okuda, H., Nagler, H.M. *et al.* (1994) Lead exposure *in vivo* alters the fertility potential of mouse sperm *in vitro*. *Toxicol. Appl. Pharmacol.*, **124**, 310–316.
- Spinelli, A., Figa-Talamanca, I. and Osborn, J. (1997) Time to pregnancy and occupation in a group of Italian women. *Int. J. Epidemiol.*, **26**, 601–609.
- Spungin, B., Margalit, I. and Breitbart, H. (1995) Sperm exocytosis reconstructed in a cell-free system: evidence for the involvement of phospholipase C and actin filaments in membrane fusion. *J. Cell Sci.*, **108**, 2525–2535.
- Sram, R.J., Benes, I., Binkova, B. *et al.* (1996) Teplice program—the impact of air pollution on human health. *Environ. Health Perspect.*, **104** (Suppl. 4), 699–714.
- Stachel, B., Dougherty, R.C., Lahl, U. *et al.* (1989) Toxic environmental chemicals in human semen: analytical method and case studies. *Andrologia*, **21**, 282–291.
- Staessen, J., Amery, A., Bernard, A. *et al.* (1991) Effects of exposure to cadmium on calcium metabolism: a population study. *Br. J. Ind. Med.*, **48**, 710–714.
- Staessen, J.A., Buchet, J.P., Ginucchio, G. *et al.* (1996) Public health implications of environmental exposure to cadmium and lead: an overview of epidemiological studies in Belgium. *J. Cardiovasc. Risk*, **3**, 26–41.
- Steen, O.P. and Pangkahila, A. (1984a) Occupational influences on male fertility and sexuality. *Andrologia*, **16**, 5–22.
- Steen, O.P. and Pangkahila, A. (1984b). Occupational influences on male fertility and sexuality. *Andrologia*, **16**, 93–101.
- Stern, R.M., Berlin, A., Fletcher, A. *et al.* (1986) International conference on health hazards and biological effects of welding fumes and gases. *Int. Arch. Occup. Environ. Health*, **57**, 237–246.
- Stoltenberg, M., Sorensen, M.B., Danscher, G. *et al.* (1997a) Autometallographic demonstration of zinc ions in rat sperm cells. *Mol. Hum. Reprod.*, **3**, 763–767.
- Stoltenberg, M., Sorensen, M.B. and Danscher, G. (1997b) Histochemical demonstration of zinc ions in ejaculated semen. *Int. J. Androl.*, **20**, 229–236.
- Svensson, B.-G., Bjornham, A., Schutz, A. *et al.* (1987) Acidic deposition and human exposure to toxic metals. *Sci. Total Environ.*, **67**, 101–115.
- Suk, W.A., Collman, G. and Damstra, T. (1996) Human biomonitoring: research goals and needs. *Environ. Health Perspect.*, **104**, 479–483.
- Sweeney, T.E., Rozum, J.S. and Gore, R.W. (1995) Alteration of testicular microvascular pressures during venous pressure elevation. *Am. J. Physiol.*, **269**, H37–H45.
- Sylora J.A. and Pryor, J.L. (1994) Varicocele. *Curr. Therapy Endocrinol. Metab.*, **5**, 309–314.
- Tas, S., Lauwerys, R. and Lison, D. (1996) Occupational hazards for the male reproductive system. *Crit. Rev. Toxicol.*, **26**, 261–307.
- Telisman, S., Cvitkovic, P., Gavella, M. *et al.* (1990) Semen quality in men with respect to blood lead and cadmium levels. In *International Symposium on Lead and Cadmium Toxicology, Peking, Peoples' Republic of China, August 18–21, 1990*, pp. 29–32.
- Tesarik, J., Guido, M., Mendoza, C. *et al.* (1998) Human spermatogenesis *in vitro*: effects of follicle-stimulated hormone and testosterone on meiosis, spermiogenesis, and Sertoli cell apoptosis. *J. Clin. Endocrinol. Metab.*, **83**, 4467–4473.
- Tielemans, E., Burdorf, A., te Velde, E.R. *et al.* (1999) Occupationally related exposures and reduced semen quality: a case-control study. *Fertil. Steril.*, **71**, 690–696.
- Todd, A.C., Wetmur, J.G., Moline, J.M. *et al.* (1996) Unraveling the chronic toxicity of lead: an essential priority for environmental health. *Environ. Health Perspect.*, **104** (Suppl. 1), 141–146.
- Tsukidate, K., Yamamoto, K., Snyder, J.W. *et al.* (1993) Microtubule antagonists activate programmed cell death (apoptosis) in cultured rat hepatocytes. *Am. J. Pathol.*, **143**, 918–925.
- Umeyama, T., Ishikawa, H., Takeshima, H. *et al.* (1986) A comparative study of seminal plasma trace elements in fertile and infertile men. *Fertil. Steril.*, **46**, 494–499.

- Vigil, P., Wohler, C., Bustos-Obregon, E. *et al.* (1994) Assessment of sperm function in fertile and infertile men. *Andrologia*, **26**, 55–60.
- Vine, M.F. (1996) Smoking and male reproduction: a review. *Int. J. Androl.*, **19**, 323–337.
- Vine, M.F., Margolin, B.H., Morrison, H.I. *et al.* (1994) Cigarette smoking and sperm density: a meta-analysis. *Fertil. Steril.*, **61**, 35–43.
- Vine, M.E., Tse, C.-K.J., Hu, P.-C. *et al.* (1996) Cigarette smoking and semen quality. *Fertil. Steril.*, **65**, 835–842.
- Voeller, H.J., Wilding, G. and Gelmann, E.P. (1991) v-rasH expression confers hormone-independent *in vitro* growth to LNCaP prostate carcinoma cells. *Mol. Endocrinol.*, **5**, 209–216.
- Waalkes, M.P. and Rehms, S. (1994) Cadmium and prostate cancer. *J. Toxicol. Environ. Health*, **43**, 251–269.
- Wagner, U., Schlebusch, H., Van der Ven, H. *et al.* (1990) Accumulation of pollutants in the genital tract of sterility patients. *J. Clin. Chem. Clin. Biochem.*, **28**, 683–688.
- Wang, X., Wang, M., Niu, T., Chen, C. *et al.* (1998) Microsomal epoxide hydrolase polymorphism and risk of spontaneous abortion. *Epidemiology*, **9**, 540–544.
- Wang, Z., Chin, T.A. and Templeton, D.M. (1996) Calcium-independent effects of cadmium on actin assembly in mesangial and vascular smooth muscle cells. *Cell Motil. Cytoskel.*, **33**, 208–222.
- Watanabe, M. (1998) Polymorphic CYP genes and disease predisposition—what have the studies shown so far? *Toxicol. Lett.*, **28**, 167–171.
- Welch, L.S., Schrader, S.M., Turner, T.W. *et al.* (1988) Effect of exposure to ethylene glycol ethers on shipyard painters: II. male reproduction. *Am. J. Ind. Med.*, **14**, 509–526.
- Winder, C. (1993) Lead, reproduction and development. *NeuroToxicol.*, **14**, 303–318.
- Wong, M.K. and Koh, L.L. (1986) Mercury, lead, and other heavy metals in Chinese medicines. *Biol. Trace Elem. Res.*, **10**, 91–97.
- Wong, O., Utidjian, H.M.D. and Karten, V.S. (1979) Retrospective evaluation of reproductive performance of workers exposed to ethylene dibromide (EDB). *J. Occup. Med.*, **21**, 98–102.
- World Health Organization (1980) *Report of a Study Group: Recommended Health-Based Limits in Occupational Exposure to Heavy Metals*. Technical Report 647. WHO, Geneva, Switzerland.
- World Health Organization (1992) The influence of varicocele on parameters of fertility in a large group of men presenting to infertility clinics. *Fertil. Steril.*, **57**, 1289–1293.
- Wu, T.-N., Yang, G.-Y., Shen, C.-Y. *et al.* (1995) Lead contamination of candy: an example of crisis management in public health. *Lancet*, **346**, 1437.
- Wyrobek, A.J., Schrader, S.M., Perreault, S.D. *et al.* (1997) Assessment of reproductive disorders and birth defects in communities near hazardous chemical sites. III. Guidelines for field studies of male reproductive disorders. *Reprod. Toxicol.*, **11**, 243–259.
- Xu, B., Chia, S.-E., Tsakok, M. *et al.* (1993) Trace elements in blood and seminal plasma and their relationship to sperm quality. *Reprod. Toxicol.*, **7**, 613–618.
- Xu, C., Johnson, J.E., Singh, P.K. *et al.* (1996) *In vivo* studies of cadmium-induced apoptosis in testicular tissue of the rat and its modulation by a chelating agent. *Toxicology*, **107**, 1–8.
- Xuezhi, J., Youxin, L. and Yilan, W. (1992) Studies on lead exposure on reproductive system: a review of work in China. *Biomed. Environ. Sci.*, **5**, 266–275.
- Yamaguchi, M., Sakotu, J. and Takihara, H. (1989) The application of intrascrotal deep body temperature measurement for the noninvasive diagnosis of varicocele. *Fertil. Steril.*, **52**, 295–301.
- Yan, H., Carter, C.E., Xu, C. *et al.* (1997) Cadmium-induced apoptosis in the urogenital organs of the male rat and its suppression by chelation. *J. Toxicol. Environ. Health*, **52**, 149–168.
- Zamboni, L. (1987) The ultrastructural pathology of the spermatozoon as a cause of infertility: the role of electron microscopy in the evaluation of semen quality. *Fertil. Steril.*, **48**, 711–734.
- Zenses, M.T. (1995) Cigarette smoking as a cause of delay in conception. *Reprod. Med. Rev.*, **4**, 189–205.
- Zhuchenko, O., Bailey, J., Bonnen, P. *et al.* (1997) Autosomal dominant cerebella ataxia (SCA6) associated with small polyglutamine expansions in the alpha-1A voltage-dependent calcium channel. *Nature Genet.*, **15**, 62–69.

Received on May 6, 1999; accepted on January 17, 2000