

Forum Original Research Communication

Antioxidant Mechanisms of Nitric Oxide Against Iron-Catalyzed Oxidative Stress in Cells

VALERIAN E. KAGAN,^{1,2} ANDREY V. KOZLOV,³ YULIA Y. TYURINA,^{1,4}
ANNA A. SHVEDOVA,⁵ and JACK C. YALOWICH²

ABSTRACT

Three distinct antioxidant pathways are considered through which iron-catalyzed oxidative stress may be regulated by nitric oxide (NO). The first two pathways involve direct redox interactions of NO with iron catalytic sites and represent a fast response that may be considered an emergency mechanism to protect cells from the consequences of acute and intensive oxidative stress. These are (i) NO-induced nitrosylation at heme and non-heme iron catalytic sites that is capable of directly reducing oxoferryl-associated radicals, (ii) formation of nitrosyl complexes with intracellular "loosely" bound redox-active iron, and (iii) an indirect regulatory pathway that may function as an adaptive mechanism that becomes operational upon long-term exposure of cells to NO. In the latter pathway, NO down-regulates expression of iron-containing proteins to prevent their catalytic prooxidant reactions. *Antioxid. Redox Signal.* 3, 189–202.

INTRODUCTION

NITRIC OXIDE (NO) has emerged in recent years as a fundamental inter- and intracellular signaling molecule essential for the maintenance of homeostasis. This is due, to a large extent, to its ability to form complexes readily with iron-containing heme and non-heme proteins and its relatively "phlegmatic" behavior toward most biomolecules. It is becoming increasingly clear that redox reactions of NO play a very important role in its regulatory functions in cells and biological fluids. In particular, anti- and prooxidant effects of NO

are said to be associated with its numerous protective and toxic functions, respectively.

Antioxidant functions of NO may be realized through a number of direct redox interactions, as well as indirectly via redox-dependent effects on major metabolic regulatory pathways. As a free radical, NO is amazingly reactive toward organic radicals. This feature is responsible for the remarkable antioxidant effectiveness of NO in scavenging lipid peroxy and alkoxy radicals and inhibiting peroxidation of different lipids, such as membrane phospholipids and cholesterol (5, 31, 43, 60). Antioxidant effects of NO may be also achieved via its direct

Departments of ¹Environmental and Occupational Health and ²Pharmacology, University of Pittsburgh, Pittsburgh, PA, U.S.A.

³University of Veterinary Medicine, Institute of Pharmacology and Toxicology, and L. Boltzmann Institute for Experimental and Clinical Traumatology, Vienna, Austria.

⁴On leave from Institute of Evolutionary Physiology and Biochemistry, Russian Academy of Sciences, St. Petersburg, 194223 Russia.

⁵Health Effects Laboratory Division, Pathology and Physiology Research Branch, NIOSH, Morgantown, WV, U.S.A.

redox interactions with iron-centers during iron-catalyzed reactions leading to oxidative stress. These include chelation of redox-active metal ions by NO[•] (31, 40), as well as reduction of oxoferryl species (22, 23). Finally, NO may affect oxidative stress indirectly by down-regulating expression of iron-containing proteins through posttranscriptional mechanisms or up-regulation of antioxidant enzymes, *i.e.*, contributing to antioxidant effects through its action as a signaling molecule. These direct and indirect interactions of NO with three types of iron species that are mainly responsible for the catalysis of oxidative stress reactions in cells and biological fluids—heme proteins, non-heme iron proteins, and “loosely bound” iron readily available for iron chelators—are the focus of our discussion here.





PROOXIDANT ACTIVITY OF HEME PROTEINS

In blood, most iron is bound to hemoglobin (Hb), whereas myoglobin (Mb) represents one of the major reservoirs of iron in some cell types, such as muscle cells (*e.g.*, cardiomyocytes). Mb and Hb react with alkylhydroperoxides (including lipid hydroperoxides) to yield several redox-active species, as well as a number of free radical intermediates, such as alkylhydroperoxyl (ROO[•]), alkyloxyl (RO[•]), and alkyl (R[•]) radicals (10, 32, 67). The oxidation of ferric (met) forms of Mb (Mb-Fe³⁺) and Hb (Hb-4Fe³⁺) by hydroperoxides is presumed to proceed via two-electron oxidation to give a long-lived oxoferryl (Fe⁴⁺=O) species, one oxidation equivalent above the ferric state, plus a transient protein radical (10, 21). There are two (or more) sites for the protein-derived radical: tyrosine (Tyr¹⁰³) or tryptophan (Trp¹⁴). The Mb Trp¹⁴ peroxy radical has been shown to react rapidly with a wide range of proteins to give long-lived secondary radicals on the target protein. Recipient peptides/proteins that contained a tyrosine and/or tryptophan amino acid residue were most susceptible to free radical transfer (11). These protein-to-protein radical transfer reactions may be responsible for protein oxidation (51). Additionally, antioxidants (GSH, ascorbate, Trolox C, vitamin E,

and urate) can directly reduce Mb-derived peroxy radical to produce antioxidant-derived radicals (35). This results in antioxidant depletion. Protein radicals have been shown to react with peribacteroid membrane fractions, with the consequent generation of lipid-derived radicals (35). Apart from short-lived radicals, longer-lived protein radicals associated with oxoferryl hemoproteins may be involved in the initiation of lipid peroxidation (8).

RELEASE OF IRON FROM HEME IN THE MET-Hb/*tert*-BUTYL HYDROPEROXIDE (*tert*-BuOOH) SYSTEM

A plethora of data indicate that two highly reactive iron species are formed during interaction of heme with peroxides, namely oxoferryl (ferryl, perferryl) species and “loosely” bound iron that can be released from the heme. In the absence of oxygen, formation of these species can be described by the following reactions:



In the presence of oxygen, the reaction of oxygen with protein radical yields the peroxy radical ([•]O-O-Hb-heme-Fe⁴⁺=O). In accordance with this scheme, “loosening” of metal is usually associated with oxidation of protein, as has been shown for different classes of superoxide dismutases (39, 73). Self-peroxidation of met-Hb (25) leading to the formation of protein radicals and subsequent protein fragmentation is likely to be the cause of “loosening” and release of iron from the heme. This scheme does not consider the redox state of released iron. It has been shown that free iron produced by heme/perox-

ide interactions is in the ferrous state (9, 33, 52). The mechanisms involved in the formation and release of ferrous (Fe^{2+}) iron from ferryl (Fe^{4+}) species are not fully understood. This, however, is important because ferrous iron is a strong promoter of oxidative stress.

Interestingly, the release of ferrous iron is more characteristic of the interaction of hemoproteins with organic hydroperoxides (cumene hydroperoxide, *tert*-BuOOH) than with hydrogen peroxide (52). This is consistent with the results of Tajima (67), which showed the existence of heme-peroxide complexes. Later, Jinno et al. (38) demonstrated that hemoprotein-*n*-butylperoxide complex is rapidly decomposed to non-heme iron. Thus, it seems that investigation of the early stages of interactions of hemoproteins with organic hydroperoxides may give the key to understanding the mechanism(s) of iron release.

It is presumed that the initial product of this reaction is the ferryl heme cation radical (25), although the species has never been directly detected by electron paramagnetic resonance (EPR) spectroscopy. Characterization of this very reactive oxygen intermediate remains a challenging area of research. Recent calculations performed for the heme of cytochromes P-450 showed that such a putative reactive oxygen intermediate should contain two parallel unpaired spins on the $\text{Fe}=\text{O}$ moiety and an antiparallel unpaired spin distributed between the $a_{2u}(\pi)$ orbital of the porphyrin ring and the π -orbital of ligand atom (46), indicating the existence of an organic radical in the heme iron moiety. Based on kinetic behavior of the peroxyl radical and the protein radical in a met-Hb/hydrogen peroxide model system, Svislitskiy et al. (66) suggested that peroxyl radicals with characteristic features at $g = 2.035$ are not formed from protein radicals ($g = 2.0042$). They postulated the existence of a radical intermediate not detectable by EPR, but capable of reacting with oxygen to yield the $g = 2.035$ protein peroxyl radical EPR signal. Notably, the peroxyl radical EPR signal was not saturable by power likely due to its close proximity to the heme iron moiety. In a recent study by Baron et al. (4), a similar EPR signal at $g \approx 2.035$ characteristic of a peroxyl radical was observed (but not assigned) in the reaction sys-

tem met-Mb/hydrogen peroxide/linoleate immediately after mixing of the reagents. Despite the assignment of the $g = 2.035$ signal to a protein peroxyl radical, an absolutely identical signal was observed during oxidation of *tert*-BuOOH in the absence of protein, clearly indicating that it is derived from the *tert*-BuOO \cdot radical (34).

Thus, it appears that there are two contradicting results on the nature of products formed during interactions of hemoproteins with organic hydroperoxides: (i) EPR-detectable peroxyl radicals do not seem to derive from the protein radical, and (ii) ferrous iron is released from ferryl (Fe^{4+}) heme species, suggesting that the reduction step takes place. At first glance, these two findings seem to be disconnected. Three sets of results, however, permit logical combination of these findings into one scheme: (i) *tert*-BuOOH-heme complex is involved in iron release; (ii) the $g = 2.035$ peroxyl radical and protein radical seem to have a different origin; and (iii) the $g = 2.035$ peroxyl radical EPR signal is detectable very early in the reaction between heme and peroxide, much earlier than the protein radical. Altogether, these facts suggest that a reductive mechanism of iron release during the interaction of heme proteins with peroxides is operational.

To support this hypothesis experimentally, we performed direct EPR studies of met-Hb/*tert*-BuOOH in expectation that this relatively slow reaction will permit detection of the key free radical intermediates, as well as release ("loosening") of ferrous iron. We found no signals in the high-field ($g = 1.95\text{--}2.10$) segment of the EPR spectrum at 0 time (Fig. 1, curve 1). After 30 s, an integral peak at $g = 2.035$ and a trough at $g = 2.004$ (species A) was detected in the EPR spectrum (Fig. 1, curve 2). The peak at $g = 2.035$ gradually decayed over time, and a new species (species B) appeared in the EPR spectrum recorded at 5 min (Fig. 1, curve 3). This species had integral peaks at $g = 2.025$ and $g = 2.017$, and a shoulder at $g = 2.010$. The signal intensity of species B was maximal at 15 min (Fig. 1, curves 4 and 5).

Species A is likely a peroxyl radical signal, whose appearance has been well documented for such systems (8, 47, 50, 55, 65, 66) and for a model system containing high concentrations





FIG. 1. High-resolution low-temperature EPR spectra of the reaction mixture met-Hb/*tert*-BuOOH acquired at different time intervals after the reagents were mixed. Conditions were as follows: met-Hb (300 μ M heme iron), *tert*-BuOOH (1.5 mM), disodium phosphate buffer (50 mM, pH 7.4 at 25°C). The reaction was started by the addition of *tert*-BuOOH. EPR spectra were recorded at 150°C; gain, 500 and 200; modulation amplitude, 0.5 mT; power, 10 mW; time constant, 0.03 s; scan time, 4 min; scan range, 15 mT. Spectra 1–5 were recorded at 0, 0.5, 5, 15, and 30 min after beginning of the reaction, respectively.

of *tert*-BuOOH and iron ions (34). If this peroxy radical is the first intermediate of the interaction between *tert*-BuOOH and heme iron, then it must be formed via reaction:



We were able to obtain similar peroxy radical signals by one-electron oxidation of *tert*-BuOOH by $\text{K}_3\text{Fe}(\text{CN})_6$, a strong one-electron oxidant (Fig. 2, inset C). To clarify whether signals observed in the met-Hb/*tert*-BuOOH system are associated with heme iron, we used an EPR power saturation technique to compare relaxation parameters of species A, species B, and peroxy radical generated in the presence of $\text{K}_3\text{Fe}(\text{CN})_6$. Species A (detected after 30 s) was not saturable by increasing the power up to 180 mW (Fig. 2, filled squares), whereas species B (Fig. 2, filled diamonds) as well as the EPR signal of peroxy radical observed in the *tert*-BuOOH/ $\text{K}_3\text{Fe}(\text{CN})_6$ system (Fig. 2, filled circles) were saturated at ~ 40 – 100 mW power.

This indicates that species A is likely associated with the heme iron moiety of met-Hb in contrast to species B, which is likely a protein-centered radical.

The power saturation technique also permits differentiation between overlapping signals, because different relaxation constants cause a change in the signal features. To clarify whether species A and species B are single signals or a mixture of different signals, we compared the shape of signals obtained at various power values. Each spectrum was divided by a constant corresponding to its intensity, resulting in a series of spectra with constant integral intensity. The series of spectra obtained for species A and species B are presented in Fig. 3. Power saturation did not affect the shape of the signal for species A. This indicates that species A represents a single signal (formed 30 s after beginning of the reaction). In contrast, species B ap-




FIG. 2. Power saturation of met-Hb/*tert*-BuOOH EPR signals. Filled squares and spectrum A: EPR signal detected after 30 s of incubation of met-Hb/*tert*-BuOOH at 25°C (species A); filled circles and spectrum C: EPR signals detected in the presence of $\text{K}_3\text{Fe}(\text{CN})_6$; filled diamonds and spectrum B: EPR signal detected after 30 min of incubation of met-Hb/*tert*-BuOOH at 25°C (species B). EPR spectra were recorded at -150°C ; gain, 500 and 200; modulation amplitude, 0.5 mT; power, 10 mW; time constant, 0.03 s; scan time, 4 min; scan range, 15 mT.




FIG. 3. Serial spectra obtained for species A (A) and B (B) at different values of power. Spectra 1–5 were recorded at 1, 10, 40, 100, and 180 mW, respectively. The spectrum of species B includes features of protein radical, $O=Fe^{4+}$ —protein \cdot (filled circle), and features of protein-centered peroxy radical, $O=Fe^{4+}$ —protein- $O\cdot$ (filled triangle).

It appears to be a mixture of at least two signals (see Fig. 3) that can be assigned to protein radical and protein-centered peroxy radical, as has been well documented (8, 47, 50, 55, 65, 66).

We suggest that species A is the primary radical intermediate formed as a result of iron “loosening” (release) independently of the pathway that yielded species B. This is supported by the following results. We observed neither *tert*-BuO \cdot radicals nor protein radicals that are expected to emerge from one-electron reduction of *tert*-BuOOH by met-Hb at the initial stage of the reaction. Instead, only species A could be seen 30 s after met-Hb and *tert*-BuOOH were mixed. If species A is the primary intermediate of the reaction between heme iron and *tert*-BuOOH, it could be formed only

through one-electron oxidation of *tert*-BOOH and reduction of heme iron, respectively. In line with this, generation of species A (but not of species B) was remarkably inhibited when oxy-Hb (Fe^{2+}) was used instead of met-Hb (data not shown). If *tert*-BuOO \cdot is the first reaction intermediate of Fe^{3+} porphyrin with *tert*-BuOOH as has been proposed by Tajima (67) for hydrogen peroxide, then *tert*-BuOO \cdot is expected to be in close proximity to the reduced iron center. In fact, our EPR power saturation experiments demonstrated that species A was indeed associated with heme.

The formation of species A may be related to heme iron reduction and the appearance of Hb(Fe^{2+}). This, however, was not observed in the system (data not shown). Another possibility is “loosening” (release) of ferrous ions from “reductive” interaction between *tert*-BuOOH and Hb. To prove this, we used the Fe^{2+} chelator, *o*-phenanthroline, to detect the ferrous iron. We found that the *o*-phenanthroline-chelatable ferrous iron accumulated over time in the incubation system, and its amount inversely correlated with the concentration of species A (Fig. 4). This implies that species A is likely associated with the “loosening” (release) of ferrous iron.

In summary, our results imply the existence





FIG. 4. Time course of met-Hb, species A, and *o*-phenanthroline- Fe^{2+} complex initiated by met-Hb/*tert*-BuOOH. Filled squares: time course of species A; filled diamonds: time course of met-Hb; open squares: time course of *o*-phenanthroline- Fe^{2+} complex.

of two reaction pathways through which *tert*-BuOOH interacts with met-Hb (Scheme 1). The first one (which is faster) produces *tert*-BuOO[•] by one-electron oxidation of *tert*-BuOOH (Scheme 1, steps I–III), whereas the second one (which is slower) yields ferryl species by two-electron oxidation of heme iron (Scheme 1, steps I, II, IV, and V). This is in agreement with earlier findings that ferryl-Hb is the only species formed at low *tert*-BuOOH/Hb ratios; with the increase of *tert*-BuOOH/Hb ratio, the



Scheme 1. Proposed reaction stages during interaction of *tert*-BuOOH with met-Hb. I: The system includes high-spin heme Fe^{3+} ($S = 5/2$) and free *tert*-BuOOH ($S = 0$). II: *tert*-BuOOH and met-Hb form a complex consisting of low-spin heme Fe^{3+} ($S = 1/2$) and *tert*-BuOOH ($S = 0$). III: *tert*-BuOOH reduces heme iron yielding species A, a peroxy radical ($S = 1/2$), and loosely bound Fe^{2+} ($S = 0$). IV: Two-electron oxidation of heme iron by *tert*-BuOOH yielding per ferryl-Hb. The system includes low-spin Fe^{5+} ($S = 1/2$). V: One-electron oxidation of protein by Fe^{5+} yielding Hb- Fe^{4+} ($S = 0$) and species B, representing two radicals, a protein radical ($S = 1/2$) and a protein peroxy radical ($S = 1/2$).




FIG. 5. Low-temperature EPR spectra of K/VP.5 cells treated with *tert*-Bu-OOH. K/VP.5 cells (250×10^6 cells/ml) were incubated in 50 mM disodium phosphate buffer, pH 7.4, in the presence of *tert*-BuOOH ($80 \mu\text{mol}/10^6$ cells) at 25°C . At the indicated time point, cell suspensions were frozen at -150°C and used for EPR measurements. 1, control (no *tert*-Bu-OOH added); 2, cells treated with *tert*-BuOOH (30 s after addition of *tert*-BuOOH); 3, cells treated with *tert*-BuOOH (15 min after addition of *tert*-BuOOH). EPR spectra were recorded at -150°C ; gain, 100 (3) and 500 (1 and 2); modulation amplitude, 0.5 mT; power, 10 mW; time constant, 0.03 s; scan time, 4 min; scan range, 15 mT.

release of free iron becomes detectable (8, 48). Simple thermodynamic considerations also indicate that *tert*-BuOOH is more likely to act as an electron donor at high *tert*-BuOOH/Hb ratios, whereas Hb is more likely to be a donor of electrons at high Hb/*tert*-BuOOH ratios. Clearly, achievable concentrations of hydroperoxides (H_2O_2 , lipid hydroperoxides) along with the concentrations of heme proteins in cells or biological fluids determine which of these pathways predominate under physiologically relevant conditions.

In separate experiments, we performed EPR studies of radical species formed by *tert*-BuOOH in K/VP.5 cells, a subclone of human erythropoietic leukemia K562 cells that contain a relatively high concentration of endogenous Hb [~ 30 – $40 \text{ pmol}/10^6$ cells (59)]. We found that after the addition of *tert*-BuOOH to K/VP.5 cells at concentrations close to those used in our model experiments with met-Hb and oxy-Hb, two major species appeared in the EPR spectra (Fig. 5). One signal ($g = 2.035$) was detectable

almost immediately after addition of *tert*-BuOOH (30 s), and the other one ($g = 2.004$) was present in the spectra at both 30 s and 15 min of incubation. Based on the values of g factors and the shape of the signals, they may be tentatively assigned to the *tert*-BuOO \cdot signal and ferryl-protein radical signal, respectively. Thus, essentially very similar mechanisms of Hb/*tert*-BuOOH interactions may operate in cells. Indeed, we were able to detect both *tert*-BuOO \cdot signal and ferryl-protein radical signal as the primary and secondary free radical species formed in the course of the reaction between endogenous (oxy)Hb and *tert*-BuOOH. Although these results strongly suggest that more than one reactive oxidant may be involved in oxidative stress induced by Hb *in vivo*, further studies are necessary to quantitate contributions of the two factors in oxidations of physiologically relevant reductants.

ANTIOXIDANT PATHWAYS OF NO AGAINST IRON-CATALYZED OXIDATIVE STRESS

There are three distinct pathways through which iron-catalyzed oxidative stress may be regulated by NO as presented in our discussion and experimental results below.

NO-induced nitrosylation at heme and non-heme iron catalytic sites protects against oxidative stress

In the first pathway, a direct chemical interaction of NO with hydroperoxides at non-heme and heme iron catalytic sites prevents generation of oxoferryl species and different free radicals, and thus precludes free radical-induced damage of critical biomolecules. The possibility of this pathway has been initially established in simple chemical systems *in vitro* (22, 40).

In the initial experiments, the reactions of met-Mb and met-Hb, oxidized to their respective oxoferryl free radical species (Mb-Fe $^{4+}=O$ and Hb-4Fe $^{4+}=O$) by *tert*-BuOOH, with NO were studied in a nitrogen atmosphere (22). It was shown that NO may reduce both oxoferryl and apoprotein free radical electrophilic centers of Mb-Fe $^{4+}=O$ /Hb-4Fe $^{4+}=O$ and eliminate *tert*-butyl(per)oxyl radicals, thus protect-

ing against oxidative damage. Moreover, NO reduced Mb-Fe $^{4+}=O$ /Hb-4Fe $^{4+}=O$ to their respective ferric (met) forms and prevented oxidation of lipids [*cis*-parinaric acid (PnA) integrated into liposomes], oxidation of luminol, and formation of the *tert*-butyl(per)oxyl adduct with the spin trap 5,5'-dimethylpyrroline *N*-oxide. NO eliminated the signals of tyrosyl radical detected by EPR and oxoferryl detected by mass spectrometry in the reaction of *tert*-BuOOH with met-Mb (22). Subsequent studies showed that a combination of Hb/*tert*-BuOOH added extracellularly to vascular smooth muscle cells (VSMCs) caused oxidative stress as evidenced by peroxidation of their major phospholipids—phosphatidylethanolamine (PE), phosphatidylcholine (PC), phosphatidylserine (PS), and phosphatidylinositol (PI). The oxidative stress was accompanied by cytotoxic effects of oxy-Hb/*tert*-BuOOH and met-Hb/*tert*-BuOOH on VSMCs (62). In the presence of an NO donor, (Z)-1-[N-(3-ammoniopropyl)-N-(*n*-propyl)amino]diazene-1-ium-1,2-diolate (NOC-15), VSMCs were protected against oxidative stress and cytotoxicity induced by Hb/*tert*-BuOOH. The protective effect of NOC-15 was most likely due to its ability to form NO-heme Hb, hence to prevent the formation of oxoferryl-Hb species (62).

Further experiments demonstrated that this antioxidant function of NO, *i.e.*, its ability to reduce oxoferryl Mb/Hb species, is realized in cells (23, 24). In particular, it was demonstrated that in K/VP.5 cells (a subline of human erythroleukemia K562 cells with elevated intracellular Hb concentrations), *tert*-BuOOH-induced formation of oxoferryl-Hb-derived free radical species was proportional to endogenous Hb concentrations (23). Further, a correlation between the production of oxoferryl-Hb-derived radicals, peroxidation of major membrane phospholipids (PC, PE, PS, PI, and cardiolipin), and cell viability was demonstrated. Most importantly, when the cells were pretreated with an NO donor, NOC-15, that caused nitrosylation of non-heme iron centers and Hb-heme, *tert*-BuOOH was not able to induce either formation of oxoferryl-Hb-derived free radical species or peroxidation of phospholipids in the cells. No cytotoxicity was observed in NOC-15-pretreated cells after incubation with *tert*-BuOOH. These results indicate that NO produces iron-ni-

trosyl complexes whose redox interactions with *tert*-BuOOH prevented generation of oxoferryl-Hb-derived free radical species.

Qualitatively very similar results were obtained with cardiac myocytes (24). Two distinct free radical species—alkoxyl radicals associated with non-heme iron catalytic sites and Mb protein-centered peroxy radicals, found in low-temperature EPR spectra of cardiac myocytes exposed to *tert*-BuOOH—were completely abolished when the cells were incubated with the combination of NOC-15 and *tert*-BuOOH. NO acted as an effective antioxidant in live cardiomyocytes as it was able to protect cardiomyocytes completely against both *tert*-BuOOH-induced phospholipid peroxidation and cytotoxicity.

Overall, these results on antioxidant effects of NO against oxidative stress induced by combinations of organic hydroperoxides with hemoproteins (Mb or Hb) are summarized in Tables 1 and 2. Notably, formation of NO-heme hemoproteins was essential for antioxidant protection via the mechanism of direct reduction of oxoferryl-derived radicals. The protective effects afforded by nitrosothiols that are known to S-nitrosylate hemoproteins were minimal (Table 2).

Intracellular "loosely" bound redox-active iron is available for NO to form nitrosyl complexes

The second pathway is based on the ability of NO to interact with "loosely" bound iron constitutively present in cells or formed as a result of ongoing oxidative stress (see above). Formation of nitrosyl iron complexes inhibits their catalytic prooxidant reactions. This mechanism is similar to the effects of iron chelators, e.g., deferoxamine (DFO).

In plasma, iron is normally bound to proteins mainly due to the concerted action of ceruloplasmin and apotransferrin. The former oxidizes ferrous ions to the ferric form, whereas the latter sequesters ferric ions to form transferrin (19). Transferrin receptors transport iron into cells, and the iron is subsequently stored within ferritin (69). Intracellular iron is mostly tightly bound by iron-storing proteins (e.g., ferritin) and only a small part of total iron, so-called "loosely" bound iron ["free iron" or tran-

sferable iron pool (36)], may be present as low-molecular-weight iron complexes with a low stability constant. This "loosely" bound iron (which may also include iron "loosely" bound to proteins) is available for chelators, such as DFO and phenanthroline. Normally, iron associated with ferritin does not possess any catalytic activity (1, 37, 57). In contrast, the pool of "loosely" bound iron is capable of catalyzing oxidative stress reactions (2, 27). Therefore, regulation of intracellular iron is deemed to be one of the important mechanisms that determines sensitivity to oxidative stress (6, 29).

One of the effective ways to protect against oxidative stress induced by "loosely" bound iron is to use iron chelators, such as *o*-phenanthroline, EDTA, diethylenetriaminepentaacetic acid, or DFO, that form stable redox-inactive complexes with iron (17, 63). DFO is the most commonly used and pharmacologically efficacious chelator (12, 15, 17, 58, 63). In addition, it is the only iron chelator currently approved for clinical use for treatment of iron overload, including acute iron poisoning and treatment of chronic iron overload in transfusion-dependent anemias, such as β -thalassemia. Based on extensive clinical use of DFO, it was concluded that DFO-available iron can promote oxidative stress (7, 18, 58). Technical difficulties (it is administered by subcutaneous infusion over 8–12 h), as well as problems with compliance, have prompted a search for alternatives to DFO, in particular, iron-chelating molecules active after oral administration (12).

Lately, inhaled NO is broadly used in clinical practice. EPR studies demonstrated that NO and DFO bind the same pools of intracellular iron (45). DFO reacts with intracellular iron to form paramagnetic iron complexes with characteristic EPR signal with $g = 4.3$ (Fig. 6) (44). Treatment of tissue homogenates with sodium nitrite (or gaseous NO) results in appearance of the signal from dinitrosyl iron complexes with $g = 2.03$ (Fig. 6) (70, 71). Administration of DFO prior to sodium nitrite resulted in the $g = 4.3$ EPR signal. Conversely, when DFO was administered after sodium nitrite, only the EPR signal with $g = 2.03$ was detectable (45).

These data demonstrate that the same endogenous iron was available for chelation with DFO or with NO to produce EPR-detectable

TABLE 1. EFFECT OF NOC-15 ON OXIDATION OF PnA-LABELLED PHOSPHOLIPIDS INDUCED BY *tert*-BuOOH IN DIFFERENT CELL LINES

Cell line	<i>tert</i> - BuOOH (μ M)	NOC-15 (μ M)	Time (min)	Oxidized PnA (ng/ μ g of total lipid phosphorus)				
				PI	PE	PS	PC	
K/VP.5 erythroleukemia cells (n = 3)	100	0	60	57.1 \pm 7.1	48.2 \pm 8.1	58.9 \pm 12.8	42.8 \pm 2.7	
	100	40	60	23.2 \pm 2.2*	25.4 \pm 7.2*	28.6 \pm 5.7*	17.8 \pm 1.5*	
	100	80	60	27.8 \pm 2.8	12.5 \pm 1.0 [†]	7.1 \pm 0.8 [†]	14.3 \pm 0.6 [†]	
K/VP.5 erythroleukemia cells treated with hemin (25 μ M) (n = 3)	100	0	60	62.3 \pm 9.3	46.4 \pm 6.2	66.1 \pm 11.6	44.6 \pm 3.4	
	100	40	60	7.6 \pm 0.3 [†]	16.1 \pm 1.4*	32.1 \pm 10.1*	11.1 \pm 0.7 [†]	
	100	80	60	7.1 \pm 0.3 [†]	15.0 \pm 1.6 [†]	7.1 \pm 1.4 [†]	11.1 \pm 0.8 [†]	
VSMCs treated with oxy-Hb (5 μ M) (n = 5)	150	0	60	56.9 \pm 2.9	39.7 \pm 4.1	56.8 \pm 3.1	30.5 \pm 3.6	
	150	50	60	40.0 \pm 3.1*	24.0 \pm 1.7*	38.4 \pm 1.3*	22.2 \pm 3.1	
	150	200	60	30.5 \pm 3.0*	15.6 \pm 1.4*	9.1 \pm 6.3 [†]	7.5 \pm 4.7*	
Cardiomyocytes (n = 3)	150	0	15		9.2 \pm 0.6	41.0 \pm 5.4	3.4 \pm 0.7	
	150	150	15		2.4 \pm 0.7	0.1 \pm 0.1 [†]	1.5 \pm 0.6	

Cells loaded with PnA (5 μ g/10⁶ cells) were exposed to *tert*-BuOOH in the presence or absence of NOC-15. NOC-15 was added 15 min prior to induction of oxidative stress. At the end of incubation, cells were harvested and washed, and lipids were extracted and resolved by HPLC. All values are means \pm SEM. The amount of intracellular Hb in K/VP.5 cells nontreated and treated with hemin was estimated to be 25 pmol of Hb/10⁶ cells and 90 pmol of Hb/10⁶ cells, respectively. The amount of intracellular oxy-Mb in cardiomyocytes was estimated to be 50 pmol/10⁶ cells.

* $p < 0.05$, [†] $p < 0.005$ versus *tert*-BuOOH.

TABLE 2. EFFECT OF DIFFERENT NO DONORS ON OXIDATION OF PnA-LABELED PHOSPHOLIPIDS INDUCED BY *tert*-BuOOH IN VSMCs

NO donors	<i>tert</i> -BuOOH (μ M)	donor (μ M)	Time (min)	Oxidized PnA (ng/ μ g of total lipid phosphorus)			
				PI	PE	PS	PC
None (n = 5)	150	0	60	7.6 \pm 0.4	26.3 \pm 2.8	6.6 \pm 0.4	72.3 \pm 8.5
GS-NO (n = 3)	150	200	60	8.0 \pm 0.8	25.6 \pm 1.5	7.1 \pm 1.1	74.4 \pm 8.3
NAC-S-NO (n = 3)	150	200	60	7.0 \pm 0.7	29.3 \pm 1.9	7.6 \pm 0.6	78.5 \pm 6.4
NOC-15 (n = 5)	150	200	60	4.0 \pm 0.4*	10.9 \pm 1.0*	2.2 \pm 1.5†	37.2 \pm 2.3*

Cells loaded with PnA (5 μ g/ 10^6 cells) were exposed to *tert*-BuOOH in the presence or absence of GS-NO, NAC-S-NO, or NOC-15. NO donors were added 15 min prior to induction of oxidative stress. At the end of incubation, cells were harvested and washed, and lipids were extracted and resolved by HPLC. All values are means \pm SEM.

* $p < 0.05$, † $p < 0.005$ versus control.

paramagnetic centers. These results also suggest that NO may protect against iron toxicity through binding catalytically redox-active iron that is also available for chelation by DFO. In other words, NO may act as an antioxidant capable of inhibiting oxidative stress induced by loosely bound (DFO-chelatable) iron.

NO down-regulates expression of iron-containing proteins to protect against oxidative stress

The third antioxidant pathway for NO is associated with its ability to regulate the intracellular content of iron-containing catalytic sites. Expression of several essential proteins involved in cellular iron homeostasis has been shown to be regulated by iron-regulatory proteins 1 and 2 (IRP-1, IRP-2) binding to specific mRNA sequences—iron responsive elements (IREs) (3, 26, 28, 41, 49, 54, 61, 68). Functional IREs have been identified in the 5' untranslated

regions of ferritin H-chain, ferritin L-chain, erythroid aminolevulinic synthase, mitochondrial aconitase, and succinate dehydrogenase, as well as the 3' untranslated regions of transferrin receptors (13, 54).

NO can directly interact with IRP-1, thus triggering IRE-dependent responses (13, 16). Two different sites on IRP-1 can be targeted by NO. One target site may be the sulfhydryl group of Cys⁴³⁷, which has been shown to regulate the binding of IRP-1 to IREs (16, 30, 64). The second site is the dynamic Fe-S cluster whose removal from the apoprotein is induced by NO binding (53). Hence, NO binding to IRP-1 initiates IRP-1/IRE interactions because only the IRP-1 apoprotein can bind to the IREs (16, 30, 64). Therefore, chronic exposure to NO donors or overproduction of endogenous NO in inducible NO synthase (iNOS)-transduced cells results in decreased cellular content of non-heme and heme-iron by coordinate translational regulatory mechanisms (14, 20, 42, 54, 56). Obviously, cells with decreased levels of endogenous redox-active iron catalytic sites would be less susceptible to oxidative stress.

This has been directly demonstrated in experiments with K/VP.5 cells transduced with a retroviral vector containing the human iNOS gene (K/VP.5-iNOS) (72). K/VP.5-iNOS cells were remarkably less sensitive to the cytotoxic effects of *tert*-BuOOH compared with K/VP.5 cells. It was determined that levels of non-heme and heme (Hb) iron were dramatically decreased in K/VP.5-iNOS cells compared with K/VP.5 cells in line with the sharply decreased intensities of EPR signals of nitrosylated species observed in the presence of endoge-




FIG. 6. EPR spectra of dinitrosyl iron complexes ($g = 2.03$) and DFO-iron complexes ($g = 4.3$).

nously produced NO (by iNOS) or upon addition of exogenous NO (incubation with NOC-15). *tert*-BuOOH-induced oxoferryl-Hb-associated protein-centered free radical species, as well as *tert*-BuO[•]-alkoxyl radicals, were readily detectable in K/VP.5 cells, but were greatly reduced in K/VP.5-iNOS cells. Susceptibility to *tert*-BuOOH-induced cytotoxicity in K/VP.5-iNOS cells was increased by 24-h treatment with an iNOS inhibitor, L-N^G-monomethylarginine (L-NMA). NOC-15-mediated protection against *tert*-BuOOH-induced cytotoxicity in L-NMA-treated cells was mediated by direct NO redox interactions with iron catalytic sites, preventing "activation" of *tert*-BuOOH to its cytotoxic reactive species, but was diminished upon addition of NOC-15 to L-NMA-treated cells (72).

In most cells, both direct and indirect regulatory mechanisms of NO are likely operational as effective protective mechanisms against oxidative stress. The pathways that involve direct redox interactions of NO with iron catalytic sites likely represent a fast response that may be considered an emergency mechanism to protect cells from the consequences of both acute and long-term oxidative stress. An indirect regulatory pathway may function as an adaptive mechanism that becomes operational upon long-term exposure of cells to NO (72).

ACKNOWLEDGMENTS

This work was supported by NIH HL-64145-1A1, NIH CA 74972 and CA 77468, and the International Neurological Science Fellowship Program (F05 NS 10669) administered by NIH/NINDS in collaboration with WHO, Unit of Neuroscience, Division of Mental Health and Prevention of Substance Abuse.

ABBREVIATIONS

tert-BuOOH, *tert*-butyl hydroperoxide; DFO, deferoxamine; EPR, electron paramagnetic resonance; Hb, hemoglobin; Hb-4Fe³⁺, methemoglobin; iNOS, inducible nitric oxide synthase; IREs, iron responsive elements; IRP-1 and IRP-2, iron regulatory proteins 1 and 2; K/VP.5-iNOS cells, K/VP.5 cells transduced

with a retroviral vector containing the human iNOS gene; Mb, myoglobin; Mb-Fe³⁺, metmyoglobin; L-NMA, L-N^G-monomethylarginine; NO, nitric oxide; NOC-15, (Z)-1-[N-(3-ammoniopropyl)-N-(n-propyl)amino]diazene-1-ium-1,2-diolate; PC, phosphatidylcholine; PE, phosphatidylethanolamine; PI, phosphatidylinositol; PnA, *cis*-parinaric acid; PS, phosphatidylserine; VSMCs, vascular smooth muscle cells.

REFERENCES

1. Aisen P, and Listowsky I. Iron transport and storage proteins. *Annu Rev Biochem* 49: 357-393, 1980.
2. Aust SD, Morehouse LA, and Thomas CE. Role of metals in oxygen radical reactions. *Free Radic Biol Med* 1: 3-25, 1985.
3. Aziz N, and Munro HN. Iron regulates ferritin mRNA translation through a segment of its 5' untranslated region. *Proc Natl Acad Sci U S A* 84: 8478-8482, 1987.
4. Baron CP, Skibsted LH, and Andersen HJ. Peroxidation of linoleate at physiological pH: hemichrome formation by substrate binding protects against metmyoglobin activation by hydrogen peroxide. *Free Radic Biol Med* 28: 549-558, 2000.
5. Bloodsworth A, O'Donnell VB, and Freeman BA. Nitric oxide regulation of free radical- and enzyme-mediated lipid and lipoprotein oxidation. *Arterioscler Thromb Vasc Biol* 20: 1707-1715, 2000.
6. Cairo G, Tacchini L, Pogliaghi G, Anzon E, Tomasi A, and Bernelli-Zazzera A. Induction of ferritin synthesis by oxidative stress. Transcriptional and post-transcriptional regulation by expansion of the "free" iron pool. *J Biol Chem* 270: 700-703, 1995.
7. Ceccarelli D, Kozlov AV, Galesi D, Tomasi A, Giovannini F, and Masini A. The role of desferrioxamine chelatable iron in rat liver mitochondrial dysfunction in chronic dietary iron overload. *Bioelectrochem Bioenerg* 42: 169-174, 1997.
8. Cooper CE, Green ES, Rice-Evans CA, Davies MJ, and Wrigglesworth JM. A hydrogen-donating monohydroxamate scavenges ferryl myoglobin radicals. *Free Radic Res* 20: 219-227, 1994.
9. D'Agnillo F, and Chang TM. Absence of hemoprotein-associated free radical events following oxidant challenge of crosslinked hemoglobin-superoxide dismutase catalase. *Free Radic Biol Med* 24: 906-912, 1998.
10. Davies MJ. Detection of peroxy and alkoxyl radicals produced by reaction of hydroperoxides with heme proteins by electron spin resonance spectroscopy. *Biochim Biophys Acta* 964: 28-35, 1988.
11. Deterding LJ, Barr DP, Mason RP, and Tomer KB. Characterization of cytochrome *c* free radical reactions with peptides by mass spectrometry. *J Biol Chem* 273: 12863-12869, 1998.
12. de Montalembert M. [Iron chelation in 1998] La chelation du fer en 1998. *Transfus Clin Biol* 5: 353-356, 1998.

13. Domachowske JB. The role of nitric oxide in the regulation of cellular iron metabolism. *Biochem Mol Med* 60: 1–7, 1997.
14. Domachowske JB, Rafferty SP, Singhania N, Mardiney M 3rd, and Malech HL. Nitric oxide alters the expression of gamma-globin, H-ferritin, and transferrin receptor in human K562 cells at the posttranscriptional level. *Blood* 88: 2980–2988, 1996.
15. Dragsten PR, Hallaway PE, Hanson GJ, Berger AE, Bernard B, and Hedlund BE. First human studies with a high-molecular-weight iron chelator. *J Lab Clin Med* 135, 57–65, 2000.
16. Drapier JC, Hirling H, Wietzerbin J, Kaldy P, and Kuhn LC. Biosynthesis of nitric oxide activates iron regulatory factor in macrophages. *EMBO J* 12: 3643–3649, 1993.
17. Elihu N, Anandasbapathy S, and Fishman WH. Chelation therapy in cardiovascular disease: ethylenediaminetetraacetic acid, deferoxamine, and dexrazoxane. *J Clin Pharmacol* 32: 101–105, 1998.
18. Ferrali M, Signorini C, Ciccoli L, Bambagioni S, Rossi V, Pompella A, and Comporti M. Protection of erythrocytes against oxidative damage and autologous immunoglobulin G (IgG) binding by iron chelator fluor-benzoil-pyridoxal hydrazone. *Biochem Pharmacol* 59: 1365–1373, 2000.
19. Frieden E, and Hsieh HS. Ceruloplasmin: the copper transport protein with essential oxidase activity. *Adv Enzymol Relat Areas Mol Biol* 44: 187–236, 1976.
20. Geng YJ, Hellstrand K, Wennmalm A, and Hansson GK. Apoptotic death of human leukemic cells induced by vascular cells expressing nitric oxide synthase in response to gamma-interferon and tumor necrosis factor-alpha. *Cancer Res* 56: 866–874, 1996.
21. Giulivi C, Romero FJ, and Cadenas E. The interaction of trolox C, a water-soluble vitamin E analog, with ferrylmyoglobin: reduction of the oxoferryl moiety. *Arch Biochem Biophys* 299: 302–312, 1992.
22. Gorbunov NV, Osipov AN, Day BW, Zayas-Rivera B, Kagan VE, and Elsayed NM. Reduction of ferrylmyoglobin and ferrylhemoglobin by nitric oxide: a protective mechanism against ferryl hemoprotein-induced oxidations. *Biochemistry* 34: 6689–6699, 1995.
23. Gorbunov NV, Yalowich JC, Gaddam A, Thampatty P, Ritov VB, Kisin ER, Elsayed NM, and Kagan VE. Nitric oxide prevents oxidative damage produced by *tert*-butyl hydroperoxide in erythroleukemia cells via nitrosylation of heme and non-heme iron. Electron paramagnetic resonance evidence. *J Biol Chem* 272: 12328–12341, 1997.
24. Gorbunov NV, Tyurina YY, Salama G, Day BW, Claycamp HG, Argyros G, Elsayed NM, and Kagan VE. Nitric oxide protects cardiomyocytes against *tert*-butyl hydroperoxide-induced formation of alkoxyl and peroxy radicals and peroxidation of phosphatidylserine. *Biochem Biophys Res Commun* 244: 647–651, 1998.
25. Gunther MR, Sturgeon BE, and Mason RP. A long-lived tyrosyl radical from the reaction between horse metmyoglobin and hydrogen peroxide. *Free Radic Biol Med* 28: 709–719, 2000.
26. Guo B, Brown FM, Phillips JD, Yu Y, and Leibold EA. Characterization and expression of iron regulatory protein 2 (IRP2). Presence of multiple IRP2 transcripts regulated by intracellular iron levels. *J Biol Chem* 270: 16529–16535, 1995.
27. Halliwell B, and Gutteridge JM. Oxygen toxicity, oxygen radicals, transition metals and disease. *Biochem J* 219: 1–14, 1984.
28. Hentze MW, Caughman SW, Rouault TA, Barriocanal JG, Dancis A, Harford JB, and Klausner RD. Identification of the iron-responsive element for the translational regulation of human ferritin mRNA. *Science* 238: 1570–1573, 1987.
29. Herbert V, Shaw S, Jayatilleke E, and Stopler-Kasdan T. Most free-radical injury is iron-related: it is promoted by iron, heme, holoferritin and vitamin C, and inhibited by desferoxamine and apoferritin. *Stem Cells* 12: 289–303, 1994.
30. Hirling H, Henderson BR, and Kuhn LC. Mutational analysis of the [4Fe-4S]-cluster converting iron regulatory factor from its RNA-binding form to cytoplasmic aconitase. *EMBO J* 13: 453–461, 1994.
31. Hogg N, Kalyanaraman B, Joseph J, Struck A, and Parthasarathy S. Inhibition of low-density lipoprotein oxidation by nitric oxide. Potential role in atherogenesis. *FEBS Lett* 334: 170–174, 1993.
32. Hogg N, Rice-Evans C, Darley-Usmar V, Wilson MT, Paganga G, and Bourne L. The role of lipid hydroperoxides in the myoglobin-dependent oxidation of LDL. *Arch Biochem Biophys* 314: 39–44, 1994.
33. Iakutova ES, Osipov AN, Kostenko OV, Arnkhol'd I, Arnol'd K, and Vladimirov IuA. Interaction of hypochlorite with oxyhemoglobin leads to liberation of iron in a catalytically active form. [Russian] *Biofizika* 37: 1021–1028, 1992.
34. Iannone A, Tomasi A, and Canfield LM. Generation of *N-tert*-butyl-alpha-phenylnitron radical adducts in iron breakdown of *tert*-butyl-hydroperoxide. *Res Chem Intermed* 22: 469–479, 1996.
35. Irwin JA, Ostdal H, and Davies MJ. Myoglobin-induced oxidative damage: evidence for radical transfer from oxidized myoglobin to other proteins and antioxidants. *Arch Biochem Biophys* 362: 94–104, 1999.
36. Jacobs A. Low molecular weight intracellular iron transport compounds. *Blood* 50: 433–439, 1977.
37. Jacobs DL, Watt GD, Frankel RB, and Papaefthymiou GC. Redox reactions associated with iron release from mammalian ferritin. *Biochemistry* 28: 1650–1655, 1989.
38. Jinno J, Shigematsu M, Tajima K, Sakurai H, Ohya Nishiguchi H, and Ishizu K. Coordination structure and chemical reactivity of hemoprotein-butyl peroxide complex. *Biochem Biophys Res Commun* 176: 675–681, 1991.
39. Kang JH, and Kim SM. Fragmentation of human Cu,Zn-superoxide dismutase by peroxidative reaction. *Mol Cells* 7: 553–558, 1997.
40. Kanner J, Harel S, and Granit R. Nitric oxide as an antioxidant. *Arch Biochem Biophys* 289: 130–136, 1991.
41. Ke Y, Wu J, Leibold EA, Walden WE, and Theil EC.

- Loops and bulge/loops in iron-responsive element isoforms influence iron regulatory protein binding. Fine-tuning of mRNA regulation? *J Biol Chem* 273: 23637–23640, 1998.
42. Kim YM, Bergonia HA, Muller C, Pitt BR, Watkins WD, and Lancaster JR Jr. Nitric oxide and intracellular heme. *Adv Pharmacol* 34: 277–291, 1995.
 43. Korytowski W, Zareba M, and Girotti AW. Nitric oxide inhibition of free radical-mediated cholesterol peroxidation in liposomal membranes. *Biochemistry* 39: 6918–6928, 2000.
 44. Kozlov AV, Vladimirov IuA, and Azizova OA. The measurement of the amount of iron (III) complexes with desferal in the perfused rat liver by an EPR method. [Russian] *Biull Eksp Biol Med* 107: 577–579, 1989.
 45. Kozlov AV, Yegorov DY, Vladimirov YA, and Azizova OA. Intracellular free iron in liver tissue and liver homogenate: studies with electron paramagnetic resonance on the formation of paramagnetic complexes with desferal and nitric oxide. *Free Radic Biol Med* 13: 9–16, 1992.
 46. Loew GH, and Harris DL. Role of the heme active site and protein environment, structure, spectra, and function of the cytochrome P450s. *Chem Rev* 100: 407–419, 2000.
 47. McArthur KM, and Davies MJ. Detection and reactions of the globin radical in hemoglobin. *Biochim Biophys Acta* 1202: 173–181, 1993.
 48. Mehlhorn RJ, and Gomez J. Hydroxyl and alkoxy radical production by oxidation products of metmyoglobin. *Free Radic Res Commun* 18: 29–41, 1993.
 49. Menotti E, Henderson BR, and Kuhn LC. Translational regulation of mRNAs with distinct IRE sequences by iron regulatory proteins 1 and 2. *J Biol Chem* 273: 1821–1824, 1998.
 50. Moore KP, Holt SG, Patel RP, Svistunenko DA, Zackert W, Goodier D, Reeder BJ, Clozel M, Anand R, Cooper CE, Morrow JD, Wilson MT, Darley-Usmar V, and Roberts LJ. A causative role for redox cycling of myoglobin and its inhibition by alkalinization in the pathogenesis and treatment of rhabdomyolysis-induced renal failure. *J Biol Chem* 273: 31731–31737, 1998.
 51. Ostdal H, Andersen HJ, and Davies MJ. Formation of long-lived radicals on proteins by radical transfer from heme enzymes—a common process? *Arch Biochem Biophys* 362: 105–112, 1999.
 52. Panter SS. Release of iron from hemoglobin. *Methods Enzymol* 231: 502–514, 1994.
 53. Pantopoulos K, and Hentze MW. Nitric oxide signaling to iron-regulatory protein: direct control of ferritin mRNA translation and transferrin receptor mRNA stability in transfected fibroblasts. *Proc Natl Acad Sci U S A* 92: 1267–1271, 1995.
 54. Paraskeva E, and Hentze MW. Iron-sulphur clusters as genetic regulatory switches: the bifunctional iron regulatory protein-1. *FEBS Lett* 389: 40–43, 1996.
 55. Patel RP, Svistunenko DA, Darley-Usmar VM, Symons MC, and Wilson MT. Redox cycling of human methemoglobin by H₂O₂ yields persistent ferryl iron and protein based radicals. *Free Radic Res* 25: 117–123, 1996.
 56. Rafferty SP, Domachowske JB, and Malech HL. Inhibition of hemoglobin expression by heterologous production of nitric oxide synthase in the K562 erythroleukemic cell line. *Blood* 88: 1070–1078, 1996.
 57. Reif DW. Ferritin as a source of iron for oxidative damage. *Free Radic Biol Med* 12: 417–427, 1992.
 58. Richardson DR, and Ponka P. Development of iron chelators to treat iron overload disease and their use as experimental tools to probe intracellular iron metabolism. *Am J Hematol* 58: 299–305, 1998.
 59. Ritke MK, Roberts D, Allan WP, Raymond J, Bergottz VV, and Yalowich JC. Altered stability of etoposide-induced topoisomerase II–DNA complexes in resistant human leukemia K562 cells. *Br J Cancer* 69: 687–697, 1994.
 60. Rubbo H, Radi R, Trujillo M, Telleri R, Kalyanaraman B, Barnes S, Kirk M, and Freeman BA. Nitric oxide regulation of superoxide and peroxynitrite-dependent lipid peroxidation. Formation of novel nitrogen-containing oxidized lipid derivatives. *J Biol Chem* 269: 26066–26075, 1994.
 61. Samaniego F, Chin J, Iwai K, Rouault TA, and Klausner RD. Molecular characterization of a second iron-responsive element binding protein, iron regulatory protein 2. Structure, function, and post-translational regulation. *J Biol Chem* 269: 30904–30910, 1994.
 62. Shvedova AA, Tyurina YY, Gorbunov NV, Tyurin VA, Castranova V, Kommineni C, Ojimba J, Gandley R, McLaughlin MK, and Kagan VE. *tert*-Butyl hydroperoxide/hemoglobin-induced oxidative stress and damage to vascular smooth muscle cells: different effects of nitric oxide and nitrosothiols. *Biochem Pharmacol* 57: 989–1001, 1999.
 63. Sigurdsson L, Reyes J, Kocoshis SA, Hansen TW, Rosh J, and Knisely AS. Neonatal hemochromatosis: outcomes of pharmacologic and surgical therapies. *J Pediatr Gastroenterol Nutr* 26: 85–89, 1998.
 64. Stamler JS, Singel DJ, and Loscalzo J. Biochemistry of nitric oxide and its redox-activated forms. *Science* 258: 1898–1902, 1992.
 65. Stolze K, and Nohl H. Reactions of reducing xenobiotics with oxymyoglobin. Formation of metmyoglobin, ferryl myoglobin and free radicals: an electron spin resonance and chemiluminescence study. *Biochem Pharmacol* 49: 1261–1267, 1995.
 66. Svistunenko DA, Patel RP, and Wilson MT. An EPR investigation of human methaemoglobin oxidation by hydrogen peroxide: methods to quantify all paramagnetic species observed in the reaction. *Free Radic Res* 24: 269–280, 1996.
 67. Tajima K. A possible model of a hemoprotein–hydrogen peroxide complex. *Inorg Chim Acta* 163: 115–122, 1989.
 68. Theil EC. Iron regulatory elements (IREs): a family of mRNA non-coding sequences. *Biochem J* 304: 1–11, 1994.
 69. Thomson AM, Rogers JT, and Leedman PJ. Iron-regulatory proteins, iron-responsive elements and ferritin mRNA translation. *Int J Biochem Cell Biol* 31: 1139–1152, 1999.

70. Vanin AF, and Chetverikov AG. Paramagnetic nitrosyl complexes of heme and nonheme iron. [Russian] *Biofizika* 13: 608–615, 1968.
71. Vanin AF, Bliumenfel'd LA, and Chetverikov AG. EPR study of non-heme iron complexes in cells and tissues. [Russian] *Biofizika* 12: 829–838, 1967.
72. Yalowich JC, Gorbunov NV, Kozlov AV, Allan W, and Kagan VE. Mechanisms of nitric oxide protection against *tert*-butyl hydroperoxide-induced cytotoxicity in iNOS-transduced human erythroleukemia cells. *Biochemistry* 38: 10691–10698, 1999.
73. Yamakura F, Rardin RL, Petsko GA, Ringe D, Hiraoka BY, Nakayama K, Fujimura T, Taka H, and Murayama K. Inactivation and destruction of conserved Trp159 of Fe-superoxide dismutase from *Porphyromonas gingivalis* by hydrogen peroxide. *Eur J Biochem* 253: 49–56, 1998.

Address reprint requests to:
Dr. Valerian E. Kagan
Department of Environmental and
Occupational Health
University of Pittsburgh
260 Kappa Dr.
RIDC Park
Pittsburgh, PA 15238, U.S.A.

E-mail: kagan@pitt.edu

Received for publication October 18, 2000; accepted December 10, 2000.

This article has been cited by:

1. Andrey V Kozlov, Soheyl Bahrami, Heinz Redl, Csaba Szabo. 2017. Alterations in nitric oxide homeostasis during traumatic brain injury. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* . [[CrossRef](#)]
2. Peter Dungal, Martin Perlinger, Adelheid Weidinger, Heinz Redl, Andrey V. Kozlov. 2015. The cytoprotective effect of nitrite is based on the formation of dinitrosyl iron complexes. *Free Radical Biology and Medicine* **89**, 300-310. [[CrossRef](#)]
3. Hanna Lewandowska, Jarosław Sadło, Sylwia Moczyńska, Tomasz M. Stępkowski, Grzegorz Wójciuk, Marcin Kruszewski. 2015. Formation of glutathionyl dinitrosyl iron complexes protects against iron genotoxicity. *Dalton Trans.* **44**:28, 12640-12652. [[CrossRef](#)]
4. Qian Li, Chuanyu Li, Harry K. Mahtani, Jian Du, Aashka R. Patel, Jack R. Lancaster. 2014. Nitrosothiol Formation and Protection against Fenton Chemistry by Nitric Oxide-induced Dinitrosyliron Complex Formation from Anoxia-initiated Cellular Chelatable Iron Increase. *Journal of Biological Chemistry* **289**:29, 19917-19927. [[CrossRef](#)]
5. Reeta Yadav, Yuval Samuni, Alex Abramson, Rephael Zeltser, Nardi Casap, Tonmoy K. Kabiraj, Maureen L. Banach, Uri Samuni. 2014. Pro-oxidative synergic bactericidal effect of NO: kinetics and inhibition by nitroxides. *Free Radical Biology and Medicine* **67**, 248-254. [[CrossRef](#)]
6. Taija S. Koskenkorva-Frank, Günter Weiss, Willem H. Koppenol, Susanna Burckhardt. 2013. The complex interplay of iron metabolism, reactive oxygen species, and reactive nitrogen species: Insights into the potential of various iron therapies to induce oxidative and nitrosative stress. *Free Radical Biology and Medicine* **65**, 1174-1194. [[CrossRef](#)]
7. Jack R. Lancaster, Gregory I. Giles. Nitrogen Monoxide (Nitric Oxide): Bioinorganic Chemistry . [[CrossRef](#)]
8. Ivan Rousseau, Monica Galleano, Susana Puntarulo. 2011. Fe Allocation in Liver during Early Stages of Endotoxemia in Fe-Overload Rats. *Toxicologic Pathology* **39**:7, 1075-1083. [[CrossRef](#)]
9. S.F. Medeiros, A.M. Santos, H. Fessi, A. Elaissari. 2011. Stimuli-responsive magnetic particles for biomedical applications. *International Journal of Pharmaceutics* **403**:1-2, 139-161. [[CrossRef](#)]
10. Jun Zhao, Tao Liu, Long Ma, Ming Yan, Zhengyi Gu, Yi Huang, Fang Xu, Yu Zhao. 2011. Antioxidant and Preventive Effects of Extract from *Nymphaea candida* Flower on In Vitro Immunological Liver Injury of Rat Primary Hepatocyte Cultures. *Evidence-Based Complementary and Alternative Medicine* **2011**, 1-8. [[CrossRef](#)]
11. José C. Toledo, Charles A. Bosworth, Seth W. Hennon, Harry A. Mahtani, Hector A. Bergonia, Jack R. Lancaster. 2008. Nitric Oxide-induced Conversion of Cellular Chelatable Iron into Macromolecule-bound Paramagnetic Dinitrosyliron Complexes. *Journal of Biological Chemistry* **283**:43, 28926-28933. [[CrossRef](#)]
12. K CHANDRASHEKAR, MURALIDHARA. 2008. Oxidative alterations induced by d-aspartic acid in prepubertal rat testis in vitro: A mechanistic study#. *Theriogenology* **70**:1, 97-104. [[CrossRef](#)]
13. A GUPTA, S SHARMA, K CHOPRA. 2008. Reversal of iron-induced nephrotoxicity in rats by molsidomine, a nitric oxide donor. *Food and Chemical Toxicology* **46**:2, 537-543. [[CrossRef](#)]
14. Konstantin B. Shumaev, Olga V. Kosmachevskaya, Alexander A. Timoshin, Anatoly F. Vanin, Alexey F. Topunov. Dinitrosyl Iron Complexes Bind with Hemoglobin as Markers of Oxidative Stress 445-461. [[CrossRef](#)]
15. Shino Magaki, Claudius Mueller, Steven M. Yellon, James Fox, Joseph Kim, Eugene Snissarenko, Vernon Chin, Manik C. Ghosh, Wolff M. Kirsch. 2007. Regional dissection and determination of loosely bound and non-heme iron in the developing mouse brain. *Brain Research* **1158**, 144-150. [[CrossRef](#)]
16. L. L. Gudkov, K. B. Shumaev, E. I. Kalenikova, S. A. Gubkina, A. F. Vanin, E. K. Ruuge. 2007. Antioxidant and prooxidant action of nitric oxide donors and metabolites. *Biophysics* **52**:3, 315-321. [[CrossRef](#)]
17. A. Biliska, M. Dubiel, M. Sokołowska-Jezewicz, E. Lorenc-Koci, L. Włodek. 2007. Alpha-lipoic acid differently affects the reserpine-induced oxidative stress in the striatum and prefrontal cortex of rat brain. *Neuroscience* **146**:4, 1758-1771. [[CrossRef](#)]

18. A GUPTA, V CHANDER, S SHARMA, K CHOPRA. 2007. Sodium nitroprusside and l-arginine attenuates ferric nitrilotriacetate-induced oxidative renal injury in rats. *Toxicology* **232**:3, 183-191. [[CrossRef](#)]
19. P CORNEJO, V FERNANDEZ, M VIAL, L VIDELA. 2007. Hepatoprotective role of nitric oxide in an experimental model of chronic iron overload. *Nitric Oxide* **16**:1, 143-149. [[CrossRef](#)]
20. A VANIN, N SANINA, V SEREZHENKOV, D BURBAEV, V LOZINSKY, S ALDOSHIN. 2007. Dinitrosyl-iron complexes with thiol-containing ligands: Spatial and electronic structures. *Nitric Oxide* **16**:1, 82-93. [[CrossRef](#)]
21. K. B. Shumaev, A. A. Gubkin, S. A. Gubkina, L. L. Gudkov, I. V. Sviryaeva, A. A. Timoshin, A. F. Topunov, A. F. Vanin, E. K. Ruuge. 2006. The interaction between dinitrosy iron complexes and intermediates of oxidative stress. *Biophysics* **51**:3, 423-428. [[CrossRef](#)]
22. Irina I. Vlasova, Vladimir A. Tyurin, Alexandr A. Kapralov, Igor V. Kurnikov, Anatoly N. Osipov, Maxim V. Potapovich, Detcho A. Stoyanovsky, Valerian E. Kagan. 2006. Nitric Oxide Inhibits Peroxidase Activity of Cytochrome c · Cardiolipin Complex and Blocks Cardiolipin Oxidation. *Journal of Biological Chemistry* **281**:21, 14554-14562. [[CrossRef](#)]
23. Jack R. Lancaster, Gregory I. Giles. Nitrogen Monoxide (Nitric Oxide): Bioinorganic Chemistry . [[CrossRef](#)]
24. Changyuan Lu, Willem H. Koppenol. 2005. Inhibition of the Fenton reaction by nitrogen monoxide. *JBIC Journal of Biological Inorganic Chemistry* **10**:7, 732-738. [[CrossRef](#)]
25. Susana Puntarulo. 2005. Iron, oxidative stress and human health. *Molecular Aspects of Medicine* **26**:4-5, 299-312. [[CrossRef](#)]
26. Pamela Cornejo, Patricia Varela, Luis A. Videla, Virginia Fernández. 2005. Chronic iron overload enhances inducible nitric oxide synthase expression in rat liver. *Nitric Oxide* **13**:1, 54-61. [[CrossRef](#)]
27. Elżbieta Lorenc-Koci, Maria Sokołowska, Inga Kwiecień, Lidia Włodek. 2005. Treatment with 1,2,3,4-tetrahydroisoquinolone affects the levels of nitric oxide, S-nitrosothiols, glutathione and the enzymatic activity of γ -glutamyl transpeptidase in the dopaminergic structures of rat brain. *Brain Research* **1049**:2, 133-146. [[CrossRef](#)]
28. Gaia Rocchitta, Rossana Migheli, Maria P. Mura, Giuseppe Grella, Giovanni Esposito, Bianca Marchetti, Egidio Miele, Maria S. Desole, Maddalena Miele, Pier Andrea Serra. 2005. Signaling pathways in the nitric oxide and iron-induced dopamine release in the striatum of freely moving rats: Role of extracellular Ca²⁺ and L-type Ca²⁺ channels. *Brain Research* **1047**:1, 18-29. [[CrossRef](#)]
29. Shahrzad Tafazoli, Dane D. Spehar, Peter J. O'Brien. 2005. Oxidative Stress Mediated Idiosyncratic Drug Toxicity. *Drug Metabolism Reviews* **37**:2, 311-325. [[CrossRef](#)]
30. Kenneth J. Smith. Nitric Oxide and Axonal Pathophysiology 255-273. [[CrossRef](#)]
31. Srdjan Stojanović, Dragana Stanić, Milan Nikolić, Mihailo Spasić, Vesna Niketić. 2004. Iron catalyzed conversion of NO into nitrosonium (NO⁺) and nitroxyl (HNO/NO⁻) species. *Nitric Oxide* **11**:3, 256-262. [[CrossRef](#)]
32. 2003. Trend of Most Cited Papers (2001-2002) in ARS. *Antioxidants & Redox Signaling* **5**:6, 813-815. [[Citation](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]
33. Martyn A. Sharpe, Sarah J. Robb, John B. Clark. 2003. Nitric oxide and Fenton/Haber-Weiss chemistry: nitric oxide is a potent antioxidant at physiological concentrations. *Journal of Neurochemistry* **87**:2, 386-394. [[CrossRef](#)]
34. Akira I. Hida, Teruyuki Kawabata, Yukiko Minamiyama, Akiko Mizote, Shigeru Okada. 2003. Saccharated colloidal iron enhances lipopolysaccharide-induced nitric oxide production in vivo. *Free Radical Biology and Medicine* **34**:11, 1426-1434. [[CrossRef](#)]
35. P FORD, L LAVERMAN, I LORKOVIC. REACTION MECHANISMS OF NITRIC OXIDE WITH BIOLOGICALLY RELEVANT METAL CENTERS 203-257. [[CrossRef](#)]
36. John J Haddad. 2002. Pharmacoredox regulation of cytokine-related pathways: from receptor signaling to pharmacogenomics. *Free Radical Biology and Medicine* **33**:7, 907-926. [[CrossRef](#)]

37. Reto Huber, Tom Deboer, Irene Tobler. 2002. Sleep deprivation in prion protein deficient mice and control mice: genotype dependent regional rebound. *Neuroreport* 13:1, 1-4. [[CrossRef](#)]