

Work Stress, Sleep Deficiency, and Predicted 10-Year Cardiometabolic Risk in a Female Patient Care Worker Population

Henrik B. Jacobsen, PsyD,^{1,2*} Silje E. Reme, PsyD, PhD,^{1,3} Grace Sembajwe, MSc, ScD,^{4,5} Karen Hopcia, NP,^{1,5} Tore C. Stiles, PsyD, PhD,² Glorian Sorensen, MPH, PhD,^{1,6,7} James H. Porter, BS,⁸ Miguel Marino, PhD,⁹ and Orfeu M. Buxton, PhD^{1,8,10}

Objectives The aim of this study was to investigate the longitudinal effect of work-related stress, sleep deficiency, and physical activity on 10-year cardiometabolic risk among an all-female worker population.

Methods Data on patient care workers ($n=99$) was collected 2 years apart. Baseline measures included: job stress, physical activity, night work, and sleep deficiency. Biomarkers and objective measurements were used to estimate 10-year cardiometabolic risk at follow-up. Significant associations ($P < 0.05$) from baseline analyses were used to build a multivariable linear regression model.

Results The participants were mostly white nurses with a mean age of 41 years. Adjusted linear regression showed that having sleep maintenance problems, a different occupation than nurse, and/or not exercising at recommended levels at baseline increased the 10-year cardiometabolic risk at follow-up.

Conclusions In female workers prone to work-related stress and sleep deficiency, maintaining sleep and exercise patterns had a strong impact on modifiable 10-year cardiometabolic risk. Am. J. Ind. Med. 57:940–949, 2014. © 2014 Wiley Periodicals, Inc.

KEY WORDS: cardiometabolic risk; nurses; sleep maintenance; physical activity; follow-up; work-family conflict

¹Harvard School of Public Health, Center for Work, Health and Wellbeing, Boston, Massachusetts

²The Norwegian University of Science and Technology, Trondheim, Norway

³Uni Health, Uni Research, Bergen, Norway

⁴CUNY School of Public Health at Hunter College, New York, New York

⁵Partners Occupational Health Service, Partners Health Care System, Boston, Massachusetts

⁶Department of Society, Health and Human Development, Harvard School of Public Health, Boston, Massachusetts

⁷Dana Farber Cancer Institute, Boston, Massachusetts

⁸Department of Medicine, Brigham and Women's Hospital, Boston, Massachusetts

⁹Department of Family Medicine, Public Health & Preventive Medicine, Division of Biostatistics, Oregon Health & Science University, Portland, Oregon

¹⁰Division of Sleep Medicine, Harvard Medical School, Boston, Massachusetts

Contract grant sponsor: National Institute for Occupational Safety and Health; Contract grant number: U19 OH008861; Contract grant sponsor: National Center for Research Resources; Grant number; Contract grant number: UL1 RR025758-04; Contract grant sponsor: Work, Family, and Health Network; Contract grant number: U01 AG5186989; Contract grant sponsor: National Heart, Lung, and Blood Institute; Contract grant number: R01 HL107240; Contract grant sponsor: Dean Hashimoto and Partners Occupational Health Service.

Disclosure Statement: The authors report no conflicts of interests.

*Correspondence to: Henrik Børsting Jacobsen, PsyD, NTNU, Faculty of Medicine, Department of Circulation and Medical Imaging, Po.box 8905 MTFS, 7491 Trondheim, Norway.
E-mail: henrik.b.jacobsen@ntnu.no

INTRODUCTION

Cardiovascular diseases are the leading cause of adult death in the world, with approximately 32% and 27% of adult female and male fatalities in 2004 attributed to cardiovascular diseases [Mathers et al., 2008]. In the U.S., recent statistics show that heart diseases and strokes caused 34% of all deaths and represented a total annual cost of 475.3 billion dollars; making cardiovascular disease the foremost cause of death and the third-most expensive disease in America [Roger et al., 2011]. The last decade has seen both increased awareness and reduction of the incidence and prevalence of cardiovascular disease, but it remains a major burden [Backe et al., 2012].

Recent research has identified sleep quality and sleep duration as important factors in cardiovascular disease risk [Buxton and Marcelli, 2010; Laugsand et al., 2013]. Indeed, insufficient sleep duration and poor sleep quality appear to contribute to increased cardiovascular disease risk [Buxton and Marcelli, 2010; Cappuccio et al., 2010], and have been linked to elevated body mass index [Hasler et al., 2004; Kohatsu et al., 2006], weight gain [Patel et al., 2004, 2008], obesity [Taheri et al., 2004; Cizza et al., 2005; Gangwisch et al., 2005; Buxton et al., 2013], and diabetes mellitus [Ayas et al., 2003a; Gottlieb et al., 2005; Buxton et al., 2013]. Additionally, modulations of cardiovascular function by sleep duration have been demonstrated showing elevations in blood pressure [Lusardi et al., 1999; Meier-Ewert et al., 2004], which is consistent with observed associations of sleep duration with blood pressure [Gottlieb et al., 2006; Stranges et al., 2010]. Perhaps most importantly, past research has shown that short sleep (<6 hr per day) predicts premature mortality [Wingard and Berkman, 1983; Kripke et al., 2002; Dew et al., 2003; Patel et al., 2004; Grandner et al., 2010]. Long sleep duration (>8 hr per day) may also be associated with cardiovascular risk [Hale and Do, 2007; Gallicchio and Kalesan, 2009; Buxton and Marcelli, 2010].

Shift work, particularly night shift work, has been associated with sleep deficiency and has shown a dose-response relationship with coronary heart disease [Kawachi et al., 1995; Åkerstedt, 2003]. A vulnerable population with regards to both sleep deficiency and shift work are patient care workers [Kawachi et al., 1995; Buxton et al., 2012]. A recent study demonstrated that 57% of a large patient care worker cohort reported sleep deficiency, specifically a lack of quality, duration and/or perceived sufficiency of sleep [Buxton et al., 2012]. The Nurses Health Study showed a 4% increase in the hazard ratio for ischemic heart disease with every 5 years nurses worked rotating night shifts [Brown et al., 2009]. Furthermore, a large-scale, prospective study on the same population of married nurses demonstrated a significant increase in mortality risk in those who reported either short or long sleep duration [Patel et al., 2004]. This increased risk was associated with hypertension [Patel

et al., 2004], and other studies have shown the independent association of short and long sleep duration with coronary events in this cohort [Ayas et al., 2003b].

Patient care workers also report significantly higher psychosocial stress than other occupational groups [Evans and Steptoe, 2002]. When compared with other workers, nurses report more job strain due to higher demands and lower skill utilization [Evans and Steptoe, 2002]. In addition, patient care workers report high levels of work-family conflict, an important cause of perceived stress [Kim et al., 2012].

There is a well-established association between work-related stress and risk of cardiovascular disease [Eller et al., 2009; Schnall et al., 2009; Backe et al., 2012], but the effect has been claimed to be significantly less than that of traditional risk factors like amount of physical activity [Kivimaki et al., 2012; Steptoe and Kivimäki, 2012].

The concept of work-related stress is constantly evolving. A recent review highlights the need for a broader understanding of how it relates to cardiovascular risk [Backe et al., 2012]. Work-family conflict has been identified as a major cause of work-related stress in several studies, and has been implicated as a contributing factor to cardiovascular risk [Bellavia and Frone, 2005; Berkman et al., 2010]. The conflict of roles at work and in one's personal life is recognized as an independent factor from job strain and effort-reward imbalance [Netemeyer et al., 1996; Bhui et al., 2012], and might contribute to our understanding of the psychosocial work-environment in cardiovascular risk research [Frone et al., 2011]. Two recent systematic reviews on cardiovascular risk and workplace stressors also underline the need for more longitudinal studies on female populations [Eller et al., 2009; Backe et al., 2012]. The association between high demands, low control and low co-worker support (iso-strain) at work and cardiovascular risk is consistently found in men [Belkic et al., 2004], but the evidence in women is less conclusive [Backe et al., 2012]. To address this, we focus the current analysis on a sample of women.

A study on a patient worker population combining several different measures of work stress controlled by other confounding factors such as physical activity [Hublin et al., 2010], sleep duration [Chandola et al., 2010], and night-work hours [Yang et al., 2006].

In the current study we test the hypothesis that work-related stress, sleep deficiency, and physical activity over time influences modifiable 10-year cardiometabolic risk in a female worker population. More specifically, we investigate the predictive associations of baseline characteristics on modifiable cardiometabolic risk factors in a female worker population, and investigate the longitudinal effects of iso-strain, work-family conflict, night work, physical activity and sleep deficiency on a non-self reported, modifiable cardiometabolic risk score [Marino et al., 2014], in a female patient care worker population.

MATERIALS AND METHODS

Study Design

This study had a longitudinal design where data was collected at two time-points from a group of patient care workers at a large hospital in Boston. Baseline collection spanned from October 2009 to January 2010 and the second was a follow-up from August to November 2011 (Fig. 1). All participants provided written informed consent. Research protocols were approved by the designated institutional review board and consistent with the principles in the Declaration of Helsinki.

Study Sample and Data Collection

Data collection at baseline has been described in previous articles [Sorensen et al., 2011; Buxton et al., 2012; Kim et al., 2012]. In brief, the data presented at baseline was collected through a cross-sectional survey of patient care workers at two major hospitals in 2009. In this initial data collection, we randomly selected 2000 eligible workers and invited them via e-mail to participate in the survey on-line. Of these patient care workers initially contacted, 1,572 participants completed at least 50% of the survey. When collecting follow-up data, only employees from one of the two hospitals were contacted. Employees who completed the initial survey at this hospital ($n=840$) were randomized and divided into six equal groups. Each group received an initial contact via e-mail, reminding employees of their participation in the original survey and asking about their interest in the second phase of the study. If not responsive after the initial contact, employees received up to seven additional contacts via email. A small number of subjects who expressed interest but subsequently became unresponsive were re-contacted by phone. Subjects who completed phase two of the study were compensated \$100. Out of 840 employees who were re-contacted after

completing the survey, a total of 99 (11.8%) females completed the phase-two visit and blood draw. A low participation rate at time two was expected as this was a much more comprehensive sampling with in-person collection of both biomarkers and self-report data. Test statistics (frequency distributions and mean differences) showed that these responders were similar to the overall sample on important baseline socio-demographic characteristics, with race/ethnicity (described below) being the only significant difference.

Measures

Registered nurses measured height, weight, and blood pressure, and collected blood samples. Blood pressure was taken using a calibrated, clinical-standard arm cuff (Welch Allyn Spot Vital Signs monitor model# 4200B-E1) and systolic blood pressure (SBP) was used as a continuous measure. Body mass index (BMI) was calculated as weight (kg) per meter squared (m^2) of height. Subjects were given a smoking status as current smoker if they answered yes to the following question “Have you smoked a cigarette, even a puff, in the last 7 days? (Yes; No).” Venous blood samples provided by all 99 subjects were assayed to determine cholesterol and glycosylated hemoglobin values.

Assay Details

All assays at follow-up were performed by CLIA certified laboratories. Glycosylated hemoglobin (Hb A_{1C}) was assayed using a Roche P-Modular Tina-Quant Immuno-assay, with an intra-assay precision coefficient of variation (CV) of 0.8–1.5%, an inter-assay CV of 1.3–2.0%, and a lower limit of detection of 2.9%. Total cholesterol was measured enzymatically in serum using a Roche/Hitachi analyzer with an intra-assay CV of 0.8%, an inter-assay CV of 1.7%, and a lower limit of detection of 3 mg/dl. HDL cholesterol was measured enzymatically in serum via a Roche/Hitachi analyzer with an intra-assay CV of 0.60–




FIGURE1. Participant flow and a conceptual model of the study, showing how baseline characteristics are hypothesized to predict 10-year cardiometabolic risk nearly 2 years later.

0.95%, an inter-assay CV of 1.2–1.3%, and a lower limit of detection of 3 mg/dl.

Assessment of 10-Year Cardiometabolic Risk

Cardiometabolic 10-year risk was assessed based on five non-self report modifiable cardiometabolic risk factors, initially developed in the Framingham Study [Wilson et al., 1998] modified by D'Agostino et al. [2008], and further developed by Marino et al. in 2013 using the Framingham offspring study to focus solely on modifiable factors [Marino et al., 2014]. The most recent modification evaluates risk through the addition of Hb A_{1C} SBP and BMI. In brief, while controlling for age and in gender-stratified risk models, the model assessed current smoking, and continuous total cholesterol, HDL-cholesterol, SBP, HbA1c levels, and BMI. Details of the model are described elsewhere [Marino et al., 2014]. Each of the items in the component score was kept continuous to keep them sensitive to small changes and to reflect the potential impact of interventions. Some (n = 2) of the participants had missing data on more than one of these risk factors and were excluded from analyses.

Covariates

The covariates were determined a priori and reported at baseline. We selected covariates that have been associated with cardiovascular health, sleep, and psychosocial stress. All covariates have been described in previous studies [Sorensen et al., 2011; Buxton et al., 2012; Kim et al., 2012].

Socio-demographic factors were obtained through participants reporting their age (years), gender, race/ethnicity (Hispanic, White, Black, and mixed race/others), occupation (staff nurse, patient care associate (PCA) and others), education (GED or less; Some College; College Degree; Graduate School), ability to pay bills (great deal of difficulty; some difficulty; a little difficulty; no difficulty; don't know; refused), height (inches), and weight (pounds). Body mass index (BMI) was measured by a nurse at follow-up using weight and height (kilograms per square meter).

Night work was quantified from detailed administrative payroll data and calculated as average night work-hours per month (between 10 PM and 6 AM), calculated from October 2008 until August 2009, making the assessment a year before initiating the survey for all workers. Excluding shifts shorter than 4 hr, the variable was trichotomized into 0–6 hr, 6–72 hr, and more than 72 hr per month, over the past year, during months worked, as described previously [Buxton et al., 2012].

Work-related stress was assessed by self-reported job demands, decision latitude, co-worker/supervisor support, and work-family conflict. A modified version of the Job

Content Questionnaire [Karasek, 1979; Karasek et al., 1998; Lusardi et al., 1999] measured job demands, decision latitude, and co-worker/supervisor support. Job demands were assessed through five items that were weighted and summed yielding a scale from 12 to 48 [Karasek et al., 1998]. Decision latitude was assessed through nine items and created as a weighted sum of decision authority and skill discretion from the Job Content Questionnaire [Karasek et al., 1998]. Co-worker/supervisor support was assessed through five items (three and two, respectively) with five response categories summed, yielding a scale from 2 to 10 [Karasek et al., 1998]. Supervisor support was summed and scaled giving a scale of 3–15 [Karasek et al., 1998]. Social support was defined as the sum of co-worker and supervisor support. Iso-strain was a composite variable assessed by combining the risk categories job demands, decision latitude, and social support and dichotomizing these three variables into low/high categories [Harvey et al., 2011; Bhui et al., 2012]. Iso-strain was defined as the combination of high job demands, low control, and low social support [Bhui et al., 2012].

Work-family conflict was assessed by the work-family conflict scale [Netemeyer et al., 1996] consisting of five items with response categories from 1 to 5 summed, scaled, and giving a range of 5–25: (1) "The demands of my work interfere with my family or personal time." (2) "The amount of time my job takes up makes it difficult to fulfill family or personal responsibilities." (3) "Things I want to do at home do not get done because of the demands my job puts on me." (4) "My job produces strain that makes it difficult to fulfill my family or personal duties." (5) "Due to work-related duties, I have to make changes to my plans for family or personal activities." Response categories ranged from 5 = "Strongly Agree" to 1 = "Strongly Disagree," making a higher score an indicator of more work-family conflict.

Physical activity (PA) outside of work was assessed with a measure adapted from the Centers for Disease Control and Prevention Behavioral Risk Factor and Surveillance System Physical Activity measure [Centers for Disease Control and Prevention, 2009]. Respondents were asked about their participation in vigorous and moderate PA of at least 10 min' duration outside of work. Each activity was defined by number of days per week and the total duration (hours and minutes) per day. Recommended PA was defined as reporting at least 30 min of moderate or vigorous activity on at least 5 days a week or at least 20 min of vigorous activity on at least 3 days a week [US Department of Health and Human Services, 2010].

Sleep deficiency [Luyster et al., 2012] in the last 4 weeks was assessed by self-report. Sleep was measured using three self-reported variables: sleep duration, sleep quality, and sleep insufficiency. Short sleep duration was assessed by asking subjects how many hours of sleep they got each night over the past 4 weeks [Sorensen et al., 2011; Buxton et al., 2012]. Sleep quality was determined based

on the frequency of sleep maintenance problems, assessed by asking subjects how often they woke up in the middle of the night or early morning, with four response categories: “not during the past month”; “Less than once per week”; “Once or twice per week”; or “three or more times a week” [Buysse et al., 1989]. Sleep insufficiency was determined based on subject response on a five-component scale ranging from “never” to “always” to the question “how often during the past 4 weeks did you get enough sleep to feel rested upon waking up?” [Sorensen et al., 2011], similar to the Centers for Disease Control measures used in the BRFSS study to assess U.S. national sleep sufficiency by state [McKnight-Eily et al., 2008]. Sleep outcomes were dichotomized into “Yes/No” based on participant response on the component scales. In brief, a response of <6 hr per night yielded a “Yes” on short sleep duration, a response of “three or more times a week” for interrupted sleep yielded a “Yes” on sleep maintenance problems and a response of “always” or “sometimes” yielded a “Yes” on sleep insufficiency. Sleep deficiency was defined as having responded “Yes” to one or more of these sleep outcomes. The distribution and overlap of the sleep variables are illustrated in Figure 2.

Data Analysis

In initial analyses, the characteristics of workers were compared on 10-year cardiometabolic risk score (%) treated continuously for analysis purposes. For dichotomous variables we used the independent sample *t*-test to compare means. For ordinal and continuously measured characteristics we used simple linear regression or a one-way ANOVA.

Sleep Deficiency, 1 of any component; n=61, 62%




FIGURE 2. Participant distribution and overlap on baseline sleep outcomes. Sleep maintenance is represented by the blue square, short sleep duration (<6 hr) by the red square and sleep insufficiency by the yellow square.

Significant variables from baseline analyses were entered into a multivariable linear regression. As baseline assessments only had self-report measures, no measure of risk change is included in the analyses.

RESULTS

Our participants (n=99) were predominantly white (91%), female (100%), nurses (68%) with a college degree (65%) and a mean age of 40.8 (SD 11.9, range 21–62) years. Fourteen (13.7%) of total respondents had iso-strain, 61 (59%) met recommended levels of PA, and 64 (63%) had sleep deficiency. The participants’ score on the individual cardiometabolic risk factors demonstrated means and standard deviations of 112.1 mmHg (SD 12.1) on SBP, 5.6% (SD 0.68) on Hb A_{1C}, 26.2 kg/m² (SD 5.7, range 17.6–46.1) on BMI and 67.6 mg/dl (SD 16.7) on HDL cholesterol. The participant characteristics measured by cardiometabolic risk (range 1–46%) are presented in Table I.

Multivariable Associations

Significant variables from baseline analyses were included in a multi-linear model in order to control for covariation. Table II reports the unadjusted regression coefficient (B). The adjusted regression coefficient (β) reported in the following paragraph, represents the relation between X¹ and Y averaged over all values of X², hence showing the most influential predictor of the significant covariates. Sleep maintenance problems at baseline significantly predicted cardiometabolic risk score at follow-up and this association remained after adjusting for covariates ($\beta = 0.26$). In addition, whether or not the patient care worker reported doing the recommended level of PA at baseline, predicted increased cardiometabolic risk at follow-up in the adjusted model ($\beta = 0.29$), as did the occupational category the worker belonged to ($\beta = 0.24$). The overall model fit was ($X^2_{\text{red}} = 0.23$) and a Durbin-Watson test yielded $d = 1.59$, which is within the critical value standards for non-autocorrelation, meaning the unobserved error terms are not correlated (see Table II for details of model results).

DISCUSSION

In this longitudinal study of 99 female patient care workers, we investigated how baseline measures of sleep deficiency, PA, work stress and night work predicted 10-year cardiometabolic risk from biomarkers at 2-year follow-up. Sleep maintenance problems, occupational category, and not meeting the CDC’s recommended amount of PA at baseline, significantly predicted increased 10-year cardiometabolic risk. These predictive associations remained when controlled for covariates.

TABLE I. Mean (SD), Correlation (r), Variance (r^2/F) of Participant Characteristics on 10-Year Cardiometabolic Risk Score Within Covariates

Work conditions/demographics	Cardiometabolic risk score (range 0–100 %)		P-value
	Independent variables		
Sleep deficiency	Yes (n = 63) 9.29 % (10.5 %)	No (n = 36) 6.03 % (6.2 %)	0.058
Sleep duration	<6 hr (n = 26) 9.5 % (9.0 %)	≥6 hr (n = 73) 7.54 % (9.3 %)	0.36
Insufficient sleep	Yes (n = 29) 6.13 % (8.8 %)	No (n = 69) 8.86 % (9.3 %)	1.000
Sleep maintenance problems	Yes (n = 39) 10.51 % (10.6 %)	No (n = 59) 6.5 % (7.8 %)	0.048*
Recommended physical activity	Yes (n = 61) 11.9 % (11.7 %)	No (n = 38) 5.6 % (6.0 %)	0.003*
Iso-strain	Yes (n = 14) 8.82 % (9.2 %)	No (n = 74) 8.36 % (9.7 %)	0.87
Job demands (12–48)	$r = 0.09 r^2 = 0.008$		0.37
Decision latitude (2–10)	$r = 0.36 r^2 = 0.001$		0.73
Coworker support (2–10)	$r = 0.14 r^2 = 0.02$		0.17
Supervisor support (3–15)	$r = 0.09 r^2 = 0.009$		0.36
Work-family conflict (5–25)	$r = 0.04 r^2 = 0.002$		0.68
Occupation (staff nurse ; patient care associate (PCA); other)	(Welch's F) 28.2 (2, 50.2)		<0.001*
Staff nurse	6.5 % (7 %)		
PCA	1.2 % (0.6 %)		
Other	13 % (12.3 %)		
Race/ethnicity (white ; hispanic; black)	$F(2, 92) 0.14$		0.87
White	8.2 % (9.2 %)		
Hispanic	5.4 % (5.1 %)		
Black	8.1 % (12.9 %)		
Education (GED or less ; some college; degree; graduate school)	(Welch's F) 1.3 (3, 4.6)		0.38
GED or less	17.4 % (20.5 %)		
Some college	7.8 % (9 %)		
Degree	6.6 % (8.1 %)		
Graduate school	13 % (11.0 %)		
Economic status (difficulty paying bills?) (little difficulty w/bills ; at least some difficulty w/bills)	$F(4, 91) 0.6$		0.7
A great deal of difficulty w/bills	5.2 % (3.6 %)		
At least some difficulty w/bills	8.4 % (8.8 %)		
A little difficulty	10 % (10.2 %)		
No difficulty	7.6 % (9.3 %)		
Refused to answer	2.9 % (2.9 %)		
Night work (0–6 hr monthly ; >6 hr but <72; 72+)	(Welch's F) 6.1 (2, 25.6)		0.007*
0–6 hr monthly	8.8 % (9.6 %)		
>6 hr but <72	3.9 % (4.8 %)		
72+	10.6 % (9.7 %)		

Covariates are either dichotomous (yes/no) or ordinal/continuous (range). Categorical variables are shown with both test statistic and mean (SD). Reference categories are bolded. All characteristics may not add up to n = 99, because of missing data.

*Significant P-values (<0.05) are bolded.

Our finding that sleep maintenance problems independently predict increased cardiometabolic 10-year risk is consistent with a recent large-scale prospective study on reduced sleep quality and incidence of heart failure which

identified a dose-response relationship between cumulative increase in insomnia symptoms and risk of heart failure [Laugsand et al., 2013]. Earlier studies on insomnia symptoms and cardiovascular risk further support a link to

TABLE II. Multivariable Linear Regression Showing Associations of Baseline Characteristics on 10-Year Cardiometabolic Risk at Follow-up

Independent variables	Cardiometabolic risk model	
	Unstandardized coefficient (95 % CI)	P-value
Sleep maintenance problems (yes; no)	0.046 (0.01, 0.048)	0.01
Occupation (staff nurse; PCA; other)	0.027 (0.006, 0.48)	0.01
Recommended physical activity (yes; no)	0.054 (0.019, 0.09)	0.003
Night work categories (≤6, 6–72, 72+)	0.015 (−0.012, 0.041)	0.27

Categories are listed in parentheses.

Significant P-values are bolded. The interpretation of the coefficient is the expected change in *y* for a one-unit change in *x* when the other covariates are held fixed—that is, the expected value of the partial derivative in *y* for a one-unit change in *x* when other covariates are held fixed. This is sometimes called the unique effect of *x* on *y*.

heart failure as well as cardiovascular disease [Newman et al., 2000; Chien et al., 2010]. A recent study of insomnia and heart failure [Laugsand et al., 2013] describes an increased relative risk of heart failure for women compared to men, which was attributed to either a higher prevalence of insomnia symptoms in females [Ohayon, 2002] or the lower baseline risk for heart failure in women [Laugsand et al., 2013]. Our results support further investigation into whether insomnia symptoms pose a greater threat to women than men when assessing cardiometabolic risk.

Short sleep duration has previously been linked to increased blood pressure [Gottlieb et al., 2006; Buxton and Marcelli, 2010]. Recent findings from a large American cohort also demonstrated that insomnia symptoms led to a fivefold elevated risk for hypertension when combined with short sleep duration (≤5 hr), whereas insomnia symptoms and adequate sleep duration did not show significant associations with hypertension [Vgontzas et al., 2009]. Our results support a predictive link between sleep maintenance problems and cardiometabolic risk, but we did not find evidence for such a “combination effect” when investigating sleep deficiency. However, this could be due to our small sample or all female population; future studies should include combination/composite variables to further the understanding of this link.

Our results failed to show a link between night/shift work hours per month and 10-year cardiometabolic risk, when controlling for covariates. Rotating night shifts has previously been associated with increased risk for ischemic heart disease [Brown et al., 2009] and exhibits a dose-response relationship with coronary heart disease in the Nurses Health Study [Kawachi et al., 1995; Brown et al., 2009]. However,

there are other studies showing no association between night shifts and ischemic heart disease [Frost et al., 2009]. A recent review on night work and ischemic heart disease looked at 16 studies with a relative risk varying from 0.64 to 2.25 [Frost et al., 2009]. However, overlap between sleep maintenance problems and working night shifts could perhaps be expected as sleep deficiency has been suggested as a mechanism explaining earlier findings [Brown et al., 2009]. In the current study, there was no significant correlation between working night shifts and reporting sleep maintenance problems. Our results are in line with a recent review showing only a marginally increased relative risk when obtaining night work hours from company records, compared to when night work is self-reported [Frost et al., 2009].

Failing to exercise moderately or vigorously for 150 min or more during a week is a well-established risk factor in cardiovascular disease [Sattelmair et al., 2011; Wen et al., 2011]. The results from our study are in accordance with numerous prospective cohort studies and meta-analyses done in several countries [Sattelmair et al., 2011; Wen et al., 2011]. Our occupational group, patient care workers, have been investigated through the Nurses Health Study, demonstrating that even light PA was associated with lower cardiovascular risk [Lee et al., 2001].

Interestingly, several studies have documented a link between short sleep duration and reduced PA [Briones et al., 1996; Weaver et al., 1997]. A recent laboratory study demonstrated how induced short sleep duration (<5.5) in 18 healthy subjects led to a 31% decrease in daily activity and 24% reduction in moderate and vigorous activity [Bromley et al., 2012]. When looking at this in our population, though not significant, we observed a trend that a lower percentage of the participants’ reporting short sleep duration met the recommended amounts of PA than participants reporting adequate sleep duration. Such a link between sleep and PA underlines the need for a specific focus on sleep adequacy, duration, and maintenance in occupational health interventions.

No measures of job stress were significant in the adjusted analyses in our study. A review on psychosocial stress and ischemic heart disease risk highlights the contradictory evidence in female populations when it comes to job stress [Eller et al., 2009].

This study has limitations, especially that the sample size is small and was not chosen randomly, which limits the generalizability of our results. However, an all female population is not common in cardiometabolic risk studies. These results should be considered a contribution to a growing research base on female cardiometabolic risk. Another limitation is that we have only two time points approximately 2 years apart that does not allow for investigations of mediating mechanisms or directionality. But, the extensive questionnaires at baseline, administrative payroll data on night work status, and lengthy follow-up with biomarkers still make a strong case for the relevance of

variables showing significant predictive value. Another limitation of note is that our prospective design does not preclude reverse causality. We do not have information on events before the beginning of the study and cannot account for other factors influencing causality. Moreover, we do not have a measure or question addressing postmenopausal status. The average menopausal age in American women is 51 years, and the range is 40–61 years, and about half of our sample is within this range, which is an important risk factor.

The relationship between sleep maintenance problems, PA and cardiometabolic risk is a widely studied connection. Yet, there is less evidence in female patient care workers, a population with very high stress levels and high prevalence of sleep deficiency. The Nurses Health Study is perhaps the largest cohort with the most extensive follow-up done on a nurse population. However, its participants consist only of married nurses and do not include PCA or single nurses, who are vulnerable workers with regards to both social support and workload. This study also implements biomarker data in a high priority population providing objective measurements to motivate change in occupational health practices. In addition, our data includes non self-report variables such as administrative payroll data on night work status, and measured cardiometabolic risk [Marino et al., 2014], a much more sensitive outcome than cardiovascular disease or death as registered by death certificates, self-report or hospitalization due to modest accuracy of diagnoses, and referral biases [Frost et al., 2009]. Sleep has previously been suggested as a third pillar of health, alongside nutrition and exercise, in cardiometabolic disease risk [Buxton et al., 2013]. Our results support this view and highlight the need for sleep deficiency to be prioritized in occupational health programs, especially those targeting cardiovascular health in women.

ACKNOWLEDGMENTS

This work was supported by a grant (Grant sponsor: National Institute for Occupational Safety and Health. Grant number: U19 OH008861) for the Harvard School of Public Health Center for Work, Health and Well-being, for the Harvard Clinical and Translational Science Center, (Grant sponsor: National Center for Research Resources. Grant number: UL1 RR025758-04), the Work, Family, and Health Network (Grant sponsor: Grant number: U01 AG5186989), (Grant sponsor: National Heart, Lung, and Blood Institute. Grant number: R01HL107240), and additional support was provided by Dean Hashimoto and Partners Occupational Health Service. This study would not have been accomplished without the participation of Partners HealthCare System and leadership from Dennis Colling, Sree Chaguturu, and Kurt Westerman. The authors would like to thank Partners Occupational Health Services

including Marlene Freeley for her guidance, as well as Elizabeth Taylor, Elizabeth Tucker O'Day, and Terry Orechia. We also thank individuals at each of the hospitals including Jeanette Ives Erickson and Jacqueline Somerville in Patient Care Services leadership, and Jeff Davis and Julie Celano in Human Resources. Additionally, we wish to thank Charlene Feilteau, Mimi O'Connor, Margaret Shaw, Eddie Tan, and Shari Weingarten for assistance with supporting databases. We also thank Chris Kenwood of NERI for his statistical and programming support. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Center For Research Resources or the National Institutes of Health.

REFERENCES

Ayas NT, White DP, Al-Delaimy WK, Manson JAE, Stampfer MJ, Speizer FE, Patel S, Hu FB. 2003a. A prospective study of self-reported sleep duration and incident diabetes in women. *Diabetes Care* 26:380–384.

Ayas NT, White DP, Manson JE, Stampfer MJ, Speizer FE, Malhotra A, Hu FB. 2003b. A prospective study of sleep duration and coronary heart disease in women. *Arch Intern Med* 163:205–209.

Backe EM, Seidler A, Latza U, Rossnagel K, Schumann B. 2012. The role of psychosocial stress at work for the development of cardiovascular diseases: A systematic review. *Int Arch Occup Environ Health* 85:67–79.

Belkic KL, Landsbergis PA, Schnall PL, Baker D. 2004. Is job strain a major source of cardiovascular disease risk? *Scand J Work Environ Health* 30:85–128.

Bellavia GM, Frone MR. 2005. Work-family conflict. In: Barling J, Kelloway EK, Frone MR, editors. *Handbook of work stress*. Thousand Oaks, CA: Sage. pp. 113–147.

Berkman LF, Buxton O, Ertel K, Okechukwu C. 2010. Managers' practices related to work–family balance predict employee cardiovascular risk and sleep duration in extended care settings. *J Occup Health Psychol* 15:316–329.

Bhui KS, Dinos S, Stansfeld SA, White PD. 2012. A synthesis of the evidence for managing stress at work: A review of the reviews reporting on anxiety, depression, and absenteeism. *J Environ Public Health* 2012, Article ID 515874, 21 pages. doi: 10.1155/2012/515874

Briones B, Adams N, Strauss M, Rosenberg C. 1996. Relationship between sleepiness and general health status. *Sleep* 19:583–588.

Bromley LE, Booth JN, III, Kilkus JM, Imperial JG, Penev PD. 2012. Sleep restriction decreases the physical activity of adults at risk for type 2 diabetes. *Sleep* 35:977–984.

Brown DL, Feskanich D, Sánchez BN, Rexrode KM, Schernhammer ES, Lisabeth LD. 2009. Rotating night shift work and the risk of ischemic stroke. *Am J Epidemiol* 169:1370–1377.

Buxton OM, Broussard JL, Zahl AK, Hall M. 2013. Effects of sleep deficiency on hormones, cytokines, and metabolism. In: Redline S, Berger NA, editors. *Energy balance and cancer volume 8: Impact of sleep and sleep disturbances on obesity and cancer*. New York: Springer.

Buxton OM, Hopcia K, Sembajwe G, Porter JH, Dennerlein JT, Kenwood C, Stoddard AM, Hashimoto D, Sorensen G. 2012. Relationship of sleep deficiency to perceived pain and functional limitations in hospital patient care workers. *J Occup Environ Med* 54:851–858.

Buxton OM, Marcelli E. 2010. Short and long sleep are positively associated with obesity, diabetes, hypertension, and cardiovascular disease among adults in the United States. *Soc Sci Med* 71:1027–1036.

Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. 1989. The Pittsburgh sleep quality index: A new instrument for psychiatric practice and research. *Psychiatry Res* 28:193–213.

Cappuccio FP, D'Elia L, Strazzullo P, Miller MA. 2010. Sleep duration and all-cause mortality: A systematic review and meta-analysis of prospective studies. *Sleep* 33:585–592.

Centers for Disease Control and Prevention. 2009. Behavioral risk factor surveillance system survey questionnaire. Atlanta, GA: Centers for Disease Control and Prevention.

Chandola T, Ferrie JE, Perski A, Akbaraly T, Marmot MG. 2010. The effect of short sleep duration on coronary heart disease risk is greatest among those with sleep disturbance: A prospective study from the Whitehall II cohort. *Sleep* 33:739–744.

Chien K-L, Chen P-C, Hsu H-C, Su T-C, Sung F-C, Chen M-F, Lee Y-T. 2010. Habitual sleep duration and insomnia and the risk of cardiovascular events and all-cause death: Report from a community-based cohort. *Sleep* 33:177–184.

Cizza G, Skarulis M, Mignot E. 2005. A link between short sleep and obesity: Building the evidence for causation. *Sleep* 28:1217–1220.

D'Agostino RB, Vasan RS, Pencina MJ, Wolf PA, Cobain M, Massaro JM, Kannel WB. 2008. General cardiovascular risk profile for use in primary care the Framingham Heart Study. *Circulation* 117:743–753.

Dew MA, Hoch CC, Buysse DJ, Monk TH, Begley AE, Houck PR, Hall M, Kupfer DJ, Reynolds CF, III. 2003. Healthy older adults' sleep predicts all-cause mortality at 4 to 19 years of follow-up. *Psychosom Med* 65:63–73.

Eller NH, Netterstrom B, Gyntelberg F, Kristensen TS, Nielsen F, Steptoe A, Theorell T. 2009. Work-related psychosocial factors and the development of ischemic heart disease: A systematic review. *Cardiol Rev* 17:83–97.

Evans O, Steptoe A. 2002. The contribution of gender-role orientation, work factors and home stressors to psychological well-being and sickness absence in male-and female-dominated occupational groups. *Soc Sci Med* 54:481–492.

Frone MR, Russell M, Cooper ML. 2011. Relation of work-family conflict to health outcomes: A four-year longitudinal study of employed parents. *J Occup Org Psychol* 70:325–335.

Frost P, Kolstad HA, Bonde JP. 2009. Shift work and the risk of ischemic heart disease—a systematic review of the epidemiologic evidence. *Scand J Work Environ Health* 35:163–179.

Gallicchio L, Kalesan B. 2009. Sleep duration and mortality: A systematic review and meta-analysis. *J Sleep Res* 18:148–158.

Gangwisch JE, Malaspina D, Boden-Albala B, Heymsfield SB. 2005. Inadequate sleep as a risk factor for obesity: Analyses of the NHANES I. *Sleep* 28:1289–1296.

Gottlieb DJ, Punjabi NM, Newman AB, Resnick HE, Redline S, Baldwin CM, Nieto FJ. 2005. Association of sleep time with diabetes mellitus and impaired glucose tolerance. *Arch Intern Med* 165:863–867.

Gottlieb DJ, Redline S, Nieto FJ, Baldwin CM, Newman AB, Resnick HE, Punjabi NM. 2006. Association of usual sleep duration with hypertension: The Sleep Heart Health Study. *Sleep* 29:1009–1014.

Grandner MA, Hale L, Moore M, Patel NP. 2010. Mortality associated with short sleep duration: The evidence, the possible mechanisms, and the future. *Sleep Med Rev* 14:191–203.

Hale L, Do DP. 2007. Racial differences in self-reports of sleep duration in a population-based study. *Sleep* 30:1096–1103.

Harvey AG, Murray G, Chandler RA, Soehner A. 2011. Sleep disturbance as transdiagnostic: Consideration of neurobiological mechanisms. *Clin Psychol Rev* 31:225–235.

Hasler G, Buysse DJ, Klaghofer R, Gamma A, Ajdacic V, Eich D, Rössler W, Angst J. 2004. The association between short sleep duration and obesity in young adults: A 13-year prospective study. *Sleep* 27:661–666.

Hublin C, Partinen M, Koskenvuo K, Silventoinen K, Koskenvuo M, Kaprio J. 2010. Shift-work and cardiovascular disease: A population-based 22-year follow-up study. *Eur J epidemiol* 25:315–323.

Karasek RA, Jr. 1979. Job demands, job decision latitude, and mental strain: Implications for job redesign. *Admin Sci Q* 24:285–308.

Karasek R, Brisson C, Kawakami N, Houtman I, Bongers P, Amick B. 1998. The Job Content Questionnaire (JCQ): An instrument for internationally comparative assessments of psychosocial job characteristics. *J Occup Health Psychol* 3:322–355.

Kawachi I, Colditz GA, Stampfer MJ, Willett WC, Manson JAE, Speizer FE, Hennekens CH. 1995. Prospective study of shift work and risk of coronary heart disease in women. *Circulation* 92:3178–3182.

Kim SS, Okechukwu CA, Buxton OM, Dennerlein JT, Boden LI, Hashimoto DM, Sorensen G. 2012. Association between work-family conflict and musculoskeletal pain among hospital patient care workers. *Am J Ind Med* 56:488–495.

Kivimaki M, Nyberg ST, Batty GD, Fransson EI, Heikkila K, Alfredsson L, Björner JB, Borritz M, Burr H, Casini A, et al. 2012. Job strain as a risk factor for coronary heart disease: A collaborative meta-analysis of individual participant data. *Lancet* 380:1491–1497.

Kohatsu ND, Tsai R, Young T, VanGilder R, Burmeister LF, Stromquist AM, Merchant JA. 2006. Sleep duration and body mass index in a rural population. *Arch Intern Med* 166:1701–1705.

Kripke DF, Garfinkel L, Wingard DL, Klauber MR, Marler MR. 2002. Mortality associated with sleep duration and insomnia. *Arch Gen Psychiatry* 59:131–136.

Laugsand LE, Strand LB, Platou C, Vatten LJ, Janszky I. 2013. Insomnia and the risk of incident heart failure: A population study. *Eur Heart J* [Epub ahead of print].

Lee IM, Rexrode KM, Cook NR, Manson JAE, Buring JE. 2001. Physical activity and coronary heart disease in women. *JAMA* 285:1447–1454.

Lusardi P, Zoppi A, Preti P, Pesce RM, Piazza E, Fogari R. 1999. Effects of insufficient sleep on blood pressure in hypertensive patients: A 24-h study. *Am J Hypertens* 12:63–68.

Luyster FS, Strollo PJ, Jr, Zee PC, Walsh JK. 2012. Sleep: A health imperative. *Sleep* 35:727–734.

Marino M, Li Y, Pencina MJ, D'Agostino RB, Berkman L, Buxton OM. 2014. Quantifying cardiometabolic risk using modifiable non-self-reported risk factors. *Am J Prev Med* In press.

Mathers C, Fat DM, Boerma J. 2008. The global burden of disease: 2004 update. Geneva, Switzerland: World Health Organization.

McKnight-Eily L, Presley-Cantrell L, Strine T, Chapman D, Perry G, Croft J. 2008. Perceived insufficient rest or sleep-four states, 2006. *MMWR Morb Mortal Wkly Rep* 57:200–203.

Meier-Ewert HK, Ridker PM, Mullington JM, Dinges DF, Price NJ, Regan MM, Rifai N. 2004. Effect of sleep loss on C-reactive protein, an inflammatory marker of cardiovascular risk. *J Am Coll Cardiol* 43:678–683.

Netemeyer RG, Boles JS, McMurrian R. 1996. Development and validation of work-family conflict and family-work conflict scales. *J Appl Psychol* 81:400–410.

Newman AB, Spiekerman CF, Enright P, Lefkowitz D, Manolio T, Reynolds CF, Robbins J. 2000. Daytime sleepiness predicts mortality and cardiovascular disease in older adults. *J Am Geriatr Soc* 48:115–123.

Ohayon MM. 2002. Epidemiology of insomnia: What we know and what we still need to learn. *Sleep Med Rev* 6:97–111.

Patel SR, Ayas NT, Malhotra MR, White DP, Schernhammer ES, Speizer FE, Stampfer MJ, Hu FB. 2004. A prospective study of sleep duration and mortality risk in women. *Sleep* 27:440–444.

Patel SR, Blackwell T, Redline S, Ancoli-Israel S, Cauley JA, Hillier TA, Lewis CE, Orwoll ES, Stefanick ML, Taylor BC. 2008. The association between sleep duration and obesity in older adults. *Int J Obes* 32:1825–1834.

Roger VL, Go AS, Lloyd-Jones DM, Adams RJ, Berry JD, Brown TM, Carnethon MR, Dai S, de Simone G, Ford ES. 2011. Heart disease and stroke statistics-2011 Update. *Circulation* 123:18–209.

Sattelmair J, Pertman J, Ding EL, Kohl HW, III. Haskell W, Lee IM. 2011. Dose response between physical activity and risk of coronary heart disease. *Circulation* 124:789–795.

Schnall PL, Dobson M, Rosskam E. 2009. *Unhealthy work: Causes, consequences, cures*. New York: Baywood Publishing Company.

Sorensen G, Stoddard AM, Stoffel S, Buxton O, Sembajwe G, Hashimoto D, Dennerlein JT, Hopcia K. 2011. The role of the work context in multiple wellness outcomes for hospital patient care workers. *J Occup Environ Med* 53:899–910.

Steptoe A, Kivimäki M. 2012. Stress and cardiovascular disease. *Nat Rev Cardiol* 9:360–370.

Stranges S, Dorn JM, Cappuccio FP, Donahue RP, Rafalson LB, Hovey KM, Freudenheim JL, Kandala N-B, Miller MA, Trevisan M. 2010. A population-based study of reduced sleep duration and hypertension: The strongest association may be in premenopausal women. *J Hypertens* 28:896–902.

Taheri S, Lin L, Austin D, Young T, Mignot E. 2004. Short sleep duration is associated with reduced leptin, elevated ghrelin, and increased body mass index. *PLoS Med* 1:e62.

US Department of Health and Human Services. *Focus area 7: Educational and community-based programs—worksite setting*. Healthy People 2010.

Vgontzas AN, Liao D, Bixler EO, Chrousos GP, Vela-Bueno A. 2009. Insomnia with objective short sleep duration is associated with a high risk for hypertension. *Sleep* 32:491–497.

Weaver TE, Laizner AM, Evans LK, Maislin G, Chugh DK, Lyon K, Smith PL, Schwartz AR, Redline S, Pack AI. 1997. An instrument to measure functional status outcomes for disorders of excessive sleepiness. *Sleep* 20:835–843.

Wen CP, Wai JPM, Tsai MK, Yang YC, Cheng TYD, Lee MC, Chan HT, Tsao CK, Tsai SP, Wu X. 2011. Minimum amount of physical activity for reduced mortality and extended life expectancy: A prospective cohort study. *Lancet* 378:1244–1253.

Wilson PWF, D'Agostino RB, Levy D, Belanger AM, Silbershatz H, Kannel WB. 1998. Prediction of coronary heart disease using risk factor categories. *Circulation* 97:1837–1847.

Wingard DL, Berkman LF. 1983. Mortality risk associated with sleeping patterns among adults. *Sleep* 6:102–107.

Yang H, Schnall PL, Jauregui M, Su TC, Baker D. 2006. Work hours and self-reported hypertension among working people in California. *Hypertension* 48:744–750.

Åkerstedt T. 2003. Shift work and disturbed sleep/wakefulness. *Occup Med* 53:89–94.