

Poster presentation

0035 CANCER RISK IN OIL REFINERY WORKERS: A MORTALITY STUDY IN FOUR ITALIAN PLANTS

^{1,2}Angela Cecilia Pesatori, ²Dario Consonni, ²Paolo Grillo, ^{1,2}Pier Alberto Bertazzi. ¹Department of Clinical Sciences and Community Health, University of Milan, Milan, Italy; ²Epidemiology Unit, Department of Preventive Medicine, IRCCS Ca' Granda Foundation, Milan, Italy

10.1136/oemed-2014-102362.195

Objectives To examine the mortality experience of workers employed in four Italian oil refineries.

Method The cohort included 5112 male workers ever employed between 1949 and 2011. The average follow-up period was 49 years. SMR and 95% CI were calculated using as reference age-gender-calendar specific regional rates. Analyses by duration of employment and latency were performed.

Results In the whole cohort, pleural (6 deaths, SMR 1.59; 95% CI 0.7–3.5), brain cancers (14 deaths, SMR 1.47; 95% CI 0.9–2.5) and lymphatic leukaemia (LL) (8 deaths, SMR 1.81; 95% CI 0.9–3.6) showed increased risks. All pleural cancers occurred after 10 years of latency and the highest risk was observed among workers with duration \geq 20 years; the brain cancer excess was confined in the shortest duration and latency. The LL excess regarded workers with latency and duration longer than 10 years. Mortality from Non-Hodgkin lymphoma (NHL) (13 deaths) and acute myeloid leukaemia (AML) did not differ from the expectation. All AML cases (4 deaths) occurred after 20 years of latency (SMR 1.55, 95% CI 0.6–4.1) and a two-fold-increased risk was observed in the longest duration. Mortality for NHL + LL (recently classified as subtypes of NHL) showed increased risks among workers with duration and latency longer than 20 years.

Conclusions Our findings confirmed recent epidemiological evidences of an increased risk for pleural cancer and are coherent with most recent meta-analyses suggesting a limited evidence of an increased risk for lymphatic neoplasms in refinery workers possibly due to past exposure to benzene.

0038 CHILDREN BELOW 5 YEARS OF EMPLOYED MOTHERS ARE LESS EXPOSED TO ACUTE POISONING IN ALEXANDRIA, EGYPT

Raed Alazab. Alazhar University, College of Medicine, Cairo, Egypt

10.1136/oemed-2014-102362.196

Objectives To identify the incidence rate and determinants of acute poisoning among children (1–60 months old) of employed mothers.

Method A study was conducted at the poisoning unit of a university hospital. The studied children were from both rural and urban areas, were a mix of boys and girls, did not suffer from any mental disabilities, were aged between 1 month old to 60 months old, and were of Egyptian nationality. Data was collected by using a clinical examination form and a questionnaire. All parents/carers of the studied children were interviewed as well. Clinical assessment of the children included: general health conditions; AVPU (alert, respond to verbal stimuli, respond to painful stimuli, unconsciousness); and clinical examinations.

Results 18.5% of total admissions were children (1–60 months old), 62.5% were males, 83.3% did not attend nursery, 79.9% were from urban areas, 33% of mothers were illiterate, and 60.2% of poisonings were due to household products. Kerosene alone was implicated in 24.3% of all cases; 37.4% of cases took

place in the kitchen; 47.4% of cases were poisoned during the period between 8am and 4pm, and 65.4% reached the poisoning unit within 2 to 4 h of accidental poisoning. Risk factors among the studied children were ordered by stepwise regression analysis as the following: non employed mothers; children who did not attend nursery; children of the male gender; and the education and literacy level of their mothers.

Conclusions Children of non employed mothers are at more risk for acute poisoning.

0041 ASSOCIATIONS OF THERMOPHILIC ACTINOMYCETES AND NON-TUBERCULOUS MYCOBACTERIA WITH RESPIRATORY HEALTH IN OCCUPANTS OF A WATER-DAMAGED OFFICE BUILDING

¹Ju-Hyeong Park, ¹Jean Cox-Ganser, ¹Sandra White, ¹Scott Laney, ²Steve Caulfield, ²William Turner, ³Austin Sumner, ¹Kathleen Kreiss. ¹National Institute for Occupational Safety and Health, Morgantown, WV, USA; ²Turner Building Science and Design, LLC, Harrison, ME, USA; ³Northwestern Occupational Health, Albans, VT, USA

10.1136/oemed-2014-102362.197

Objectives We examined microbial correlates of health outcomes in employees in a building with a sarcoidosis cluster and excess asthma.

Method We offered employees a questionnaire and pulmonary function tests (PFT) and collected 120 floor dust samples by vacuuming workstation areas or the exterior rim of the floor. We analysed dust for culturable fungi, bacteria, actinomycetes, non-tuberculous mycobacteria, and cell wall components of fungi [(1 \rightarrow 3)- β -D-glucan] and Gram-negative bacteria (endotoxin).

Results Among 136 employees, 77% participated in the questionnaire, 64% in PFTs, and 49% in both. Of the 105 questionnaire participants, 50 (48%) reported at least one hypersensitivity pneumonitis (HP)-like symptom (shortness of breath on exertion, flu-like achiness, or fever and chills) in the last 4 weeks; 69 (66%) reported at least one HP-like symptom in the last 12 months. PFT results were abnormal or borderline for 13 participants (6 obstruction; 2 restriction; 2 mixed pattern; and 3 borderline obstruction). In multivariate logistic regressions, log-transformed levels of thermophilic actinomycetes (GM=580 CFU/m²) were positively associated with HP-like symptoms in the last 4 weeks [odds ratio (OR)=1.6, 95% confidence interval (CI)=1.15–2.35] and 12 months (OR=1.6, 95% CI=1.09–2.32), and with an abnormal/borderline PFT result (OR=1.7, 95% CI=1.04–2.88). Log-transformed mycobacteria levels (GM=960 CFU/m²) were marginally positively associated with HP-like symptoms in the last 4 weeks that improved when away from the building (OR=1.9; 95% CI=0.95–3.85). These associations were not confounded by smoking, gender, age, or building tenure.

Conclusions Our study suggests that thermophilic actinomycetes and non-tuberculous mycobacteria may have played a role in the respiratory illness among occupants of this water-damaged building.

0043 BENIGN PLEURAL ABNORMALITIES AND MESOTHELIOMA AMONG ASBESTOSIS WORKERS IN HONG KONG

¹Lap Ah (Shelly) Tse, ²Ronald KF Au, ¹Minghui Chen, ¹Feng Wang, ¹Ignatius TS Yu. ¹JC School of Public Health and Primary Care, The Chinese University of Hong Kong, Hong Kong, China; ²Pneumoconiosis Clinic, Department of Health, Hong Kong, China

10.1136/oemed-2014-102362.198



0041 Associations of thermophilic actinomycetes and non-tuberculous mycobacteria with respiratory health in occupants of a water-damaged office building

Ju-Hyeong Park, Jean Cox-Ganser, Sandra White, et al.

Occup Environ Med 2014 71: A63
doi: 10.1136/oemed-2014-102362.197

Updated information and services can be found at:
http://oem.bmjjournals.org/content/71/Suppl_1/A63.3

These include:

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:
<http://group.bmjjournals.org/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmjjournals.org/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmjjournals.org/subscribe/>