



Evidence of Health Risks Associated with Prolonged Standing at Work and Intervention Effectiveness

Thomas R. Waters¹, PhD & Robert B. Dick¹, PhD

¹ National Institute For Occupational Safety and Health, Cincinnati, OH, USA

Keywords

Prolonged standing; low back pain; leg pain; cardiovascular problems; subjective measures; biomechanical/physiological measures; interventions (floor mats, shoe inserts, sit-stand chairs/workstations); guidelines.

Correspondence

Robert B. Dick, PhD, Guest Researcher, National Institute for Occupational Safety and Health, 4676 Columbia Parkway, MSC24, Cincinnati, OH 45226-1998.
E-mail: rbd1@cdc.gov

The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the National Institute for Occupational Safety and Health.

Accepted April 06, 2014.

doi: 10.1002/rnj.166

Abstract

Purpose: Prolonged standing at work has been shown to be associated with a number of potentially serious health outcomes, such as lower back and leg pain, cardiovascular problems, fatigue, discomfort, and pregnancy-related health outcomes. Recent studies have been conducted examining the relationship between these health outcomes and the amount of time spent standing while on the job. The purpose of this article was to provide a review of the health risks and interventions for workers and employers that are involved in occupations requiring prolonged standing. A brief review of recommendations by governmental and professional organizations for hours of prolonged standing is also included.

Findings: Based on our review of the literature, there seems to be ample evidence showing that prolonged standing at work leads to adverse health outcomes. Review of the literature also supports the conclusion that certain interventions are effective in reducing the hazards associated with prolonged standing. Suggested interventions include the use of floor mats, sit-stand workstations/chairs, shoes, shoe inserts and hosiery or stockings. Studies could be improved by using more precise definitions of prolonged standing (e.g., duration, movement restrictions, and type of work), better measurement of the health outcomes, and more rigorous study protocols.

Conclusion and Clinical Relevance: Use of interventions and following suggested guidelines on hours of standing from governmental and professional organizations should reduce the health risks from prolonged standing.

Introduction

Many workers are required to stand for long periods of time without being able to walk or sit during the work shift. In operating rooms, for example, nurses and doctors must stand for many hours during surgical procedures. Similarly, direct care nurses, hairdressers, and store clerks spend large fractions of their working time standing without the ability to sit down. Briefly, a short summary below outlines the scope of the prolonged standing situation in working populations.

McCulloch (2002) summarized findings from 17 studies that involved standing for more than 8 hours per day. Major health risks identified were chronic venous insufficiency, musculoskeletal pain of the lower back and feet, preterm birth, and spontaneous abortions. Best et al. (2002) reported on the findings from a self-reported questionnaire administered to 204 hairdressers. Back pain was the most reported musculoskeletal disorder followed by neck and shoulder discomfort. Duration of standing was reported to be between 82% and 99% of total work time. Tissot, Messing, and Stock (2005) reported that the

standing at work prevalence rate is 58% in the Quebec working population and more common in men, workers >25 years of age, and lower income workers. Meijssen and Knibbe (2007) reported that the average standing time for Dutch peri-operative personnel [equivalent to nurses in the United States] was 2.5 hours per day, and that 18% of respondents exceeded 4 hours of standing per day and 47% were in the Amber zone, and 17% were classified into the Red zone. According to Dutch ergonomic guidelines for prolonged standing, exposure is classified into one of three zones: Green (safe—continuous standing ≤ 1 hour and total/day ≤ 4 hours), Amber (action recommended—continuous standing > 1 hour *or* total/day > 4 hours), or Red (direct action required—continuous standing > 1 hour *and* total/day > 4 hours). Werner, Gell, Hartigan, Wiggermann, and Keyserling (2010) in a cross-sectional evaluation of workers at an engine manufacturing plant in jobs that necessitated prolonged standing and walking reported that 24% met the case definition for foot/ankle disorder and that 52% had the symptoms. Sitting, standing, and type of work surface did not change the prevalence.

To emphasize the importance of this topic and concern regarding worker safety, the Association for peri-operative Registered Nurses (AORN) recently published guidelines and solutions for reducing health risks associated with prolonged standing in peri-operative environments (Hughes, Nelson, Matz, & Lloyd, 2011). In the guideline adopted by AORN, it is recommended that caregivers should not stand more than 2 hours continuously or for more than 30% of the work day without some type of fatigue-reducing interventions, such as antifatigue mats, specially designed footstools, sit-stand stools or chairs, or supportive footwear. The AORN guideline also suggests that if the caregiver must wear a lead apron during prolonged standing, that exposure should be limited to 1 hour without some type of intervention.

Additionally, the Canadian Centre for Occupational Health and Safety (CCOHS) (2014) has reported that working in a standing posture on a regular basis can cause sore feet, swelling of the legs, varicose veins, general muscular fatigue, and low back pain, stiffness in the neck and shoulders, and other health problems. According to the CCOHS report, prolonged standing effectively reduces the blood supply to the muscles resulting in the acceleration of the onset of fatigue and causes pain in the muscles of the legs, back and neck, as well as pooling of blood in the legs and feet which leads to varicose veins. The CCOHS suggests that job design can reduce the ill effects

of working in a standing position by changing working positions frequently, avoiding extreme bending, stretching, and twisting, pacing work appropriately, and allowing workers suitable rest periods. The CCOHS report also suggests use of floor mats, shoe inserts, compression hosiery, and ergonomic seating to avoid exposure.

The International Labour Organization (ILO) (2011) has also published guidelines for prevention of health effects associated with exposure to prolonged standing at work. According to the ILO, if a job must be done in a standing position, a chair or stool should be provided for the worker and he or she should be able to sit down at regular intervals. The ILO also suggests use of floor mats and good shoes to avoid standing on a hard surface, as well as the availability of footrests to help reduce the strain on the back and to allow the worker to change positions by shifting weight from time to time. Finally, the ILO suggests that the height of the work surface should be adjustable or that the worker should be able to adjust their height relative to the work surface, so that the arms should not be held in awkward and extreme positions.

Reid, Bush, Karwowski, and Durrani (2010) in a review of several published studies on occupational body postures and the lower extremity body region affected, developed a lower extremity discomfort guideline for standing based on published research. Standing > 2 hours/incident affected the hip and > 3 hour affected the overall lower extremity. Halim and Omar (2012) developed a Prolonged Standing Strain Index to attempt to quantify the risk levels with standing jobs and other workplace factors (e.g., posture, injuries, vibration, air quality) with minimum risk levels proposed. The Prolonged Standing Strain Index provides an overall numerical score that can be used to assign risk for a specific job into a "Safe," "Slightly unsafe," or "Unsafe" category.

The purpose of this article is to review existing scientific literature examining the potential health consequences resulting from exposure to prolonged standing at work and to document the effectiveness of various interventions aimed at reducing potential health risks. The review encompasses studies examining a variety of health consequences including musculoskeletal disorders (MSDs), such as low back and lower limb discomfort and pain, local and whole body fatigue, cardiovascular disorders, cardiovascular insufficiency (CVI), and pregnancy outcomes. The review also examines the effectiveness of interventions to reduce risk of these health outcomes, such as various floor surfaces and use of floor

mats, shoe and shoe inserts, use of support or compression hosiery (e.g., stockings), and sit-stand chairs/workstations. The following inclusion/exclusion criteria were used for assessing studies for this review: (1) Review articles and single studies must be available in English and were published in the peer reviewed literature since 1990; (2) Single studies had a clearly identifiable study population and purpose with study designs using independent/dependent variable paradigms; (3) Outcome measurements were focused primarily on prolonged standing and either one of the health issues listed above; and, (4) Intervention studies evaluated methodology aimed at reducing risk due to prolonged standing. Listed below are short narratives for each study. At the end of each section is a table that describes the study population and summarizes the major results from each study. Use of volunteers usually indicates laboratory studies. Abbreviations used in text and tables are defined initially.

Evidence of Negative Health Outcomes

Low Back Pain

A number of low back pain (LBP) measures have been associated with prolonged standing. By far, the most measured outcome is low back fatigue and discomfort. In a study of bank tellers, Roelofs and Straker (2002) reported low back discomfort with prolonged standing and Drury et al. (2008) reported that those who stand for long periods during the day reported significantly greater body parts discomfort compared to those who sit most of the day.

A number of studies have been conducted examining potential biomechanical indicators of risk of LBP due to prolonged standing. Researchers have suggested that risk of LBP is increased due to excessive coactivation of muscles involved in postural stability during prolonged standing (Marshall, Patel, & Callaghan, 2011; Nelson-Wong, Gregory, Winter, & Callaghan, 2008). Specifically, Nelson-Wong et al. (2008) postulated that prolonged standing results in a significant increase in coactivity of the gluteus medius (GM) muscles, a muscle group that serves to stabilize the pelvis during standing by abducting, medially rotating, and laterally rotating the thigh at the hip.

To investigate whether there is evidence of increased GM activity during prolonged standing, studies have been conducted (Marshall et al., 2011 and Nelson-Wong et al., 2008). Using electromyography (EMG) recordings, the

researchers found that subjects who reported low back pain showed higher coactivation of the left and right GM muscles versus those who did not report LBP during the standing task. Marshall et al. (2011) used EMG to study the endurance and strength of the GM muscles, as well as the coactivation patterns in subjects free of back pain. Their results indicated that endurance and coactivation were affected by prolonged standing and that this influenced reports of LBP.

Additional evidence of increased symptoms of LBP due to prolonged standing has been shown in studies by Nelson-Wong and Callaghan (2010a) and Nelson-Wong, Howarth, and Callaghan (2010) who examined subjects' acute biomechanical responses during a set of functional movements following prolonged standing tasks (right single leg stance, forward flexion while standing, unloaded squats and sloped surfaces). The researchers reported that following prolonged standing tasks, (1) 40% of participants developed LBP, (2) there was a decrease in vertebral joint rotation stiffness in lateral bending, (3) there was an increased excursion postural stability center of pressure measurements during unilateral stance, (4) there was no effect of standing on forward flexion, and (5) men had greater center of pressure excursion than women on the single leg-standing task. In a similar study that examined the impact of sloped surfaces and its effect on biomechanical responses to prolonged standing (Nelson-Wong & Callaghan, 2010b) found that, for those who developed LBP, standing on sloped surface reduced pain subjective reports. This finding suggests that standing on a sloped surface in some way modifies the way the GM muscles are recruited during prolonged standing to maintain pelvic stability and decreased subjective LBP reports during prolonged standing.

The remaining studies focused on the epidemiological associations between prolonged standing and reported LBP (Andersen, Haahr, & Frost, 2007; Engels, van der Gulden, Senden, & van't Hof, 1996; Tissot, Messing, & Stock, 2009; Yip, 2004). Some of the studies showed a moderate positive relationship between exposure to prolonged standing and development of LBP, whereas some concluded that there was little or no relationship. Engels et al. (1996) found that workers who reported being "hampered by standing" at work had increased risks of LBP and leg pain. Lack of definition of the term "hampered by standing," makes it difficult to determine if their exposure variable was equivalent to the term "prolonged standing" at work. Yip (2004) did not show a significant relationship between prolonged standing and risk of LBP

at work. The study, however, had some weaknesses. The hours of prolonged standing for the comparison may have been insufficient to cause a detectable difference in health outcomes and the level of severity of the LBP outcome measure in the study was very low. In a 24-month prospective study of LBP and other MSD outcomes Andersen et al. (2007) reported that prolonged standing increased reports of LBP and leg pain. In a cross-sectional study examining the relationship between LBP and prolonged standing and sitting, Tissot et al. (2009) reported increased reports of LBP in individuals who reported being constrained during standing. Constrained sitting, however, was not a significant risk factor for LBP in the study.

In a systematic review of scientific citations focusing on the potential causality of LBP resulting from exposure to prolonged standing, Roffey, Wai, Bishop, Kwon, and Dagenais (2010) concluded that they were not able to find any high-quality studies that met more than two Bradford-Hill causation criteria. The inclusion criteria for causation in the review were very restrictive and many of the studies that did show a relationship between prolonged standing and development of low back pain were excluded from their review.

We reviewed 11 studies in this section that are presented in Table 1 with the study populations and significant findings. Most studies do report symptoms of low back pain with prolonged standing.

Cardiovascular Problems

A number of studies have investigated the effects of prolonged standing on cardiovascular health outcomes. The measures studied have included carotid arteriosclerosis, leg edema, orthostatic symptoms (light headedness or dizziness), heart rate, blood pressure, and venous diseases (varicose veins, chronic venous disease and chronic venous insufficiency).

Tomei, Baccolo, Tomao, Palmi, & Rosati (1999), investigated the relationship between major venous pathologies in the legs and prolonged standing. Clinical tests indicated that prolonged standing and age were related to increased risk of venous pathologies. In a cross-sectional study examining the relationship between prolonged standing and risk of CVI, Krijnen, de Boer, Adèr, and Bruynzeel (1997a) found that 18% were diagnosed with minor CVI and that 11% were diagnosed with major CVI symptoms. In a second paper focusing on the same workers, Krijnen, de Boer, Adèr and Bruynzeel (1997b)

showed increases in leg volume with prolonged standing. Excess risk of varicose vein occurrence was reported in Danish workers working mostly in a standing position that were followed for three years after first hospitalization (Tüchsen, Krause, Hannerz, Burr & Kristensen, 2000).

In a review of studies examining the health risks associated with prolonged standing for over 8 hours, McCulloch (2002) found significant occurrence of CVI. CVI can lead to more serious health complications. McCulloch recommended employers undertake preventive measures, such as modifying job tasks to reduce risk of cardiovascular health effects due to prolonged standing.

Krause et al. (2000) examined the effect of prolonged standing on workers who were actively working over a 4-year period. Carotid intima media thickness over a 4-year period was significantly greater for men who stand at work “very much,” compared to those who do not stand at work. Partsch, Winiger, and Lun (2004) examined the effects of standing on leg edema before and after standing at work. Venous edema increased in the evening and was more pronounced in individuals with evidence of varicose veins.

Ngomo, Messing, Perrault, and Comtois (2008) measured the effect of prolonged standing on self-reported orthostatic symptoms, heart rate, and blood pressure in workers who stand for significant periods of time during their workday (i.e., average of 84–95% of the workday) and reported blood pressure changes. Sudoł-Szopińska, Panorska, Koziński, and Błachowiak (2007) conducted a prospective study examining differences in risk of chronic venous disorders for workers who are exposed to prolonged standing compared with workers who primarily are exposed to prolonged sitting in the workplace. Prolonged standing did increase chronic venous disorder symptoms. In a follow up study with similar methodology but different workers, the findings were much the same (Sudoł-Szopińska, Bogdan, Szopiński, Panorska, & Kołodziejczak, 2011). It should be noted that the term chronic venous disorder in this study might be equivalent to the term CVI in other studies. Kraemer et al. (2000) reported on cardiovascular measures from volunteers in a complex standing fatigue protocol while performing various work tasks. Results showed that after 8 hours of standing, there were significant increases in most of the physiological measures and discomfort ratings. Bahk, Kim, Jung-Choi, Jung, and Lee (2012) in a questionnaire survey of several companies reported significantly elevated risks of varicose veins and nocturnal leg cramps in workers standing >4 h/d.

Table 1 Studies Examining Prolonged Standing and LBP

Author (Year)	Study Population	Brief Summary
Engels et al. (1996)	846 nursing staff in Netherlands	Workers who reported being “hampered by standing” at work had an increased risk of LBP (OR = 3.07) and leg pain (OR = 4.9) compared to those who were not hampered by standing at work.
Roelofs and Straker (2002)	30 Australian bank tellers (24 F, 6 M)	Self-reported discomfort was highest for the low back due to standing, compared to those who sit or use a combination of sitting and standing.
Yip (2004)	144 Hong Kong nurses	In a 12-month prospective study of prolonged standing at work, authors did not show a positive relationship between standing and risk of LBP ($p = 0.19$).
Andersen et al. (2007)	5,604 Danish workers from industrial and service companies	In a 24-month prospective study of LBP and other MSD outcomes, the authors found that prolonged standing of more than 30 min/hr was associated with an increased OR for LBP (OR = 2.1) and leg pain (OR = 1.7).
Drury et al. (2008)	United States TSA baggage screeners (7 M, 5F)	Those who stand for long periods during the day reported statistically significant greater body parts discomfort in the back, legs, and feet compared to those who sit most of the day.
Nelson-Wong et al. (2008)	23 (12 M, 11 F) Canadian volunteers	Subjects who reported low back pain showed higher coactivation of the left and right GM muscles versus those who did not report LBP during the standing task.
Tissot et al. (2009)	4,517 M + 3,213 F responding to the Quebec Social + Health Survey criteria	Self-reported standing without freedom to sit was associated with increased reports of LBP for men, but not for women.
Roffey et al. (2010)	Review of 2,766 citations (only 18 met review criteria)	Concluded that it was unlikely that occupational standing or walking is independently causative of LBP.
Nelson-Wong and Callaghan (2010a,b), Nelson-Wong et al. (2010)	2010a—43 (22 M, 21 F) 2010b—16 (8 M, 6 F) Canadian volunteers	The authors reported evidence of low back problems following exposure to prolonged standing tasks. Standing on 16° sloped surface reduced LBP scores for pain development group.
Marshall et al. (2011)	24 (8 M, 16 F) Canadian volunteers	Evidence suggesting that GM endurance and coactivation were affected by prolonged standing and this influenced reports of LBP.

There seems to be agreement among the studies that prolonged standing plays a significant and potentially dangerous role in development of vascular problems for workers who must stand for long periods during the workday. Whether these exposures lead to long-term chronic leg problems is not clear, but acute health effects clearly should be prevented to the extent feasible. We identified 11 studies focusing on the effects of prolonged standing on CV problems which are listed in Table 2.

Fatigue/Discomfort

A number of studies have shown that exposure to prolonged standing tasks can increase the physical fatigue

and discomfort reported by workers (Balasubramanian, Adalarasu, & Regulapati, 2009; Drury et al., 2008; Flore et al., 2004; and Jorgensen, Hansen, Lundager, & Winkel, 1993) in several body regions. Physical fatigue is generally assessed by various physiological/biomechanical measures such as muscle electromyography, postural stability using force platforms, or muscle surface temperatures. Discomfort is typically measured using subjective rating scales that ask individuals to rate their level of pain or fatigue using a body location diagram. Findings from studies that have examined the relationship between prolonged standing and fatigue are summarized below.

Jorgensen et al. (1993) investigated the physiological/biomechanical effects and perceived discomfort/fatigue

Table 2 Studies Examining Prolonged Standing and Cardiovascular Problems

Author (Year)	Study Population	Brief Summary
Krijnen et al. (1997a)	387 M Dutch workers in a standing profession	Age and body weight were risk factors for presence of CVI and the number of years having a standing profession was identified as a risk factor for severity of CVI.
Krijnen et al. (1997b)	387 M Dutch workers in a standing profession	Leg volume increased significantly after 2 days of exposure to prolonged standing and the increase in leg volume was associated with subjective complaints in the legs.
Tomei et al. (1999)	336 M Italian workers (industrial, office + stonemasons)	Proportion of workers standing for $\geq 50\%$ of the work shift was higher in phlebopathic than in nonphlebopathic workers, and being over 40 years of age increased risk of phlebopathy.
Krause et al. (2000)	584 Finish men in Kuöpio Ischemic Heart Disease study	Prolonged standing at work associated with increased risk of development of carotid atherosclerosis, and those with stenosis or Ischemic Heart Disease (IHD) are at significantly increased risk.
Tüchsen et al. (2000)	5,940 Danish workers ages 20–59	Increased risk ratio for varicose veins for men (1.85, 95% CI 1.33–2.36) and women (2.63, 95% CI 2.25–3.02) when working mostly in standing position. Risk adjusted for age, social group and smoking.
Kraemer et al. (2000)	12 US F volunteers	Prolonged standing significantly increases orthostatic stress on workers (i.e., increased body mass and total body water, increase in popliteal and posterior vein size, significantly increase in systolic and diastolic blood pressure) and increased discomfort.
McCulloch (2002)	17 studies of workers in jobs requiring standing >8 hours from many countries	In a review of studies examining the health risks associated with prolonged standing for >8 hours, found evidence that prolonged standing was associated with a significantly occurrence of CVI.
Partsch et al. (2004)	12 Austrian workers (8 F, 4 M) who worked in compression stocking factory	Edema in the legs increased between 10.2 and 220.3 mL after standing an average of 3.2 h/d, and patients with varicose veins and with venous edema had more pronounced evening edema than individuals without visible veins.
Ngomo et al. (2008)	Study 1—34 (11 M, 23 F) health-care workers; study 2—36 (21 M, 24 F) factory + laundry from Canada	Prolonged static standing affects arterial blood pressure (BP) and may result in orthostatic intolerance (OI) and other hemodynamic changes.
Sudoł-Szopińska et al. (2007, 2011)	2007 — 160 office + bakery workers (97 F, 63 M). 2011 — 126 office + laboratory workers from Poland	Occurrence of chronic venous disorder symptoms were significantly higher for workers who work in a standing position compared with workers who primarily work in a sitting position.
Bahk et al. (2012)	2,165 (1,203 F, 962 M) South Korean workers	Significant ORs for varicose veins in women (2.99, 95% CI 1.26–7.08) and men (7.93, 95% CI 3.15–19.95) with prolonged standing >4 h/d. Nocturnal leg cramps were significant only for men (2.93, 95% CI 1.73–4.97).

with different shoes and floor surfaces during prolonged standing. The physiological/biomechanical measures were not affected by exposure time or floor/shoe condition but the subjective ratings were. Unfortunately, with the duration of standing limited to only 2 hours, the effects on the biomechanical/physiological measures may not have occurred. Flore et al. (2004) evaluated the changes in venous pressure (before work compared to after work) on standing workers compared to the controls. Standing

workers had significantly higher measures of oxidative stress before work and after work, and also the pre–post difference was higher compared to the control group. Drury et al. (2008) examined the effect of work postures on subjective fatigue in baggage security screening workers. Work posture did have an effect on ratings of body part discomfort. Balasubramanian et al. (2009) measured muscle fatigue and perceived discomfort in workers using static and dynamic standing (move between work

stations) work tasks. Fatigue rate and discomfort reports were higher in the stationary static standing posture as compared to the dynamic standing posture. Freitas, Wieczorek, Marchetti, and Duarte (2005) reported increases on measures of postural stability from adults and elderly individuals after 30 minutes of prolonged standing. Increased sway can indicate less postural control, an indicator of physical fatigue.

There appears to be general agreement among the study findings that prolonged standing without dynamic movement, even for periods as short as 30 minutes, leads to physical fatigue, discomfort, and pain in several body regions. It is also apparent that age affects how individuals respond to prolonged standing. The studies addressing the effects of prolonged standing on fatigue/discomfort are reported below in Table 3.

Pregnancy Issues

Several studies have evaluated pregnancy-related health issues associated with prolonged standing during work. Most of the studies below investigated the effects of the combination of prolonged standing and pregnancy on

birth related health outcomes, such as stillbirths, spontaneous abortions, birth weights, and preterm deliveries. One study investigated ergonomic issues related to pregnancy and prolonged standing.

Two Canadian studies evaluated several occupational factors and pregnancy outcomes from working women (McDonald, McDonald, Armstrong, Cherry, Côté et al., 1988; McDonald, McDonald, Armstrong, Cherry, Nolin et al., 1988). Increased risk ratios were associated with prolonged standing for several health outcomes, but there was no elevated risk of preterm delivery or low birth weights. Prolonged standing was defined as standing >8/day. Teitelman, Welch, Hellenbrand, and Bracken (1990) examined the impact of maternal work activity, including prolonged standing, on preterm births and low birth weight. Prolonged standing on the job was significantly associated only with increased preterm births. Klebanoff, Shiono, and Carey (1990) evaluated the effect of physical activity during pregnancy on preterm delivery and birth weight. Only standing ≥8 hours showed a moderate increase in preterm delivery rates. Eskenazi et al. (1994) studied the association of standing at work in women whose pregnancies ended in spontaneous abortion and

Table 3 Studies Examining Prolonged Standing and Fatigue/Discomfort

Author (Year)	Study Population	Brief Summary
Jorgensen et al. (1993)	8 F Danish volunteers performing letter sorting task	There was a 4-to 6-fold increase in subjective discomfort after 2 hours of work during prolonged standing.
Flore et al. (2004)	62 F Italian surgery room workers and 65 F Italian outpatient workers	Workers with predominantly standing occupations increased venous pressure (before work compared to after work) was significantly higher for the standing workers compared to the controls compared to workers who stand less, suggesting that these workers are likely at higher risk of CVI.
Freitas et al. (2005)	14 elderly volunteers and 14 adult volunteers from Brazil	Postural sway was significantly greater following a prolonged standing task and the increase in sway was attributed to fatigue. Lack of mobility had greater effects on the elderly compared to the adults.
Drury et al. (2008)	US TSA baggage screeners (7 M, 5F)	Standing posture was rated as having the highest level of discomfort, followed by sitting on a high stool, then by sitting at a desk.
Balasubramanian et al. (2009)	9 M volunteers performing assembly/disassembly task, from India	Fatigue rates in leg and lower back muscles were significantly higher ($p < .05$) in stationary standing posture as compared to dynamic standing posture, and perceived discomfort in the legs, shoulders, and overall rating of discomfort were significantly higher for static posture than for dynamic posture.

women who delivered live births. Similar to the McDonald et al. results, women standing >8 h/day at work showed risks of adverse birth outcomes. A study of Danish women between 1989 and 1991 showed significant effects on preterm delivery rates and birth weight with hours of standing (Henriksen, Hedegaard, & Secher, 1995; Henriksen, Hedegaard, Secher, & Wilcox, 1995). Mozurkewich, Luke, Avni, and Wolf (2000) examined the effects of several working conditions on adverse pregnancy outcomes. Prolonged standing was associated with slight increased risk of preterm birth. The authors estimated that if women discontinued prolonged standing one preterm birth could be eliminated for every 27–80 women. Ha et al. (2002) reported results of a study of infant birth weight and standing at work. Although prolonged standing showed a significant association with reduced birth weight, it was not clear whether standing ≥ 3 hours per day was compared with no standing. Recently, Palmer, Bonzini, Harris, Linaker, and Bonde (2013) reported an updated review with a meta-analysis of work activities and birth outcomes. Newer studies have shown absence of large effects, but there were still small increased risks for preterm delivery, low birth weight and small birth size for gestational age with prolonged standing.

Pompeii, Savitz, Evenson, Rogers, and McMahon (2005) did not find a significant relationship between physical exertion at work and risk of preterm birth or small birth size for gestational age, but exposure to prolonged standing was not assessed as h/d, but as h/week. This resulted in the variable for prolonged standing being different from several other studies, which used h/d. In some of those studies, standing >8 h/d showed a significant association with adverse pregnancy outcomes. The difference in measuring the exposure variable may be an important factor when studies report conflicting results.

Paul and Frings-Dresen (1994) investigated some of the changes that occur in working postures due to pregnancy. Results showed that the pregnant subjects would stand further from the table, with hips positioned more backwards with increased trunk flexion and with arms more extended. Postural differences in pregnant subjects were the smallest at the self-selected table height. An adjustable workplace was recommended by the authors to minimize postural changes due to pregnancy.

Most of the studies (Table 4) that investigated the effects of the combination of prolonged standing and pregnancy did indicate that prolonged standing increased

the incidences of stillbirths, spontaneous abortions and preterm deliveries. Low birth weight, however, was not consistently affected by prolonged standing and the recent meta-analysis by Palmer et al. (2013) indicates a low risk. Generally, most studies involved >3 hours of prolonged standing, but exposure duration did vary between studies.

Effectiveness of Interventions

Compression Stockings

Several studies have investigated the wearing of support stockings or hosiery during standing at work using several subjective and biomechanical/physiological measures. Krijnen, de Boer, Adèr, Osinga, and Bruynzeel (1997) examined the effects of wearing compression stockings and floor mats on workers who had been diagnosed with CVI and jobs which required prolonged standing. Only the compression stockings showed intervention effectiveness (reduced leg swelling). Kraemer et al. (2000) evaluated the effects of commercial hosiery rated light to moderate compression on reducing lower body edema and discomfort. All commercial hosiery evaluated was rated effective. McCulloch (2002) in a review of several studies done before 2000 concluded that support hose did play a positive role in reducing the symptoms of CVI and leg complaints, but the selection of the type of hose and the severity of CVI was important. Consultation with a qualified health professional was recommended, because some support hose may actually be more harmful than helpful if they restrict lower leg venous return. Jungbeck, Peterson, Danielsson, and Norgren (2002) reported a 4-week study to evaluate the effects of compression hosiery in workers who worked in standing professions. Subjective symptom ratings (leg pain, ankle swelling, tired and heavy leg, night cramps) and foot volumetry (foot volume, expelled volume, refilling rate) were objective measurements collected. Only the subjective ratings were statistically significant. Study problems may have contributed to the lack of definitive findings for the objective measures as not all participants wore the stockings every day.

Partsch et al. (2004) evaluated types of support stockings that were most effective in reducing evening edema which occurs after prolonged standing. Evening edema was significantly reduced with compression stockings. Moseley, Piller, and Walsh (2006) compared nurses and factory workers who stood for greater than 4 h/d for

Table 4 Studies Examining Prolonged Standing and Pregnancy Issues

Author (Year)	Population Studied	Brief Summary
McDonald, Armstrong, Cherry, Cote, Lavoie, et al. (1988)	56,067 Montreal, Canada, women from 11 obstetrical units	Elevated rate of spontaneous abortions for standing ≥ 8 h/d for all occupations when grouped by work requirements. When broken down by occupation (managerial, health, and clerical), workers in clerical occupations had a significantly elevated risk ($1.37 p < .01$). Risk ratios were also elevated for stillbirths in workers standing ≥ 8 h/d.
McDonald, Armstrong, Cherry, Nolin & Robert (1988)	22,761 Montreal, Canada, single live births	Did not find increased risk of preterm delivery or low birth weights from prolonged standing ≥ 8 h/d.
Teitelman et al. (1990)	1,206 pregnant Connecticut, USA, women.	Standing on the job was significantly associated with increased preterm births (OR = 2.72, 95% CI = 1.24–5.95). Birth weights were reduced in the standing group, but the association was not significant OR = 1.58, 95% CI = 0.51–4.94).
Klebanoff et al. (1990)	7,101 receiving prenatal care from NY, WA, OK, TX and LA clinical centers in USA	Prolonged standing ≥ 8 h/d showed a moderate increased OR of 1.31 (95% CI 1.10–1.71) for preterm delivery, but did not affect preterm delivery rates or birth weight.
Eskenazi et al. (1994)	1,894 women (607 spontaneous abortions + 1,287 live births) from California, USA	Women standing >8 h/d at work showed an increased adjusted OR of 1.6 (95% CI 1.1–2.3) for spontaneous abortion when compared to standing 3 h/d per day.
Henriksen, Hedegaard, Secher, & Wilcox (1995)	8,711 Danish women with single pregnancies	Women who reported >5 h/d of both standing and walking had an adjusted OR of 3.3 (95% CI 1.4–8.0) for preterm delivery when compared to women reporting ≤ 2 h/d of standing and walking. Standing only for 5 h/d when compared to ≤ 2 h/d of standing was not significant (OR = 1.2, 95% CI 0.6–2.4).
Henriksen, Hedegaard & Secher (1995)	8,711 Danish women with single pregnancies	Women who reported standing at work >5 h/d had birth weights 49 g lower than women standing ≥ 2 h/d (95% CI –108 to 10). Higher birth weights were reported, however, for women walking >2 but ≤ 5 h/d (35 g, 95% CI 8–63).
Mozurkewich et al. (2000)	160,988 women from Asian, European, and North American countries	In a meta-analysis of 29 studies (case-control, cross-sectional, prospective cohort) involving 160,988 women, authors found that prolonged standing was significantly associated with an increased risk of preterm birth (OR = 1.26, 95% CI 1.13–1.40).
Ha et al. (2002)	1,222 Chinese women working at petrochemical plant	Adjusting for confounders the association between maternal standing hours and reduced birth weight using a multivariate GAM model showed a significant ($p = 0.01$) reduction of 16.8 g.
Pompeii et al. (2005)	1,908 North Carolina, USA, pregnant women	There were no significant associations for preterm delivery outcomes or small birth size for gestational age birth for pregnant women standing >30 h per week when compared to standing 6–15 hours per week.

(continued)

Table 4 (Continued)

Author (Year)	Population Studied	Brief Summary
Palmer et al. (2013)	Meta-analysis of articles from 1966–2011; 28 studies involved standing	Preterm delivery median RR was 1.16 (1.00–1.35); small gestational age median RR was 1.00 (0.93–1.26); low birth weight RR was 1.13 (0.70–1.58) with prolonged standing >4 h/d. Excess risks are low, but still exist.

3 weeks while alternately wearing or not wearing stockings. Leg fluid volume was significantly lowered with stockings. Chiu and Wang (2007), however, in a study of footwear (nurse shoes) with prolonged standing that included a condition where the subjects wore compression hosiery, did not find that the physiological/biomechanical measurements were significantly changed with compression hosiery but only discomfort ratings were reduced. Flore et al. (2007) examined the reactive oxygen metabolites in workers wearing compression stockings in jobs requiring prolonged standing. Compression stockings showed limited effects on reactive oxygen metabolites in only the operating room nurses compared to the other workers.

Most of the studies reviewed support the use of compression stockings in the reduction of subjective complaints of leg fatigue, pain, and swelling in work requiring prolonged standing. The findings with the physiological/biomechanical measures are less convincing, although some positive findings in reduction of leg swelling and leg fluid volume have been reported. The benefits of using compression stockings are most pronounced in workers with CVI. Based on the results from these studies it would be difficult to recommend what the pressure ranges for effective stockings would be, although the higher compression surgical stockings may be more harmful and whether the wearing of stockings would have a preventive benefit in healthy workers. Major findings from the studies reviewed are listed in Table 5.

Floor Surfaces, Mats, Shoes, and Shoe Inserts

The use of floor mats and shoe inserts during prolonged standing compared to prolonged standing on hard surfaces has been evaluated as an intervention procedure to reduce symptoms of discomfort, muscle pain, leg swelling, and tiredness. Biomechanical/physiological measurements have also been investigated for their usefulness as objective measures.

Redfern and Chaffin (1995) evaluated seven flooring conditions and one shoe insert condition in comparison

with a concrete floor surface on factory workers who had jobs requiring standing for an entire 8-hours shift. The floor surfaces ranged from several types of vinyl and viscoelastic materials. Flooring did affect workers perception of discomfort and all floor surfaces were rated better than concrete. Cham and Redfern (2001) studied the relationship between workers' subjective measures of discomfort and objective measures of fatigue and discomfort on different mat flooring conditions compared with a hard vinyl floor. Floor mats with increasing elasticity, decreased energy absorption and increased stiffness rated less discomfort and fatigue with prolonged standing. Krumwiede, Konz, and Hinnen (1998) investigated the effects of floor surfaces on comfort ratings in 3 hours of prolonged standing with 1 hour on each type of floor surface. Mat compressibility (ranged from 2.2 to 8.9%) was important in the comfort ratings and all surface types rated better than concrete. Madeleine, Voigt, and Arendt-Nielsen (1998) reported physiological and subjective measurements in a group of volunteers that stood on a hard surface versus soft surface for 2 hours while performing manual repetitive work tasks (preparing letters for sales campaign). Standing on the soft surface produced less pain and discomfort, which was supported by several of the physiological and biomechanical measurements. King (2002) and Orlando and King (2004) studied assembly line workers standing for 8 h/d using different flooring conditions and also comparing floor mats and insoles. Mats and wearing insoles were rated as more comfortable than standing on the hard floor. Age, height, and job tenure showed strong correlations with some measurements. Zander, King, and Ezenwa (2004) measured changes in leg volume following an 8 h shift with different flooring conditions. Floor conditions did not significantly affect lower leg volume measurements. Recently, Lin, Chen, and Cho (2012) used a study protocol similar (4 hours prolonged standing) to the Cham and Redfern (2001) study, but added a field test component along with the laboratory study. Both floor type and shoe condition lowered foot discomfort

Table 5 Studies Examining Effectiveness of Wearing Support Stockings or Hosiery

Author (Year)	Study Population	Brief Summary
Krijnen et al. (1997)	114 Dutch workers diagnosed with CVI in 14 meat packing plants	For workers wearing compression stockings, there were significant reductions in leg complaints and leg swelling when compared to a control group. Floor mats, however, were not effective.
Kraemer et al. (2000)	12 F US volunteers	Commercial hosiery were effective in reducing edema in the ankles and legs, and also reduced the amount of venous pooling and discomfort in the lower body following prolonged standing.
McCulloch (2002)	17 studies of workers in various jobs requiring standing >8 hours from many countries	Support hose did play a positive role in reducing the symptoms of CVI and leg complaints, but the selection of the type of hose and the severity of CVI was important.
Jungbeck et al. (2002)	52 Swedish F (36 employed in department stores, hotel and restaurants and 16 surgery staff)	Subjective ratings for leg symptoms were all significantly reduced with the use of compression stockings and although the objective measures showed some improvements they were not significant. Compliance with wearing compression stockings every day was erratic and not monitored.
Partsch et al. (2004)	12 Austrian volunteers (8 F, 4 M)	In workers who stood more than 3.2 h/d, evening edema was significantly reduced with compression stockings, with the reduction being fairly linear with increasing amount of stocking compression pressure. Pressure ranges between 11 and 21 mmHg were recommended.
Moseley et al. (2006)	27 nurses (25 F, 2 M), 30 factory workers (16 F, 14 M) from Australia	Below the knee support stockings significantly lowered leg fluid volume and body part discomfort (neck, shoulder, low back) in nurses and industry workers who stood for greater than 4 h/d, but only leg discomfort symptoms were significantly lower for nurses.
Chiu and Wang (2007)	12 F volunteers in Taiwan	Compression hosiery did not affect physiological or biomechanical measurements but did reduce subjective discomfort feelings in the lower back, knee, calf and in the metatarsal and heel regions.
Flore et al. (2007)	55 operating theater nurses, 23 ironers, 65 outpatient nurses, 35 laundry workers; all Italian F.	Lower limb venous pressure increased significantly after the work shift for nurses and controls, but only the operating room nurses showed significantly higher levels of reactive oxygen metabolites when not wearing stockings. When wearing compression stockings no significant differences were noted in venous pressures or reactive oxygen metabolites levels in either group.

and shank circumference increased linearly over the test periods. Significant changes were evident after 1 h of prolonged standing in contrast to the Cham and Redfern (2001) study which reported changes in the 3rd and 4th hours.

Redfern and Cham (2000) reviewed 11 studies dating from 1972 to 2000 that had investigated prolonged standing with flooring types that used several different kinds of subjective and measurements. The authors concluded that the mixed and sometimes conflicting results from the many studies were most likely due to methodological

differences, primarily with the duration of prolonged standing. Additionally, the biomechanical/physiological measures used have led to conflicting results so there is no consensus regarding the reliability/validity for any biomechanical or physiological measures.

Sahar et al. (2007) reviewed an extensive number of studies that had included insoles for the prevention of back pain. Only randomized control trials and crossover trials were considered. Using fairly rigid acceptance criteria, only 6 of 325 citations met the selection criteria and were reviewed for intervention effectiveness. Effectiveness

of insoles was not evident but this conclusion may be influenced by not including studies with the authors' acceptable methodology. Bahk et al. (2012) found that wearing nonheeled shoes reduced the prevalence of varicose veins but not the prevalence of nocturnal leg cramps.

The benefits of floor mats/shoe inserts appear primarily to be in reducing discomfort and fatigue after several hours of prolonged standing. The use of shoe inserts is rated about the same as the most comfortable floor mats and the greatest benefits from mats/shoe inserts may occur after several hours of prolonged standing. Table 6 lists the important results from the studies reviewed.

Sit-Stand Chairs and Workstations

A number of national guidelines have suggested that sit-stand chairs with or without footrests/foot rails and/or adjustable workstations should be used for workers who must work in jobs requiring prolonged standing during the work shift. Some early studies reported conflicting results.

Irving (1992) in a survey of surgeons who stood for long periods of time during surgical procedures reported improvements in low back discomfort when using a standing/sitting pelvic-tilt chair. Nerhood and Thompson (1994) reported the results from a year-long study of workers that had sit-stand workstations installed at their company. Employees were categorized according to risk for discomfort which was determined by computer use and mobility. Discomfort ratings, injuries and illnesses were reduced. Although mobility was included in the assignment of the adjustable sit-stand workstations, this factor was not reported in the results. Oude Vrielink, Cloosterman, van der Bunt, Krijnen, & van Dieën, 1994; reported that sit-stand chairs were not an effective intervention during a 4-hour monotonous task when compared to standing. Both physiological/biomechanical and discomfort measures were evaluated but leg movement was not controlled in the experiment making it difficult to compare the conditions. Chester, Rys, & Konz, 2002, compared sitting, standing and a sit-stand chair and reported that the sit-stand chair condition increased leg volume. Study design problems (e.g., sit-stand chair position, tilt and no back rest, limited subject movement) suggest that the design of the sit-stand chair condition was not an effective intervention.

Recently more controlled studies and review articles have been published that have evaluated the intervention

effectiveness of sit-stand work stations, but not necessarily with prolonged standing but also with prolonged sitting. Husemann, Von Mach, Borsotto, Zepf, & Scharnbacher (2009) reported from a randomized controlled trial study that a sit-stand workstation (50% sitting; 25% standing; 25% break-office tasks) reduced physical and psychological complaints compared to a control group (75% seated; 25% break-office tasks). Robertson, Cirello, and Garabet (2013) conducted a randomized controlled trial study that evaluated sit-stand workstations and ergonomics pre-training. Ergonomics trained participants (more mobility with mandatory standing) reported less musculoskeletal and visual discomfort than minimally trained participants. Both groups used sit-stand workstations, but there was no prolonged standing condition.

Karakolis and Callaghan (2014) reported a systematic review of studies that included a sit-stand workstation intervention on prolonged sitting, prolonged standing or a combination of both types. Although the studies reviewed involved office environments and not manufacturing processes, the results showed sit-stand workstations lowered subjective discomforts. The review also determined that no optimal sit-stand ratio currently exists and that the optimal ratio may be specific for each worker depending on their personal and job requirements. The important results from the seven studies reviewed representative of the area reviewed are listed in Table 7.

Upon examination of the limited studies investigating the effectiveness of sit-stand chairs and workstations to reduce risk of MSDs, the effectiveness of these interventions is not clear. Conflicting results are likely due to study design weaknesses, such posture evaluation, measures of mobility, and ratios of standing to sitting. For example, none of the studies in manufacturing settings allowed the worker to use a dynamic posture during testing that allowed the worker to sit, stand, or lean against the chair, but rather the subject had to adopt a fixed posture (typically seated) for the duration of the testing. Data do suggest that use of sit-stand workstations should be considered as an ergonomic intervention for workers who must stand or sit for long periods of time in their work shift.

Summary/Conclusions

Based on review of the literature there appears to be ample evidence that prolonged standing in the work place leads to a number of negative health outcomes. The stud-

Table 6 Studies Examining Effectiveness of Floor Surfaces, Mats, Shoes, and Inserts

Author (Year)	Study Population	Brief Summary
Jorgensen et al. (1993)	8 F Danish volunteers performing letter sorting task	The effect of different shoes and floor surfaces was negligible after 2 hours of prolonged standing. Longer durations were not tested.
Redfern and Chaffin (1995)	14 US workers (8 M, 6 F)	Ratings of perceived hardness of flooring affected perception of discomfort and were correlated with measured stiffness. Slip resistant mats and extremely soft mats did not offer marked improvement over mats with more hardness. Use of shoe inserts was perceived as similar to the better-rated cushioning mats.
Krumwiede et al. (1998)	12 US college students	Mats with the highest compressibility were rated as the most comfortable, but specific mat characteristics that influenced comfort were not apparent.
Madeleine et al. (1997)	13 M Danish volunteers	Standing on the soft surface produced lower reports of unpleasantness, whereas the hard surface showed increased trunk swelling, increased EMG activity in the muscle groups, increased sway displacement, and increased muscle fatigue. Induced muscle pain was lower on the soft surface than hard surface.
Redfern and Cham (2000)	11 studies from 1972 to 2000 that investigated prolonged standing and floor types	In a review of 11 studies, authors concluded that softer floors generally resulted in reduced discomfort and fatigue compared to a hard floor, primarily in the lower extremities. Key flooring characteristics influencing discomfort include elasticity, stiffness, and thickness.
Cham and Redfern (2001)	10 US volunteers (5 M, 5 F)	Prolonged standing and floor type had significant effects on both the subjective and objective fatigue and discomfort, but only during the 3rd and 4th hour of prolonged standing.
King (2002)	22 US assembly line workers (5 M, 17 F)	A mat alone, insoles alone, and insoles with mat were rated as significantly more comfortable and less fatiguing than a hard floor condition. No significant differences, however, were noted when comparing the effects of standing on a mat versus wearing insoles or the combination of standing on a mat with insoles.
Zander et al. (2004)	16 US assembly line workers (2 M, 14 F)	Lower leg volume increased significantly from pretest to posttest while standing for 8 hours, but there were no significant differences between any flooring conditions (wood block, anti-fatigue mat, shoe insole).
Orlando and King (2004)	16 US assembly line workers (2 M, 14 F)	Use of mats and insoles reduced ratings of general fatigue, leg fatigue, and discomfort but there were no significant differences between the flooring conditions (wood block, mat, insoles) compared.
Sahar et al. (2007)	6 randomized controlled trial studies from USA, Denmark, Israel, and South Africa	Based on a review of studies examining effectiveness of insoles, authors concluded that there was no strong evidence that using insoles was effective in the prevention or treatment of back pain.
Bahk et al. (2012)	2,165 (1,203 F, 962 M) South Korean workers	Nonheeled shoes reduced prevalence of varicose veins in women (OR 0.69, 95% CI 0.49–0.97) and men (OR 0.58, 95% CI 0.35–0.96), but not on nocturnal leg cramps with prolonged standing >4 h/d.

(continued)

Table 6 (Continued)

Author (Year)	Study Population	Brief Summary
Lin et al. (2012)	Experiment 1—13 Taiwan volunteers (10 M, 3 F). Experiment 2—14 Taiwan workers (6 M, 8 F)	Expt. 1—Foot discomfort ($p < .01$) and shank circumference ($p < .01$) significantly lower on soft floor (mat). Shoe condition only significant for discomfort ($p < .01$). significant effects > 1 hour standing. Expt. 2—Thigh circumference ($p < .014$) and shank circumference ($p < .032$) greater on hard floor. Negative effects after 1 h standing.

Table 7 Studies Examining Effectiveness of Sit-stand Chairs and Workstations

Author (Year)	Study Population	Brief Summary
Irving (1992)	55 South African surgeons	Use of a sit-stand chair during surgery improved ratings of low back discomfort compared with sitting or standing only. 100% reported an improvement after operations lasting more than 2 hours and 88% of surgeons reported that they would use the sit-stand chair frequently in the future if it was available. Only 12% said they would use it occasionally.
Nerhood and Thompson (1994)	US United Parcel workers; number not stated	Body part discomfort decreased by an average of 62% and occurrence of injuries and illnesses by >50%. No significant changes on absenteeism.
Oude Vrielink et al. (1994)	6 Dutch M volunteers	Sit-stand chairs were not effective in mitigating fatigue, leg volume or discomfort ratings when compared to a standing only task.
Chester et al. (2002)	18 US (7 F, 11 M) volunteers	Found that the sit-stand chair resulted in higher leg volume changes than standing or sitting only and the most discomfort in the hips.
Husemann et al. (2009)	60 M German college volunteers	Sit-stand workstation across a 1-week period reduced musculoskeletal complaints while performing a data entry task. No significant effects noted on data entry task performance with sit-stand workstation.
Robertson et al. (2013)	22 F US volunteers	Minimal musculoskeletal and visual discomfort over a 15 day experimental period for participants who used sit-stand workstations with ergonomics training compared to participants with minimal training. 7 body region pain measures showed significant differences ($p < 0.01$) between regions. Low back, neck, and shoulder highest reported pain areas.
Karakolis and Callaghan (2014)	14 articles from various countries published between 1950 and 2011 that met review criteria	Sufficient evidence to conclude that sit-stand workstations are effective in reducing local discomfort in the low back. Some evidence that sit-stand workstations may increase reported discomfort in hand and wrist. No optimal sit-stand time ratio. No decrease in productivity noted.

ies consistently reported increased reports of low back pain, physical fatigue, muscle pain, leg swelling, tiredness, and body part discomfort due to prolonged standing. The findings from these studies were supported by intervention studies (e.g., compression stockings, floor surfaces, mats, insoles, sit-stand chairs) that show significant

reductions in the subjective measures after implementation of the intervention. The findings from objective biomechanical and physiological studies of the effects of prolonged standing frequently correlate with subjective findings, but have not been conclusive. Considerable research has been devoted to trying to establish a biome-

Key Practice Points

- Prolonged standing of ≤ 1 hour and a total per day of ≤ 4 hours is considered a safe zone according to Dutch ergonomic guidelines.
- Workplaces should not be static, but allow for mobility and use of various postures.
- Interventions such as floor mats, shoe inserts, adjustable chairs, sit-stand workstations, and compression stockings can be effective in reducing subjective effects (e.g., pain, discomfort, fatigue) from prolonged standing.
- Improved studies are needed with more precise definitions of prolonged standing, better measurements, and more rigorous study protocols.

chanical pathway for low back pain, but the final explanation is still elusive. EMG recordings of trunk muscles, center of pressure measurements using force platforms (posturography), and body kinematics have been used and the most promising result seems to be research involving the GM muscle group. The studies of prolonged standing and measurements of physical fatigue have shown some significant changes with the biomechanical/physiological measurements, but again there has been inconsistency.

Finally, there was general agreement in the reviewed literature that interventions can be effective, at least in reducing the subjective complaints from prolonged standing. Interventions that reduce the reports of LBP and venous disorders would be the most effective in prevention of chronic health problems that could occur in older workers. In reviewing the studies examining the effectiveness of interventions, we concluded that dynamic movement appeared to be the best solution for reducing risk of these health problems due to prolonged standing. The ability for workers to “have movement” during work, such as walking around, or being able to easily shift from standing to sitting or leaning posture during the work shift seemed to be a common suggestion in nearly all of the literature but needs more research. In summary, we concluded the following:

1. There is significant evidence that prolonged standing at work (primarily in one place) increases risk of LBP, cardiovascular problems, and pregnancy outcomes.
2. Interventions designed to reduce risk of adverse health outcomes due to prolonged standing can be effective. Research, however, with scientifically operational definitions of prolonged standing (e.g., hours/day, week,

month) and improved design protocols are needed to determine which biomechanical/physiological measures are reliable and valid and correlate best with the subjective reports of effects from prolonged standing. Especially important would be experimental designs that would incorporate morning and evening measurements into biomechanical or physiological measures of prolonged standing over the course of a work period (e.g., day, week, month) while manipulating the dynamics of standing (e.g., mobility, sit-stand). Morning and evening measurements would allow for the evaluation of whether outcome measurements are within normal limits of recovery or developing into chronic conditions. An ideal study protocol would take measurements daily for a typical workweek, followed by 2 days rest with measurements retaken the day of work return. Evaluation of extended work shifts beyond the normal 8 h day is also essential.

3. Workplaces should not be static. Work should be designed so that the worker can adopt various postures during the work. For example, the worker should be able to walk around rather than stand in one place, or a sit-stand chair/workstation should be provided. Also, workers should be encouraged to modify their posture so that they mix sitting, standing, leaning, and use of a foot rail or footrest as much as possible.

4. A reliable characterization of prolonged standing is needed based on a standard workday. Various groups, such as AORN and the Dutch researchers, have suggested time limits for prolonged standing, which would be effective, but we also believe that a proper work design that includes all aspects of ergonomic intervention is also necessary.

References

- Andersen, J.H., Haahr, J.P., & Frost, P. (2007). Risk factors for more severe regional musculoskeletal symptoms: A two-year prospective study of a general working population. *Arthritis & Rheumatism*, 56(4), 1355–1364.
- Bahk, J.W., Kim, H., Jung-Choi, K., Jung, M., & Lee, I. (2012). Relationship between prolonged standing and symptoms of varicose veins and nocturnal leg cramps among women and men. *Ergonomics*, 55(2), 133–139.
- Balasubramanian, V., Adalarasu, K., & Regulapati, R. (2009). Comparing dynamic and stationary standing postures in an assembly task. *International Journal of Industrial Ergonomics*, 39(5), 649–654.
- Best, M., French, G., Ciantar, J., Didzys, D., Fitzgerald, C., Moore, K. et al. (2002). Work-related musculoskeletal

- disorders in hairdressers. *Journal of Occupational Health and Safety Australia and New Zealand*, 18(313), 67–76.
- Canadian Centre for Occupational Health and Safety (CCOHS) (2014). Basic information on Standing at work. Retrieved from http://www.ccohs.ca/oshanswers/ergonomics/standing/standing_basic/html
- Cham, R., & Redfern, M.S. (2001). Effect of flooring on standing comfort and fatigue. *Human factors*, 43(3), 381–391.
- Chester, M.R., Rys, M.J., & Konz, S.A. (2002). Leg swelling, comfort and fatigue when sitting, standing, and sit/standing. *International Journal of Industrial Ergonomics*, 29(5), 289–296.
- Chiu, M.-C., & Wang, M.-J.J. (2007). Professional footwear evaluation for clinical nurses. *Applied Ergonomics*, 38(2), 133–141.
- Drury, C.G., Hsiao, Y.L., Joseph, C., Joshi, S., Lapp, J., & Pennathur, P.R. (2008). Posture and performance: Sitting vs. standing for security screening. *Ergonomics*, 51(3), 290.
- Engels, J.A., van der Gulden, J.W., Senden, T.F., & van't Hof, B. (1996). Work related risk factors for musculoskeletal complaints in the nursing profession: Results of a questionnaire survey. *Occupational and Environmental Medicine*, 53(9), 636–641.
- Eskenazi, B., Fenster, L., Wight, S., English, P., Windham, G.C., & Swan, S.H. (1994). Physical exertion as a risk factor for spontaneous abortion. *Epidemiology*, 5(1), 6–13.
- Flore, R., Gerardino, L., Santoliquido, A., Catananti, C., Pola, P., & Tondi, P. (2007). Reduction of oxidative stress by compression stockings in standing workers. *Occupational Medicine*, 57, 337–341.
- Flore, R., Gerardino, L., Santoliquido, A., Pola, R., Flex, A., Di Campli, C. et al. (2004). Enhanced oxidative stress in workers with a standing occupation. *Occupational and Environmental Medicine*, 61, 548–550.
- Freitas, S.M.S.F., Wiczorek, S.A., Marchetti, P.H., & Duarte, M. (2005). Age-related changes in human postural control of prolonged standing. *Gait and Posture*, 22(4), 322–330.
- Ha, E., Cho, S.-I., Park, H., Chen, D., Chen, C., Wang, L. et al. (2002). Does standing at work during pregnancy result in reduced infant birth weight? *Journal of Occupational and Environmental Medicine*, 44(9), 815–821.
- Halim, I., & Omar, A.R. (2012). Development of prolonged standing strain index to quantify risk levels of standing jobs. *International Journal of Occupational Safety and Ergonomics-JOSE*, 18(1), 85–96.
- Henriksen, T.B., Hedegaard, M., & Secher, N.J. (1995). Standing and walking at work and birthweight. *Acta Obstetrica et Gynecologica Scandinavica*, 74(7), 509–516.
- Henriksen, T.B., Hedegaard, M., Secher, N.J., & Wilcox, A.J. (1995). Standing at work and preterm delivery. *British Journal of Obstetrics and Gynaecology*, 102(3), 198–206.
- Hughes, N.L., Nelson, A., Matz, M.W., & Lloyd, J. (2011). AORN Ergonomic Tool 4: Solutions for prolonged standing in perioperative settings. *AORN Journal*, 93(6), 767–774.
- Husemann, B., Von Mach, C.Y., Borsotto, D., Zepf, K.I., & Scharnbacher, J. (2009). Comparisons of musculoskeletal complaints and data entry between a sitting and sit-stand workstation paradigm. *Human Factors*, 51(3), 310–320.
- International Labour Organization. (2011). Your health and safety at work: Ergonomics. Retrieved from <http://actrav.ilo.org/actrav-english/telearn/osh/ergo/ergonomi.htm>
- Irving, G. (1992). A standing/sitting pelvic tilt chair—new hope for back-weary surgeons? *South African Medical Journal*, 82(2), 131–132.
- Jorgensen, K., Hansen, L., Lundager, K., & Winkel, J. (1993). Low back muscle reactions to constrained standing in relation to shock absorbing properties of floor and shoes. In R. Nielsen, & K. Jorgensen (Eds.), *Advances in industrial ergonomics and safety V*, (pp. 279–283). Bristol, PA: Taylor & Francis.
- Jungbeck, C., Peterson, K., Danielsson, G., & Norgren, L. (2002). Effects of compression hosiery in female workers with a standing profession. *Phlebology*, 16(3), 117–120.
- Karakolis, T., & Callaghan, J.P. (2014). The impact of sit-stand office workstations on worker discomfort and productivity: A review. *Applied Ergonomics*, 45, 799–806.
- King, P.M. (2002). A comparison of the effects of floor mats and shoe in-soles on standing fatigue. *Applied Ergonomics*, 33(5), 477–484.
- Klebanoff, M.A., Shiono, P.H., & Carey, J.C. (1990). The effect of physical activity during pregnancy on preterm delivery and birth weight. *American journal of obstetrics and gynecology*, 163(5), 1450–1456.
- Kraemer, W.J., Volek, J.S., Bush, J.A., Gotshalk, L.A., Wagner, P.R., Gómez, A.L. et al. (2000). Influence of compression hosiery on physiological responses to standing fatigue in women. *Medicine and Science in Sports and Exercise*, 32(11), 1849–1858.
- Krause, N., Lynch, J.W., Kaplan, G.A., Cohen, R.D., Salonen, R., & Salonen, J.T. (2000). Standing at work and progression of carotid atherosclerosis. *Scandinavian journal of work, environment & health*, 26(3), 227–236.
- Krijnen, R.M., de Boer, E.M., Adèr, H.J., & Bruynzeel, D.P. (1997a). Venous insufficiency in male workers with a standing profession, Part 1: Epidemiology. *Dermatology*, 194(2), 111–120.
- Krijnen, R.M., de Boer, E.M., Adèr, H.J., & Bruynzeel, D.P. (1997b). Venous insufficiency in male workers with a standing profession. Part 2: Diurnal volume changes of the lower legs. *Dermatology*, 194(2), 121–126.

- Krijnen, R.M., de Boer, E.M., Adèr, H.J., Osinga, D.S., & Bruynzeel, D.P. (1997). Compression stockings and rubber floor mats: Do they benefit workers with chronic venous insufficiency and a standing profession? *Journal of Occupational and Environmental Medicine*, 39(9), 889–894.
- Krumwiede, D., Konz, S., & Hinnen, P. (1998). Standing comfort on floor mats. *Occupational Ergonomics*, 1(2), 135–143.
- Lin, Y.-H., Chen, C.-Y., & Cho, M.-H. (2012). Influence of shoe/floor conditions on lower leg circumference and subjective discomfort during prolonged standing. *Applied Ergonomics*, 43, 965–970.
- Madeleine, P., Voigt, M., & Arendt-Nielsen, L. (1998). Subjective, physiological and biomechanical responses to prolonged manual work performed standing on hard and soft surfaces. *European Journal of Applied Physiology and Occupational Physiology*, 77(1–2), 1–9.
- Marshall, P.W.M., Patel, H., & Callaghan, J.P. (2011). Gluteus medius strength, endurance, and co-activation in the development of low back pain during prolonged standing. *Human Movement Science*, 30(1), 63–73.
- McCulloch, J. (2002). Health risks associated with prolonged standing. *Work*, 19(2), 201–205.
- McDonald, A.D., McDonald, J.C., Armstrong, B., Cherry, N.M., Côté, R., Lavoie, J. et al. (1988). Fetal death and work in pregnancy. *British Journal of Industrial Medicine*, 45(3), 148–157.
- McDonald, A.D., McDonald, J.C., Armstrong, B., Cherry, N.M., Nolin, A.D., & Robert, D. (1988). Prematurity and Work in Pregnancy. *British Journal of Industrial Medicine*, 45(1), 56–62.
- Meijssen, P., & Knibbe, H.J.J. (2007). Prolonged standing in the OR: A Dutch Research Study. *AORN Journal*, 86(3), 399–414.
- Moseley, A., Piller, N., & Walsh, J. (2006). The effectiveness of below-the-knee support stockings for standing professions. *Journal of Occupational Health and Safety Australia and New Zealand*, 22(4), 335–340.
- Mozurkewich, E.L., Luke, B., Avni, M., & Wolf, F.M. (2000). Working conditions and adverse pregnancy outcome: A meta-analysis. *Obstetrics & Gynecology*, 95(4), 623–635.
- Nelson-Wong, E., & Callaghan, J.P. (2010a). Changes in muscle activation patterns and subjective low back pain ratings during prolonged standing in response to an exercise intervention. *Journal of Electromyography and Kinesiology*, 20(6), 1125–1133.
- Nelson-Wong, E., & Callaghan, J.P. (2010b). The impact of a sloped surface on low back pain during prolonged standing work: A biomechanical analysis. *Applied Ergonomics*, 41(6), 787–795.
- Nelson-Wong, E., Gregory, D.E., Winter, D.A., & Callaghan, J.P. (2008). Gluteus medius muscle activation patterns as a predictor of low back pain during standing. *Clinical Biomechanics*, 23(5), 545–553.
- Nelson-Wong, E., Howarth, S.J., & Callaghan, J.P. (2010). Acute biomechanical responses to a prolonged standing exposure in a simulated occupational setting. *Ergonomics*, 53(9), 1117–1128.
- Nerhood, H.L., & Thompson, S.W. (1994). Adjustable sit-stand workstations in the office. *Proceedings of the Human Factors and Ergonomics Society Annual meeting*, 38, 668–672.
- Ngomo, S., Messing, K., Perrault, H., & Comtois, A. (2008). Orthostatic symptoms, blood pressure and working postures of factory and service workers over an observed workday. *Applied Ergonomics*, 39(6), 729–736.
- Orlando, A., & King, P. (2004). Relationship of demographic variables on perception of fatigue and discomfort following prolonged standing under various flooring conditions. *Journal of Occupational Rehabilitation*, 14(1), 63–76.
- Oude Vrielink, H.H.E., Cloosterman, S.G.M., van der Bunt, J.A., Krijnen, R.M.A., & van Dieën, J.H. (1994). Is a sit-stand seat an appropriate alternative in standing work situations? In F. Aghazadeh (Ed.), *Advances in Industrial Ergonomics and Safety VI* (pp. 223–228). Bristol, PA: Taylor & Francis.
- Palmer, K.T., Bonzini, M., Harris, E.C., Linaker, C., & Bonde, J.P. (2013). Work activities and risk of prematurity, low birthweight, and pre-eclampsia: an updated review and meta-analysis. *Occupational and Environmental Medicine*, 70(4), 213–222.
- Partsch, H., Winiger, J., & Lun, B. (2004). Compression stockings reduce occupational leg swelling. *Dermatologic Surgery*, 30(5), 737–743.
- Paul, J.A., & Frings-Dresen, M.H.W. (1994). Standing working posture compared in pregnant and non-pregnant conditions. *Ergonomics*, 37(9), 1563–1575.
- Pompeii, L.A., Savitz, D.A., Evenson, K.R., Rogers, B., & McMahon, M. (2005). Physical exertion at work and the risk of preterm delivery and small-for-gestational-age birth. *Obstetrics and Gynecology*, 106(6), 1279–1288.
- Redfern, M.S., & Chaffin, D.B. (1995). Influence of flooring on standing fatigue. *Human Factors*, 37(3), 570–581.
- Redfern, M.S., & Cham, R. (2000). The influence of flooring on standing comfort and fatigue. *American Industrial Hygiene Association*, 61(5), 700–708.
- Reid, C.M., Bush, P.M., Karwowski, W., & Durrani, S.K. (2010). Occupational postural activity and lower extremity discomfort: A review. *International Journal of Industrial Ergonomics*, 40, 247–256.
- Robertson, M.M., Cirello, V.M., & Garabet, A.M. (2013). Office ergonomics training and a sit-stand workstation: Effects on musculoskeletal and visual symptoms and performance of office workers. *Applied Ergonomics*, 44, 73–85.

- Roelofs, A., & Straker, L. (2002). The experience of musculoskeletal discomfort amongst bank tellers who just sit, just stand or sit and stand at work. *Ergonomics SA*, 14 (2), 11–29.
- Roffey, D.M., Wai, E.K., Bishop, P., Kwon, B.K., & Dagenais, S. (2010). Causal assessment of occupational standing or walking and low back pain: Results of a systematic review. *The Spine Journal*, 10(3), 262–272.
- Sahar, T., Cohen, M.J., Ne'eman, V., Kandel, L., Odebiyi, D.O., Lev, I. et al. (2007). Insoles for prevention and treatment of back pain. *Cochrane Database of Systematic Reviews*, 4, 1–29.
- Sudoł-Szopińska, I., Bogdan, A., Szopiński, T., Panorska, A.K., & Kołodziejczak, M. (2011). Prevalence of chronic venous disorders among employees working in prolonged sitting and standing postures. *International Journal of Occupational Safety and Ergonomics*, 17(2), 165–173.
- Sudoł-Szopińska, I., Panorska, A.K., Koziński, P., & Błachowiak, K. (2007). Work-related chronic venous disease in office and bakery workers. *Occupational Ergonomics*, 7(2), 125–137.
- Teitelman, A.M., Welch, L.S., Hellenbrand, K.G., & Bracken, M.B. (1990). Effect of maternal work activity on preterm birth and low birth weight. *American Journal of Epidemiology*, 131(1), 104–113.
- Tissot, F., Messing, K., & Stock, S. (2005). Standing, sitting and associated working conditions in the Quebec population in 1998. *Ergonomics*, 48(3), 249–269.
- Tissot, F., Messing, K., & Stock, S. (2009). Studying the relationship between low back pain and working postures among those who stand and those who sit most of the working day. *Ergonomics*, 52(11), 1402–1418.
- Tomei, F., Baccolo, T.P., Tomao, E., Palmi, S., & Rosati, M.V. (1999). Chronic venous disorders and occupation. *American Journal of Industrial Medicine*, 36(6), 653–665.
- Tüchsen, F., Krause, N., Hannerz, H., Burr, H., & Kristensen, T.S. (2000). Standing at work and varicose veins. *Scandinavian Journal of Work, Environment & Health*, 26(5), 414–420.
- Werner, R.A., Gell, N., Hartigan, A., Wiggermann, N., & Keyserling, W.M. (2010). Risk factors for foot and ankle disorders among assembly plant workers. *American Journal of Industrial Medicine*, 53, 1233–1239.
- Yip, V.Y.B. (2004). New low back pain in nurses: Work activities, work stress and sedentary lifestyle. *Journal of Advanced Nursing*, 46(4), 430–440.
- Zander, J.E., King, P.M., & Ezenwa, B. (2004). Influence of flooring conditions on lower leg volume following prolonged standing. *International Journal of Industrial Ergonomics*, 34, 279–288.

Earn nursing contact hours



Rehabilitation Nursing is pleased to offer readers the opportunity to earn nursing contact hours for its continuing education articles by taking a posttest through the ARN website. The posttest consists of questions based on this article, plus several assessment questions (e.g., how long did it take you to read the articles and complete the posttest?). A passing score on the posttest and completing of the assessment questions yield one nursing contact hour for each article.

To earn contact hours, go to www.rehabnurse.org and select the "Education" page. There you can read the article again, or go directly to the posttest assessment by selecting "RNJ online CE." The cost for credit is \$10 per article. You will be asked for a credit card or online payment service number.

The contact hours for this activity will not be available after **June 30, 2017**.

The Association of Rehabilitation Nurses is accredited as a provider of continuing nursing education by the American Nurses Credentialing Center's Commission on Accreditation (ANCC-COA).