



LETTER TO THE EDITOR

Editorial Note. Letters to the Editor are peer reviewed to ensure that the arguments are reasonable and clearly expressed. However, letters may express a particular opinion rather than a balanced interpretation. Authors of papers commented on are invited to reply, but neither the journal nor peer reviewers should be assumed to support the arguments made.

Human and Animal Evidence Supports Lower Occupational Exposure Limits for Poorly-Soluble Respirable Particles

Letter to the Editor re: 'Low-Toxicity Dusts: Current Exposure Guidelines Are Not Sufficiently Protective' by Cherrie, Brosseau, Hay and Donaldson

Eileen D. Kuempel¹, Michael D. Attfield²,
Leslie T. Stayner³ and Vincent Castranova^{4,5}

1.National Institute for Occupational Safety and Health, Education and Information Division, 1090 Tusculum Avenue, Cincinnati, OH, USA;

2.National Institute for Occupational Safety and Health, Division of Respiratory Disease Studies, Morgantown, WV, USA;

3.University of Illinois at Chicago, School of Public Health, Division of Epidemiology and Biostatistics, Chicago, IL, USA;

4.National Institute for Occupational Safety and Health, Health Effects Laboratory Division, Morgantown, WV, USA;

5.Department of Basic Pharmaceutical Sciences, School of Pharmacy, West Virginia University, Morgantown, WV, USA

E-mail: ekuempel@cdc.gov

Submitted 10 April 2014; revised 30 June 2014 revised version accepted 8 July 2014

We commend the overall evaluation by [Cherrie *et al.* \(2013\)](#) of the current occupational exposure limits (OELs) for respirable poorly-soluble low toxicity (PSLT) particles. As described in that paper, the epidemiological studies provide compelling evidence that exposure to PSLT at the current OELs has been associated with adverse health effects,

including pulmonary fibrosis and lung function deficits. In contrast to [Cherrie *et al.* \(2013\)](#), we discuss here that the chronic inhalation studies in animals also provide evidence of the adverse pulmonary effects of PSLT.

For example, we would like to clarify or correct the following statements (p. 688, 2nd column):

(1) ...the phenomenon of rat lung overload has little relevance for human lung response at high lung burden of low-toxicity dust.

This statement is not entirely supported by the scientific evidence. While it is correct that differences have been observed in the rat and human lung clearance and retention kinetics for respirable particles, these differences have been well described, and can be accounted for, using lung dosimetry models in humans (Kuempel *et al.*, 2001a,b; Gregoratto *et al.*, 2010, 2011) and rats (Tran *et al.*, 1999, 2000; Anjilvel and Asgharian, 1995). Because of the slower pulmonary clearance in humans, particles can build up in the lungs at exposures below those that would cause overloading in rats (Snipes, 1989; Kuempel *et al.*, 2000; Kuempel and Tran, 2002). This build up is associated with the movement of particles into the alveolar interstitium of the mammalian lungs (Nikula *et al.*, 1997, 2001). Only at overloading do the particle lung burdens in rats reach the higher levels that have been reported in coal miners, i.e., up to 10 mg g^{-1} lungs or more in rats (Morrow, 1988; Muhle *et al.*, 1990; Bellmann *et al.*, 1991; Oberdörster *et al.*, 1992) and in humans (Attfield *et al.*, 1994; Kuempel *et al.*, 1997; Tran and Buchanan, 2000).

In addition, the lung responses to these high lung burdens of PSLTs can be qualitatively similar in rats and humans (Castranova, 2000; Attfield *et al.*, 2012). As described in Cherrie (2013), the lung inflammatory and fibrotic response to PSLT has been observed to be greater in rats than that in mice or hamsters (Bermudez *et al.*, 2002, 2004; Elder *et al.*, 2005). Consistent with a persistent inflammation and cell proliferation mode-of-action, rats (but not mice or hamsters) showed increased lung cancer response to chronic inhalation of PSLT particles (ILSI, 2000; Schins and Knaapen, 2007). Yet, studies in mice and hamsters were negative for some particles that have been classified as known human carcinogens (Mauderly, 1997). The proportion of lung tumor types has been observed to differ in humans and rats exposed to respirable particles (ILSI, 2000), although some of the differences may be due to smoking-related tumors in humans and terminology differences (Maronpot *et al.*, 2004; NIOSH, 2011b). An ILSI (2000) expert panel concluded that the rat is a useful model for non-neoplastic lung responses to poorly-soluble particles and that (in the absence of mechanistic data to the contrary) it is also relevant to identifying potential carcinogenic hazards in humans.

A comparative pathology study (Green *et al.*, 2007) showed that humans had a more severe centriacinar fibrotic response than rats exposed to silica or coal dust; while rats had a more severe intra-alveolar response (i.e., inflammation, lipoproteinosis, and alveolar epithelial hyperplasia) after chronic exposures to silica, talc, or coal dust. Yet in both species, silica and talc were more inflammogenic than coal dust, and the severity of response was associated with the dose (i.e., increased centriacinar fibrosis at high versus low coal dust). Thus, in humans a high dust load in the alveolar interstitium can lead to septal thickening and fibrosis even in the absence of an observed macrophage-mediated decline in clearance.

(2) As discussed herein, high lung burdens of low-toxicity dust in humans cause COPD primarily and pneumoconiosis only in the very highly exposed coalmining populations (Kuempel *et al.*, 2001a,b).

In contrast to this statement that COPD and pneumoconiosis can develop 'only in very highly exposed coalmining populations...', significant risks of both COPD and pneumoconiosis have been predicted and observed in US coal miners exposed at the current US coal mine dust standard of 2 mg m^{-3} (and as low as 0.5 mg m^{-3}) over a working lifetime (NIOSH, 1995, 2011a; Attfield and Seixas, 1995; Suarthana *et al.*, 2011).

(3) There is some evidence for the increased retention of dust in the alveolar interstitial compartment in the lungs of coalminers at the very high lung burdens they experience (Kuempel *et al.*, 2001a,b; Gregoratto *et al.*, 2010).

This statement is only partially correct. In addition to the evidence from coal miners with relatively high dust exposures (Kuempel, 2000; Tran and Buchanan, 2000), the increased long-term retention of dust in human lungs has also been observed in workers with relatively low dust exposures (Gregoratto *et al.*, 2010, 2011).

(4) However, this is in contrast with the virtual cessation of clearance and subsequent rapid buildup of dust that accompanies rat

lung overload, nor do we see the pathological consequences of rat lung overload – proliferation, fibrosis, and cancer – in humans.

The second part of this statement is inaccurate because the human lung responses to respirable particles can be similar to those observed in rats exposed to overloading doses of PSLT (as discussed above). Also, the related statement on p. 686 of [Cherrie et al. \(2013\)](#) that coal workers' pneumoconiosis (CWP) is characterized by 'minimal fibrosis (macules) and occasional nodule formation' is incomplete because it does not mention the severe form of CWP, progressive massive fibrosis – which is associated with disability and early death – or the evidence that both simple CWP and PMF are associated with significantly increased mortality from nonmalignant respiratory diseases ([Attfield and Kuempel, 2008](#)).

In addition, a quantitative comparison of rat- and human-based risk estimates for lung cancer associated with exposure to different types of poorly-soluble particles (coal mine dust, carbon black, titanium dioxide, or crystalline silica) shows statistically consistent risk estimates, given consideration of the imprecision in the animal and human data ([Kuempel et al., 2009](#); [NIOSH, 2011b](#)). An increase in lung cancer has also been observed in recent studies of coal miners in the USA ([Graber et al., 2014](#)) and UK ([Miller and MacCalman, 2010](#)). Although the excess was most strongly related to silica in the UK study, it was most strongly associated with coal dust in the US study (in which the silica exposure estimates were also more uncertain).

In summary, we agree with [Cherrie et al. \(2013\)](#) that the epidemiologic literature provides adequate evidence that workplace exposure to PSLT has been associated with occupational lung disease and impairment at concentrations below the current OELs. This view has been accepted by some regulatory institutions. For example, in the USA, the regulatory exposure limit for respirable coal mine dust is to be reduced from 2 mg m^{-3} as a multi-shift average concentration to 1.5 mg m^{-3} as a single-shift (8-hr time-weighted average) concentration, starting in August 2016 ([MSHA, 2014](#)). In addition, the rat studies provide supporting evidence with qualitatively and quantitatively consistent findings to those in the human studies. Yet, more research is needed to reduce uncertainty about the rodent dose-response models compared to humans.

Such evidence is important for risk assessment and OEL development of other types of airborne particles (e.g., nanoparticles) for which sufficient human data may not be available.

DISCLAIMER

The findings and conclusions in this letter are those of the authors and do not necessarily represent the views of the National Institute for Occupational Safety and Health.

REFERENCES

Anjilvel S, Asgharian B. (1995) A multiple-path model of particle deposition in the rat lung. *Fundam Appl Toxicol*; 28: 41–50.

Attfield MD, Vallyathan V, Green FHY. (1994) Radiographic appearances of small opacities and their correlation with pathology grading of macules, nodules, and dust burden in the lungs. *Ann Occup Hyg*; 38 (Suppl. 1): 783–9.

Attfield MD, Kuempel ED. (2008) Mortality among U.S. underground coal miners: a 23-year follow-up. *Am J Ind Med*; 51: 231–45.

Attfield MD, Seixas NS. (1995) Prevalence of pneumoconiosis and its relationship to dust exposure in a cohort of U.S. bituminous coal miners and ex-miners. *Am J Ind Med*; 27: 137–51.

Attfield M, Castranova V, Kuempel E et al. (2012). "Coal." In: *Patty's industrial hygiene*. 6th edn. Wiley online. Available at <http://onlinelibrary.wiley.com/doi/10.1002/0471435139.tox016.pub2/pdf>.

Bellmann B, Muhle H, Creutzberg O et al. (1991) Lung clearance and retention of toner, utilizing a tracer technique, during chronic inhalation exposure in rats. *Fundam Appl Toxicol*; 17: 300–13.

Bermudez E, Mangum JB, Asgharian B et al. (2002) Long-term pulmonary responses of three laboratory rodent species to subchronic inhalation of pigmentary titanium dioxide particles. *Toxicol Sci*; 70: 86–97.

Bermudez E, Mangum JB, Wong BA et al. (2004) Pulmonary responses of mice, rats, and hamsters to subchronic inhalation of ultrafine titanium dioxide particles. *Toxicol Sci*; 77: 347–57.

Castranova V. (2000) From coal mine dust to quartz: mechanisms of pulmonary pathogenicity. *Inhal Toxicol*; 12 (Suppl. 3): 7–14.

Cherrie JW, Brosseau LM, Hay A et al. (2013) Low-toxicity dusts: current exposure guidelines are not sufficiently protective. *Ann Occup Hyg*; 57: 685–91.

Elder A, Gelein R, Finkelstein JN et al. (2005) Effects of subchronically inhaled carbon black in three species. I. Retention kinetics, lung inflammation, and histopathology. *Toxicol Sci*; 88: 614–29.

Graber JM, Stayner LT, Cohen RA *et al.* (2014) Respiratory disease mortality among US coal miners; results after 37 years of follow-up. *Occup Environ Med*; 71: 30–9.

Green FH, Vallyathan V, Hahn FF. (2007) Comparative pathology of environmental lung disease: an overview. *Toxicol Pathol*; 35: 136–47.

Gregoratto D, Bailey MR, Marsh JW. (2010) Modelling particle retention in the alveolar-interstitial region of the human lungs. *J Radiol Prot*; 30: 491–512.

Gregoratto D, Bailey MR, Marsh JW. (2011) Particle clearance in the alveolar-interstitial region of the human lungs: model validation. *Radiat Prot Dosimetry*; 144: 353–6.

ILSI (International Life Sciences Institute). (2000) The relevance of the rat lung response to particle overload for human risk assessment: a workshop consensus report. *Inhal Toxicol*; 12: 1–17.

Kuempel ED, O'Flaherty EJ, Stayner LT *et al.* (1997) Relationships between lung dust burden, pathology, and lifetime exposure in an autopsy study of U.S. coal miners. *Annals Occup Hygiene* 41 (Suppl. 1): 384–9.

Kuempel ED. (2000) Comparison of human and rodent lung dosimetry models for particle clearance and retention. *Drug Chem Toxicol*; 23: 203–22.

Kuempel ED, O'Flaherty EJ, Stayner LT *et al.* (2001) A biomathematical model of particle clearance and retention in the lungs of coal miners. *Regul Toxicol Pharmacol*; 34: 69–87.

Kuempel ED, Tran CL, Smith RJ *et al.* (2001) A biomathematical model of particle clearance and retention in the lungs of coal miners. II. Evaluation of variability and uncertainty. *Regul Toxicol Pharmacol*; 34: 88–101.

Kuempel ED, Tran CL. (2002) Comparison of human lung dosimetry models: implications for risk assessment. *Ann Occup Hyg*; 46 (Suppl. 1): 337–41.

Kuempel ED, Smith RJ, Dankovic DA *et al.* (2009) Rat- and human-based risk estimates of lung cancer from occupational exposure to poorly-soluble particles: a quantitative evaluation *J Phys: Conf Ser, Inhaled Particles*; 151 (1): 012011. X 2009 Mar.

Maronpot RR, Flake G, Huff J. (2004) Relevance of animal carcinogenesis findings to human cancer predictions and prevention. *Toxicol Pathol*; 32 (Suppl. 1): 40–8.

Mauderly JL. (1997) Relevance of particle-induced rat lung tumors for assessing lung carcinogenic hazard and human lung cancer risk. *Environ Health Perspect*; 105 (Suppl. 5): 1337–46.

Miller BG, MacCalman L. (2010) Cause-specific mortality in British coal workers and exposure to respirable dust and quartz. *Occup Environ Med*; 67: 270–6.

Morrow PE. (1988) Possible mechanisms to explain dust overloading of the lungs. *Fundam Appl Toxicol*; 10: 369–84.

MSHA (2014). 30 CFR Parts 70, 71, 72, *et al.* Lowering miners' exposure to respirable coal mine dust, including continuous personal dust monitors; Final Rule. Mine Safety and Health Administration, U.S. Department of Labor. *Fed Reg*; 79, May 1, pp. 24813–994. Available at <https://www.federalregister.gov/articles/2014/05/01/2014-09084/>

lowering-miners-exposure-to-respirable-coal-mine-dust-including-continuous-personal-dust-monitors. Accessed 27 June 2014.

Muhle H, Creutzenberg O, Bellman B *et al.* (1990) Dust overloading of lungs: investigations of various materials, species differences, and irreversibility of effects. *J Aerosol Med*; 3 (Suppl. 3): S111–28.

Nikula KJ, Avila KJ, Griffith WC *et al.* (1997) Lung tissue responses and sites of particle retention differ between rats and cynomolgus monkeys exposed chronically to diesel exhaust and coal dust. *Fundam Appl Toxicol*; 37: 37–53.

Nikula KJ, Vallyathan V, Green FHY *et al.* (2001) Influence of dose on the distribution of retained particulate material in rat and human lungs. In *Proceedings of Particulate Matter 2000, January 23–28, 2000*. Charleston, SC: Air and Waste Management Association.

NIOSH. (1995) *Criteria for a recommended standard: occupational exposure to respirable coal mine dust*. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 95-106.

NIOSH. (2011a) *Current intelligence bulletin 64: coal mine dust exposures and associated health outcomes: a review of information Published Since 1995*. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 2011-172.

NIOSH. (2011b). *Current intelligence bulletin 63: occupational exposure to titanium dioxide*. Cincinnati OH: U.S. Department of Health and Human Services, Centers for Disease Control, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 2011-160.

Oberdörster G, Ferin J, Morrow PE. (1992) Volumetric loading of alveolar macrophages (AM): a possible basis for diminished AM-mediated particle clearance. *Exp Lung Res*; 18: 87–104.

Schins RP, Knaapen AM. (2007) Genotoxicity of poorly soluble particles. *Inhal Toxicol*; 19 (Suppl. 1): 189–98.

Snipes MB. (1989) Long-term retention and clearance of particles inhaled by mammalian species. *Crit Rev Toxicol*; 20: 175–211.

Suarthana E, Laney AS, Storey E *et al.* (2011) Coal workers' pneumoconiosis in the United States: regional differences 40 years after implementation of the 1969 Federal Coal Mine Health and Safety Act. *Occup Environ Med*; 68: 908–13.

Tran CL, Buchanan D. (2000). *Development of a biomathematical lung model to describe the exposure-dose relationship for inhaled dust among U.K. coal miners*. Edinburgh: Institute of Occupational Medicine. IOM Research Report TM/00/02.

Tran CL, Jones AD, Cullen RT *et al.* (1999) Mathematical modeling of the retention and clearance of low-toxicity particles in the lung. *Inhal Toxicol*; 11: 1059–76.

Tran CL, Buchanan D, Cullen RT *et al.* (2000) Inhalation of poorly soluble particles. II. Influence of particle surface area on inflammation and clearance. *Inhal Toxicol*; 12: 1113–26.