

# Lung Transplantation Is Increasingly Common Among Patients With Coal Workers' Pneumoconiosis

**David J. Blackley, Dr.,<sup>1</sup>\*** **Cara N. Halldin, PhD,<sup>1</sup>** **Kristin J. Cummings, MD,<sup>2</sup>**  
**and A. Scott Laney, PhD<sup>1,3</sup>**

**Background** *The prevalence of coal workers' pneumoconiosis (CWP) in U.S. coal miners has increased, and severe presentations are increasingly common.*

**Methods** *We describe trends in lung transplantation during 1996–2014 for recipients with a primary diagnosis of CWP or pneumoconiosis unspecified, and we summarize recipient characteristics and estimate survival.*

**Results** *A total of 47 transplants were included; nearly three-quarters were performed during 2008–2014. All recipients were male, 96% were white, and the mean age was 56 years. Mean FEV<sub>1</sub>% was 35%; mean FVC% was 53%. Mean time on a waitlist was 155 days, and 60% of transplants were bilateral. Median survival was 3.7 years.*

**Conclusions** *These transplants reflect the use of a scarce resource for an entirely preventable disease, and highlight the need for enhanced efforts to reduce coal mine dust exposures.* Am. J. Ind. Med. Published 2015. This article is a U.S. Government work and is in the public domain in the USA.

**KEY WORDS:** *coal workers' pneumoconiosis; lung transplantation; occupational lung disease; coal mining*

## INTRODUCTION

The prevalence of coal workers' pneumoconiosis (CWP) in U.S. coal miners has increased since the late 1990s [Laney and Weissman, 2014]. This increase has been most pronounced in central Appalachia, and evidence

suggests severe presentations of the disease are becoming more common. Studies of these miners during the last decade have described rapid onset and progression of CWP [Antao et al., 2005; Wade et al., 2011], increases in the prevalence of progressive massive fibrosis (the most severe form of CWP) [Blackley et al., 2014] and CWP resembling silicosis [Laney et al., 2010; Cohen et al., 2015], changes in radiographic and histopathologic disease patterns [Petsonk et al., 2013], severe disease among surface miners [Halldin et al., 2015], increases in years of potential life lost attributable to CWP [Mazurek et al., 2009], and rising numbers of state and federal black lung disability compensation claims [U.S. Department of Labor, 2014].

There is no cure for CWP and other forms of coal mine dust lung disease, which include chronic obstructive pulmonary disease (COPD) and dust-related diffuse fibrosis. Supportive therapies include bronchodilators for airflow limitation, antibiotics for respiratory infections, supplemental oxygen to manage hypoxemia, and smoking cessation programs. Those with radiographic evidence of CWP should

\*National Institute for Occupational Safety Health, Respiratory Health Division, Surveillance Branch, Morgantown, West Virginia

<sup>2</sup>National Institute for Occupational Safety and Health, Respiratory Health Division, Field Studies Branch, Morgantown, West Virginia

<sup>3</sup>Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, Respiratory Health Division, Morgantown, West Virginia

Contract grant sponsor: Health Resources and Services Administration; Contract grant number: 234-2005-37001C.

\*Correspondence to: Dr. David J. Blackley, National Institute for Occupational Safety and Health, Respiratory Health Division, Surveillance Branch, 1095 Willowdale Rd, Mailstop HG900.2, Morgantown, WV 26505. E-mail: dblackley@cdc.gov

Accepted 23 November 2015

DOI 10.1002/ajim.22551. Published online in Wiley Online Library  
 (wileyonlinelibrary.com).

be advised to move to a less-dusty work environment, but this is not always possible, and disease can continue to progress even after cessation of coal mine dust exposure [Kimura et al., 2010; Mason et al., 2010]. Supportive therapies do not slow progression of CWP, and recent reports have described the use of lung transplantation among patients with end-stage CWP [Enfield et al., 2012; Hayes et al., 2012].

Little is known about lung transplantation among U.S. coal miners, and trends for the procedure have not been previously reported.

## METHODS


We analyzed de-identified Organ Procurement and Transplantation Network patient-level data provided by the United Network for Organ Sharing. To be eligible for inclusion, patients must have had a primary diagnosis of CWP or pneumoconiosis unspecified and received a single or bilateral lung transplant in the U.S. during 1996–2014, because the first lung transplant for CWP was performed in 1996 [Enfield et al., 2012]. We describe the trend in lung transplants for these patients, summarize recipient demographic and clinical characteristics, and estimate median survival using the Kaplan–Meier method.

## RESULTS

Twenty-four individuals with CWP and 23 with pneumoconiosis unspecified received lung transplants during the study period (Fig. 1). Thirty-four (72%) of these transplants were performed during 2008–2014. Occupational history data were not available, but 19 of the 23 patients with pneumoconiosis unspecified were residents of West Virginia, Virginia, Kentucky, or Pennsylvania—Appalachian states that are leading coal producers. All 47 patients with CWP or pneumoconiosis unspecified were male, 96% were white, and the mean age was 56 years. Mean FEV<sub>1</sub>% predicted at transplant registration was 35%, and mean FVC% predicted was 53%. These patients were on a waitlist for a mean of 155 days, and 28 (60%) received a bilateral transplant. Medicare paid for 14 (30%) of the procedures and private insurance paid for 16 (34%); other government insurance schemes paid for 17 transplants (36%). Twenty-two recipients were deceased as of December 31, 2014, and median post-transplant survival was 3.7 years.

## DISCUSSION

Although transplants for CWP are relatively rare compared to the total number performed in the U.S., the procedure is becoming increasingly common, and its



**FIGURE 1.** Frequency of lung transplants in patients with a primary diagnosis of coal workers' pneumoconiosis or pneumoconiosis unspecified, United States, 1996–2014; based on organ procurement and transplantation network data provided by the United Network for organ sharing; data current as of March 6, 2015.

success with respect to survival is not entirely clear. Insufficient data exist to determine survival times while appropriately controlling for competing factors. Two studies on survival following lung transplantation for CWP have been reported. Hayes and colleagues identified eight cases of lung transplantation for CWP in Kentucky and reported good clinical outcomes and survivability post transplantation [Hayes et al., 2012]. In contrast, Enfield and colleagues examined data for 30 recipients determined to have CWP and reported lower survival compared to patients undergoing lung transplants for COPD, silicosis, and idiopathic pulmonary fibrosis [Enfield et al., 2012].

The pattern of lung transplants performed for patients with CWP and pneumoconiosis unspecified, which likely represents CWP, is consistent with the recent resurgence of progressive massive fibrosis in coal miners [Blackley et al., 2014]. Though commonly thought of as a well characterized disease of antiquity, novel questions regarding our understanding of this disease continue to emerge. These results add to a growing body of evidence pointing to increased severity of coal mine dust lung disease in the U.S. and highlight the mounting consequences of this ongoing epidemic. A lung transplant for end-stage CWP is an extreme intervention, but in the absence of proven curative treatments, U.S. coal miners will continue to undergo the procedure. These transplants reflect the use of a scarce resource for an entirely preventable disease, and highlight the need for enhanced efforts to reduce coal mine dust exposures.

## AUTHORS' CONTRIBUTIONS

DJB analyzed and interpreted the data, drafted the report, approved the final version, and is accountable for its accuracy and content. CNH interpreted the data, provided critical feedback during writing, and approved the final version. KJC interpreted the data, provided critical feedback during writing, and approved the final version. ASL analyzed and interpreted the data, drafted the letter, and approved the final version.

## ETHICS REVIEW AND APPROVAL

The NIOSH IRB determined this study did not require IRB review (HSRB 14-DRDS-NR05), and activities were covered by a signed data use agreement with UNOS.

## ACKNOWLEDGMENTS

This work was supported in part by Health Resources and Services Administration contract 234-2005-370011C. The content is the responsibility of the authors alone and does not necessarily reflect the views of the Department of Health and Human Services.

## DISCLOSURE BY AJIM EDITOR OF RECORD

Steven Markowitz declares that he has no conflict of interest in the review and publication decision regarding this article. This paper was prepared and written by NIOSH employees as part of their employment.

## REFERENCES

Antao VC, Petsonk EL, Sokolow LZ, Wolfe AL, Pinheiro GA, Hale JM, Attfield MD. 2005. Rapidly progressive coal workers' pneumoconiosis in the United States: Geographic clustering and other factors. *Occup Environ Med* 62:670–674.

Blackley DJ, Halldin CN, Laney AS. 2014. Resurgence of a debilitating and entirely preventable respiratory disease among working coal miners. *Am J Respir Crit Care Med* 190:708–709.

Cohen R, Petsonk E, Rose C, Young B, Regier M, Najmuddin A, Abraham J, Churg A, Green F. 2015. Lung pathology in U.S. coal workers with rapidly progressive pneumoconiosis implicates silica and silicates. *Am J Respir Crit Care Med* DOI: 10.1164/rccm.201505-1014OC

Enfield KB, Floyd S, Barker B, Weder M, Kozower BD, Jones DR, Lau CL. 2012. Survival after lung transplant for coal workers' pneumoconiosis. *J Heart Lung Transplant* 31:1315–1318.

Halldin CN, Reed WR, Joy GJ, Colinet JF, Rider JP, Petsonk EL, Abraham JL, Wolfe AL, Storey E, Laney AS. 2015. Debilitating lung disease among surface coal miners with no underground mining tenure. *J Occup Environ Med* 57:62–67.

Hayes D, Diaz-Guzman E, Davenport DL, Zwischenberger JB, Khosravi M, Absher KJ, Hoopes CW. 2012. Lung transplantation in patients with coal workers' pneumoconiosis. *Clin Transplant* 26:629–634.

Kimura K, Ohtsuka Y, Kaji H, Nakano I, Sakai I, Itabashi K, Igarashi T, Okamoto K. 2010. Progression of pneumoconiosis in coal miners after cessation of dust exposure: A longitudinal study based on periodic chest X-ray examinations in Hokkaido, Japan. *Intern Med* 49:1949–1956.

Laney AS, Petsonk EL, Attfield MD. 2010. Pneumoconiosis among underground bituminous coal miners in the United States: Is silicosis becoming more frequent? *Occup Environ Med* 67:652–656.

Laney AS, Weissman DN. 2014. Respiratory diseases caused by coal mine dust. *J Occup Environ Med* 56:S18–S22.

Mason RJ, Broaddus VC, Martin T, King Jr, TE, Schraufnagel D, Murray JF, Nadel JA. 2010. Murray and Nadel's textbook of respiratory medicine: 2-volume set. Elsevier Health Sci.

Mazurek J, Laney A, Wood J. 2009. Coal workers' pneumoconiosis-related years of potential life lost before age 65 years—United States, 1968–2006. *Morb Mortal Wkly Rep* 58:1412–1416.

Petsonk EL, Rose C, Cohen R. 2013. Coal mine dust lung disease. New lessons from old exposure. *Am J Respir Crit Care Med* 187:1178–1185.

U.S. Department of Labor, Office of Workers' Compensation Programs, Accessed October 2014, <http://www.dol.gov/owcp/dcmwc/statistics/PartCClaimsFiled.htm>

Wade W, Petsonk E, Young B, Mogri I. 2011. Severe occupational pneumoconiosis among West Virginian coal miners: One hundred thirty-eight cases of progressive massive fibrosis compensated between 2000 and 2009. *Chest* 139:1458–1462.