

The Feasibility of Using Electronic Health Records to Describe Demographic and Clinical Indicators of Migrant and Seasonal Farmworkers

Christina Socias, Yuanyuan Liang, George Delclos, Julie Graves, Ed Hendrikson & Sharon Cooper

To cite this article: Christina Socias, Yuanyuan Liang, George Delclos, Julie Graves, Ed Hendrikson & Sharon Cooper (2016) The Feasibility of Using Electronic Health Records to Describe Demographic and Clinical Indicators of Migrant and Seasonal Farmworkers, *Journal of Agromedicine*, 21:1, 71-81, DOI: [10.1080/1059924X.2015.1074633](https://doi.org/10.1080/1059924X.2015.1074633)

To link to this article: <http://dx.doi.org/10.1080/1059924X.2015.1074633>



Accepted author version posted online: 19 Oct 2015.



Submit your article to this journal 



Article views: 53



View related articles 



View Crossmark data 

The Feasibility of Using Electronic Health Records to Describe Demographic and Clinical Indicators of Migrant and Seasonal Farmworkers

Christina Socias^a, Yuanyuan Liang^a, George Delclos^a, Julie Graves^b, Ed Hendrikson^c, and Sharon Cooper^a

^aSchool of Public Health, The University of Texas Health Sciences Center, Houston, Texas, USA; ^bAmerican University of the Caribbean, Cupecoy, Dutch Lowlands St. Maarten; ^cSalud Family Health Center, Fort Lupton, Colorado, USA

ABSTRACT

Few extensive, national clinical databases exist on the health of migrant and seasonal farmworkers (MSFWs). Electronic health records (EHRs) are increasingly utilized by Federally Qualified Health Centers (FQHCs) and have the potential to improve clinical care and complement current surveillance and epidemiologic studies of underserved working populations, such as MSFWs. The aim of this feasibility study was to describe the demographics and baseline clinical indicators of patients at an FQHC by MSFW status. The authors described 2012 patient demographics, social history, medical indicators, and diagnoses by MSFW status from the de-identified EHR database of a large, multisite Colorado Migrant Health Center (MHC). Included in the study were 41,817 patients from 2012: 553 (1.3%) MSFWs, 20,665 (49.4%) non-MSFWs, and 20,599 (49.3%) who had no information in the MSFW field. MSFWs were more often male, married, employed, Hispanic, and Spanish-speaking compared with non-MSFWs. The most frequent diagnoses for all patients were hypertension, overweight/obesity, lipid disorder, type 2 diabetes, or a back disorder. Although there were significant missing values, this feasibility study was able to analyze medical data in a timely manner and show that Meaningful Use requirements can improve the usability of EHR data for epidemiologic research of MSFWs and other patients at FQHCs. The results of this study were consistent with current literature available for MSFWs. By reaching this vulnerable working population, EHRs may be a key data source for occupational injury and illness surveillance and research.

KEYWORDS

Electronic health records; farmworkers; Meaningful Use; occupational health

Background

US farmworkers

Farmworkers represent a substantial number of US workers, with estimates ranging from approximately 750,000¹ to over 3 million.² Many farmworkers are seasonal and/or migrant workers and are diverse in nationality, socioeconomic status, and country of birth. According to the National Agriculture Worker Survey (NAWS) by the US Department of Labor, approximately 81% of farmworkers reported being Hispanic or Latino, with about 75% born in Mexico, and an estimated 53% unauthorized to work in the United States. Data from NAWS indicated that the average income of a farmworker family is between \$15,000 and \$17,499 and only 23% have health insurance.³ Due to their unique laborious working situations and variety of backgrounds, farmworkers have an increased risk of adverse health

outcomes and experience many barriers to health care access such as literacy, language, financial, or even transportation.^{2,3}

An estimated 862,808 migrant and seasonal farmworkers and their family members received care at a Migrant Health Center (MHC) in 2011.⁴ Clinical data acquired at MHCs may be an important source for surveillance of the health and safety risks faced by this population, beyond workers' compensation or other occupational health data sources that do not always capture the full picture of farmworker health issues.⁵

Occupation information in electronic health records

In 2011, the Institute of Medicine (IOM) released several detailed Meaningful Use recommendations, including incorporating occupation and industry

information into electronic health records (EHRs). The committee determined that including occupation and industry information into health records may contribute to Meaningful Use.⁶ Meaningful Use, as defined by the Centers for Medicare and Medicaid Services (CMS), stipulates that EHRs should be used in a meaningful way to improve health care delivery according to certain quality and quantity indicators.⁷ The first two stages of Meaningful Use focus on adapting EHRs and participating in Health Information Exchange (HIE). HIE is the process of safely securing and exchanging patient health information across two or more providers in support of Meaningful Use quality measures. Meaningful Use guidance of Stage 3 focuses on improving population health “through better-informed diagnoses, more focused treatment plans, and improved and streamlined return-to-work guidance.”^{6,7} Occupation information will be an important demographic indicator to facilitate the population health activities of Stage 3.

EHR acceptance provides an opportunity for many researchers across various fields in the United States to increase understanding of disease, injury, surveillance, and other public health efforts among patient populations.⁸ US providers and patients generally support adoption of EHRs and HIE,^{9–11} including among economically disadvantaged groups.¹² Public health researchers have already begun using EHR data for surveillance^{13–16} and population studies of infectious disease.⁸ HIE has shown promise in improving patient care and reducing superfluous cost by connecting providers to share patient information in a secure manner.^{17–19} Some researchers even found reduced work burden on small providers that use EHRs.²⁰ Patients are now able to access their own health information using “patient portals” through secure Internet access.²¹ The increase in use and acceptance contributes to the availability of clinical data for public health research in populations that were previously difficult to reach.

Given the availability of clinical data from clinics that serve farmworkers, this may be an efficient way to study specific injuries and illnesses related to farmworkers as compared with other patients served at these clinics. MHCs offer a unique view of farmworkers from an occupational health perspective if coupled with technical, clinical, and

information support from nonprofits such as the National Center for Farmworker Health²² and the Migrant Clinicians Network.²³ MHCs allow farmworkers to access health care and may be vital to understanding the health issues of this vulnerable population. Clinical data available from MHC providers may contribute to the national discussion on demonstrating meaningful use by the routine inclusion of occupation and industry within certified EHRs.⁶ This study used a merged, de-identified HIE database to describe the patients that visited a Colorado Federally Qualified Health Center (FQHC) in 2012 by farmworker status, including demographics, medical indicators, and the most frequent medical diagnoses.²⁴

Methods

EHR data source and population

The medical records data in this study were obtained from the Community-Based Research Network (CBRN), a National Institute of Environmental Health Sciences (NIEHS)-funded research collaborative.²⁴ This cross-sectional study captured a snapshot of adult patient EHR data from January 1 through December 31, 2012. Operating as nine community health clinics and a mobile unit, the MHC prioritizes service to all low-income, medically underserved migrant and seasonal farmworker community members.²⁵ The database included patient demographics, vital sign measurements, diagnosis codes, and a free-text variable with some occupation and/or industry information.

Data acquisition and use

A data use agreement was approved and signed by the three principal investigators of the CBRN project to outline specific data handling responsibilities and intended use. Data were acquired using a Virtual Private Network to securely transfer data from the clinic to the HIE manager as a pilot, through both SQL and HL7 coding.²⁶ The data were stored in secure warehousing units and accessed by the HIE management analysts only for database building and reporting. The HIE assigned encrypted, unique identification codes for each patient and removed identifying information.

Human subjects considerations

After institutional review board (IRB) approval from the University of Texas Health Science Center Committee for the Protection of Human Subjects and CBRN steering committee approval, a de-identified clinical data set was transferred for analysis from the HIE management organization to the investigator using password-protected, University of Texas, secure network storage. This study analyzed the EHR database as though the records were abstracted from paper health records. Patients received their usual medical care at the clinics with no active participation required by the patients. A previous IRB-approved Waiver of Authorization was established for release of records (in place of an individual informed consent).

Patient descriptors

Encrypted records for patients aged 18 years or older were reviewed for errors and completeness and matched on unique identification codes to distinguish and remove any duplicate records. Demographic factors available from the patient record included age, marital status, sex, primary language, race, ethnicity, insurance, and status of the following—employment, migrant farmworker, and seasonal farmworker. Migrant and seasonal farmworkers were grouped together as “MSFW.” Medical indicators from the patient encounter included body mass index (BMI), temperature, heart rate (pulse), and systolic and diastolic blood pressures. BMI was calculated in the medical record, provided in the data set, and edited for out-of-range values. Additionally, a binary variable was generated to indicate patients with a $BMI \geq 30 \text{ kg/m}^2$ to reflect obesity versus other. After de-duplication, the last value of the medical indicator in 2012 was included in the study (for example, the latest encounter’s BMI value and similarly for other variables). If multiple values existed for the last encounter, an average was taken. International Classification of Disease—Clinical Modification (ICD-CM) diagnosis codes available from the EHR database were also de-duplicated and counted by illness, excluding codes for preventive/nonillness

procedures such as vaccination or pregnancy. From the diagnosis codes, binary variables were created to indicate patients with a 2012 diagnosis of one of the most frequently counted diagnoses for all patients, including hypertension (e.g., 401.X-405.X), overweight/obesity, type 2 diabetes, lipid disorder, back disorder, anxiety, joint disorders, abdominal pain, drug abuse, and soft tissue disorders. A binary variable was generated for patients with a history of any surgical procedures. The following variables were available but excluded due to text entry and a lack of systematic coding scheme: allergies, education, income, family history, medication list, and social history (drug use, alcohol use, sexual orientation).

The patient demographic population is described as a whole and by MSFW status in [Table 1](#). Since a large portion of patient records were missing MSFW status information, we categorized patients with null values for MSFW status as a third comparison group to understand whether these patients were different. Medical indicators including the 10 most frequent clinical diagnoses based on ICD-9-CM code counts for all patients and by MSFW status were summarized in [Table 2](#). The chi-square test or Fisher’s exact test was used to examine differences for categorical variables (e.g., sex, race, etc.) between MSFWs versus non-MSFW patients and patients with no information (No-Info) regarding MSFW status. The Bonferroni adjustment was used to adjust the resulting P value for pairwise multiple comparisons for the three MSFW groups (MSFW vs. Non-MSFW, MSFW vs. No-Info, and Non-MSFW vs. No-Info) to make sure that the overall Type I error rate was controlled at .05. For continuous variables (e.g., blood pressure, BMI, etc.), the Kruskal-Wallis nonparametric test was used to assess the equality of the median across the three MSFW groups. Further, the Dunn procedure was used to conduct pairwise multiple comparisons of MSFW, Non-MSFW, and No-Info groups, and Bonferroni-adjusted P values were computed for each pairwise comparison. All statistical tests were conducted at a significance level of .05 (two-sided). Statistical analyses were conducted using Stata version 11.0 (StataCorp, College Station, TX).²⁷

Table 1. Demographics of All Patients and Patients by MSFW Status (Colorado Health Center, 2012).

Characteristic	All patients	MSFW	Non-MSFW	No-Info	P value ^a
Number of patients	41,817	553	20,665	20,599	
Age, years					
Median (IQR)	39 (28,52)	39 (29,51)	44 (31,55)*	36 (26,48)*	<.001
Total patients (N)	41,817	553	20,665	20,599	
Sex					
Males, n (%)	15,930 (38.5)	280 (50.6)	7,713 (37.3)*	7,937 (39.3)*	<.001
Total patients (N)	41,421	553	20,665	20,203	
Marital status					
Married, n (%)	8,598 (41.2)	306 (56.6)	8,242 (40.8)*	50 (42.0)*	<.001
Total patients (N)	20,847	541	20,187	119	
Employment					
Employed, n (%)	6,659 (34.5)	174 (46.4)	6,451 (34.2)*	34 (39.1)	<.001
Total patients (N)	19,302	375	18,840	87	
Language					
English, n (%)	14,231 (67.0)	124 (22.5)	14,063 (68.3)*	44 (57.1)*	<.001
Total patients (N)	21,233	552	20,604	77	
Race/ethnicity, n (%)					
All Hispanic	21,172 (53.7)	452 (85.8)	9,704 (49.1)*	11,016 (57.6)*	<.001
Non-Hispanic: White	16,020 (40.6)	70 (13.3)	8,984 (45.4)	6,966 (36.4)	
Non-Hispanic: Other	2,260 (5.73)	5 (0.95)	1,095 (5.5)	1,160 (6.1)	
Total patients (N)	39,452	527	19,783	19,142	
Insurance, n (%)					
Self-pay	734 (4.0)	175 (42.6)	556 (3.1)*	3 (3.0)*	<.001
Federal/state	15,027 (81.7)	209 (50.9)	14,746 (82.5)	72 (72.7)	
Private Insurance	2,626 (14.3)	27 (6.6)	2,575 (14.4)	24 (24.2)	
Total patients (N)	18,387	411	17,877	99	

^aOverall difference using χ^2 test (categorical variables) or Kruskal-Wallis test (continuous variable, age).*Significant difference compared with the MSFW group using Dunn Procedure (Kruskal-Wallis), Fisher's Exact test, or χ^2 test with the Bonferroni correction for multiple comparisons.**Table 2.** Description of Medical Indicators (Colorado Health Center, 2012).

Characteristic	All patients	MSFW	Non-MSFW	No-Info	P value ^a
Number of patients	41,817	553	20,665	20,599	
Obesity					
BMI ≥ 30 , n (%)	14,832 (39.3)	227 (42.5)	8,525 (43.9)	6,080 (34.3)*	<.001
Total patients (N)	37,702	534	19,433	17,735	
BMI, kg/m ²					
Mean (SD)	29.3 (7.0)	29.7 (5.8)	30.1 (7.3)	28.5 (6.6)*	<.001
Total patients (N)	37,702	534	19,433	17,735	
Systolic BP, mm Hg					
Median (IQR)	120 (110,130)	122 (112,132)	120 (110,128)*	120 (110,132)*	<.001
Total patients (N)	41,554	551	20,605	20,398	
Diastolic BP, mm Hg					
Median (IQR)	78 (70,82)	78 (70,84)	79 (70,84)*	77 (70,82)*	<.001
Total patients (N)	41,554	551	20,605	20,398	
BP categories, n (%)					
Normal	15,213 (36.6)	180 (32.7)	6,667 (32.4)	8,366 (41.0)*	<.001
Prehypertension	18,017 (43.4)	257 (46.6)	9,058 (44.0)	8,702 (42.7)	
Hypertension stage 1	6,220 (15.0)	81 (14.7)	3,519 (17.1)	2,620 (12.8)	
Hypertension stage 2	2,105 (5.1)	33 (6.0)	1,361 (6.6)	710 (3.5)	
Total patients (N)	41,554	551	20,605	20,398	
Heart rate, beats/min					
Median (IQR)	80 (72,84)	77 (70,84)	80 (72,86)*	78 (72,84)*	<.001
Total patients (N)	40,143	549	20,333	19,261	
Temperature, °F					
Mean (SD)	98.03 (0.6)	98.07 (0.6)	98.01 (0.6)	98.06 (0.6)	<.001
Total patients (N)	37,381	526	18,683	18,172	
Past surgery					
Yes, n (%)	27,657 (66.1)	311 (56.2)	15,126 (73.2)*	12,220 (59.3)*	<.001
Total patients (N)	41,817	553	20,665	20,599	

^aOverall difference using χ^2 test (categorical) or Kruskal-Wallis test (continuous).*(Pairwise) significant difference compared with MSFWs based on Fisher's Exact test, χ^2 test, ANOVA, or Dunn procedure (Kruskal-Wallis).

Results

We identified 98,175 patient records from a Colorado FQHC in 2012 (Figure 1). Patient records were excluded from the analysis if the patient was less than 18 years old ($n = 28,979$) or if there was no provider visit during 2012 ($n = 27,379$). As a result, a total of 41,817 patients were included in the study. Of these, 553 were indicated as MSFW patients; 20,665 were identified as non-MSFW patients; and 20,599 had no information in the MSFW field.

Patient demographics

Patient demographics (age, sex, marital status, employment, preferred language, race/ethnicity, and insurance) were described and compared across MSFW status groups in Table 1. Among the 41,817 study patients, the median age was 39 years, with males constituting 38.5% of the population. Less than half of patients (41.2%) identified as married; 34.5% were designated as employed; and the majority of patients identified English as their primary language (67.0%). Most patients identified as either Hispanic (53.7%) or non-Hispanic white (40.6%). The majority of patients utilized federal or state funding as payment for medical service (81.7%) compared with self-pay (4.0%) or private insurance (14.3%).

As compared with non-MSFWs, MSFWs were significantly younger, had a higher proportion of males, and a higher percentage of patients

identified as married. MSFWs were identified as employed more often than non-MSFWs, and they identified their primary language as English significantly less often. The majority of MSFWs were Hispanic (85.8%), compared with about half of non-MSFWs (49.1%). MSFWs were more likely to use self-pay rather than the federal or state funding to pay for their care compared with the Non-MSFW group.

As compared with the No-Info group, MSFWs were significantly older and had a significantly higher proportion of male patients. MSFWs had a significantly higher percentage of married patients, a somewhat higher employed percentage, and a significantly lower percentage of primary English speakers. MSFWs had a significantly higher percentage of Hispanic patients, with only half (50.9%) of MSFWs utilizing federal/state funding for medical compared with 72.7% of the No-Info group using federal/state funding. The No-Info group had a high number of missing values in the marital status, employment, language, and insurance variables. Although not displayed in a table, we also compared the Non-MSFW group with the No-Info group. Compared with the Non-MSFW group, the No-Info group was significantly younger ($P < .001$) and had significantly more Hispanic patients ($P < .001$). Other comparisons were not significantly different.

Patient medical indicators

Table 2 displays patient medical indicators, and Table 3 displays the most frequent diagnoses.

BMI and obesity

Overall, the MSFW group had a lower mean BMI compared with the Non-MSFW group, but had a significantly higher mean BMI compared with No-Info group. The percentage of obesity was significantly higher in the MSFW and Non-MSFW groups compared with the No-Info group.

Blood pressure (BP) and BP categories

Median blood pressure was 120/78 mm Hg for all patients at the last visit. The MSFW group had significantly higher median systolic BP compared with Non-MSFW and No-Info group patients.




Figure 1. Electronic patient record inclusion diagram (Colorado Health Center, 2012).

Table 3. Top 10 Most Common Diagnoses Among Patients With Available Diagnosis Codes, Excluding Diagnostic or Preventive Codes (Colorado Health Center, 2012).

Diagnosis	All patients	MSFW	Non-MSFW	No-Info	P value ^a
Number of patients	41,817	553	20,665	20,599	
Prevalence of top diagnoses, ^b n (%)					
Hypertension	8,087 (19.3)	122 (22.1)	5,860 (28.4)*	2105 (10.2)*	<.001
Overweight/obese	6,047 (14.5)	190 (34.4)	4,543 (22.0)*	1,314 (6.4)*	<.001
Lipid disorder	5,604 (13.4)	77 (13.9)	4,069 (19.7)*	1,458 (7.1)*	<.001
Type 2 diabetes	4,954 (11.9)	73 (13.2)	3,416 (15.5)	1,465 (7.1)*	<.001
Back disorder	4,861 (11.6)	68 (12.3)	1,576 (7.5)	3,217 (15.6)*	<.001
Anxiety	4,434 (10.6)	44 (8.0)	3,203 (15.5)*	1,187 (5.8)	<.001
Joint disorder	3,735 (8.9)	44 (8.0)	2,482 (12.0)*	1,209 (5.9)	<.001
Abdominal pain	3,591 (8.6)	41 (7.41)	2,145 (10.4)*	1,405 (6.8)	<.001
Drug abuse	3,519 (8.42)	47 (8.5)	2,741 (13.3)*	731 (3.6)*	<.001
Soft tissue disorder	3,158 (7.55)	39 (7.05)	2,080 (10.1)*	1,039 (5.0)	<.001

^aOverall difference using χ^2 test.^bEach patient may have had more than one diagnoses, but multiple diagnoses of the same health outcome were removed. The top 10 diagnoses were based on all patients rather than each individual group.

*Significant difference compared with MSFWs based on Fisher's Exact test.

MSFW patients had significantly lower diastolic BP compared with Non-MSFW but significantly higher compared with No-Info. When BP was categorized according to normal, prehypertension, hypertension stage 1, and hypertension stage 2,²⁸ more than half of all patients were categorized as prehypertensive (43.4%) or hypertensive (20.1%). The percentage of normal BPs was similar between MSFW and the Non-MSFW group patients, but the No-Info group had a significantly higher percentage of patients with a normal BP.

Vitals

Median heart rate was 80 beats per minute and mean patient temperature was 98.03°F. Overall, 66.1% of patients had a previous surgical history. MSFWs had significantly lower heart rate, lower percentage of past surgery, and no difference in temperature compared with Non-MSFW or No-Info group patients.

Diagnoses

Each patient had a list of diagnoses associated with 2012 clinic visits. A patient may have had more than one diagnosis for a given visit, so we removed any duplicate diagnosis code entries. The most frequent diagnosis codes for all patients are displayed in Table 3. The composition of the most frequent diagnoses in all groups was similar, but the leading diagnosis in MSFWs was overweight/obesity, followed by hypertension. The prevalence of each of the 10 most frequent indicators was lower in the No-Info group, except for back

disorder diagnoses, which were the leading diagnosis (15.6%). Note that diagnoses were not mutually exclusive; rather, it was possible for one patient to have multiple diagnoses. Therefore, prevalence percentages across diagnoses rows cannot be added.

Discussion

No national HIE currently exists in the United States. As potentially the first anchor for a national MSFW-focused HIE, this study addresses a lack of current, geographically diverse medical data available on farmworkers around the country and encourages progress in HIE. The goal of this study was to characterize MSFW population served at a Colorado clinic by demographic and medical indicators. The first clinic added to the HIE was a Colorado FQHC that serves as a MHC to nine communities in northeastern Colorado. Patient record data from 2012 were transferred from the health center to the HIE manager. Few comprehensive studies exist to characterize demographics of MSFWs nationally, and specifically in Colorado.

Summary of results

The demographic findings from this study were similar to those found in the NAWS, including the high percentage of Hispanics among MSFWs and using self-pay to cover medical expenses.^{3,29} The No-Info group had a significant number of

missing records in marital status, employment, language, and insurance (greater than 99% missing). These missing values made it difficult to interpret significant differences between the No-Info group and the other patient groups. The No-Info group is suspected to have a combination of MSFW and Non-MSFW group patients and was, therefore, still included in the patient description as a separate patient group.

In regard to medical indicators, a high percentage of patients were obese (39.3%), and the mean BMI was just under obese classification. Median systolic and diastolic blood pressures were close to the pre-hypertension cutoff, with only 36.6% of patients below the prehypertension cutoff at the last visit (120/80 mm Hg). Most patients had a normal heart rate and temperature; over 66% of patients reported some form of previous surgery. The most frequent diagnoses assigned to patients in 2012 were chronic diseases, with similarly high diagnosis prevalence for MSFW, Non-MSFW, and No-Info group patients. Most differences between patient groups observed in regards to medical indicators were statistically significant, but not always clinically significant. The large sample size and use of nonparametric statistical methods may have resulted in statistical significance rather than clinical significance.

The results of this study are consistent with the sparse literature available on MSFWs.³⁰⁻³³ Most of the information available regarding the health of MSFWs is specific to certain regions, such as in California,³³ Michigan,^{32,34} New York,³⁰ or Texas.³² Limited national information is available from the NAWS.^{3,29} In the medical literature available, type 2 diabetes and hypertension were always among the top diagnoses, as shown in this study.^{30,32} Most studies found similar demographics among MSFWs, especially the high percentage of Hispanic ethnicity and lower percentage of health insurance coverage. Nearly 25 years ago, type 2 diabetes and hypertension were among the most frequent diagnoses for MSFW patients at MHCs in Michigan, Indiana, and Texas.³² However, our feasibility study examined adult patients by MSFW status for Colorado. For hypertension, the 2005–2006 National Health and Nutrition Examination Survey (NHANES)³⁵ estimated that 29% of US adults were stage 1 hypertensive or higher (above 140/90 mm Hg). In our study, the

prevalence for all patients was (20.1%), which was lower than the NHANES, but we did not have information about medication use or controlled hypertension. Future expansion of this analysis to other FQHCs will provide the opportunity to fill a gap in data available for research and greater geographic representation. Upcoming EHR requirements regarding Meaningful Use may help bridge the gap between medicine and epidemiologic research. Rather than conducting extensive studies in patient populations, epidemiologists may be able to utilize EHR and HIE databases to establish exposure and health outcome associations while addressing Meaningful Use Stage 3 requirements, including to “improve population health.”⁶ Systematically incorporating industry and occupation into EHRs can streamline public health research for vulnerable populations as outlined in Stage 3 of Meaningful Use.

Study limitations

As a feasibility study, we identified some limitations regarding EHR data. A substantial number of data were missing for nearly half of all patients in 2012, including information about the MSFW variable that we used for comparison. Depending on the distribution of missing values, this could reduce the validity of our results. Further research in this database revealed that missing values may be due to errors during the data transfer from the clinic to the HIE and subsequently to the researcher, rather than a lack of collection by clinic staff³⁶ or refusal by the patient to provide information upon request. Speculation of possible errors included inconsistency in file selection or file corruption during transfer from the HIE formatting to a data set. Another possibility for error is in variability of data collection methods between different providers, which has been found as an issue in other hospital departments.³⁷ The records from this study come from a Colorado patient population, rely on the ICD-9-CM coding scheme, and may not be generalizable to all MSFWs, FQHC users, general clinic patients, or the general US population. When conducting research in a working population, the healthy worker effect should also be considered as a potential selection bias where workers may be healthier than nonworkers.³⁸

In regard to the MSFW category, information about standard clinic practice was obtained from CBRN clinic and community partners in order to understand the function and reliability of this variable. The MSFW category is currently documented in the EHR system as a billing assignment. Instead, there should be a special population designation category in addition to the billing field. This would allow the documentation of both the MSFW status and, in the event that the patient has third-party coverage, the billing information. Providers at this clinic track MSFW status using a non-payor-related variable that was not transferred with this database but is critical in future analyses to ensure the distinction between the two identifiers and better understand the percentage of MSFWs who are utilizing the different payor systems (personal communications Ed Hendrikson, April 18, 2013; E. Roberta Ryder, June 13, 2013). Using a non-payor-related variable may reduce misclassification of MSFWs and improve understanding of MSFW medical needs.

On a national basis, FQHCs designated as a MHC receive a portion of their annual grant as PHS 330(g) funding in order to address the cost of care for MSFW patients. The amount of this grant is calculated on the basis of a projection of the number of patients to be seen. Historically, new start awards are made on an average of between \$250 and \$400 per MSFW user per year, seldom enough to cover the cost of care for more than two or three medical encounters, let alone provision of dental, behavioral, or ancillary services such as outreach, transportation, interpretation, or environmental services, which are essential to serving this population. Therefore, if a MSFW qualifies for third-party reimbursement, such as Medicare, the Colorado Indigent Care Program (CICP), or has commercial insurance, the clinic staff might not document a patient's MSFW status, thinking to shift the cost to that other payor in order to receive a higher reimbursement. This is probably not intentional, but merely a difference in eligibility.

The quality of data received may also benefit from improved coding or consistency in collection. For example, information in the education variable was not systematically collected and was, therefore, difficult to analyze. The field was an open text field, rather than a prescribed selection.

In the insurance field, there were specific options: self-pay, federal programs (Medicaid/Medicare), state programs (CICP, Children's State Insurance program), or private insurance. Obtaining patient level of educational attainment can easily be conducted in a systematic coding scheme modeled after any national survey. Other examples of variables obtained but inconsistently coded included income, allergies, family histories, and medication. Additionally, certain variables were collected but missing completely, such as smoking status. Smoking status is particularly important when examining hypertension.²⁸ Including information on demographics is part of Meaningful Use standards, and refining these variables using coding schemes and improved quality will be important at later stages of Meaningful Use. Using information such as education, smoking status, family histories, etc., the provider can improve their care delivery, and epidemiologists can improve recommendations for understanding health risk factors and future interventions. Beyond Meaningful Use, including demographics is important in epidemiologic research to understand health issues of MSFWs, since few resources currently exist.

Study strengths

Generally speaking, health information technology has a positive impact on care,³⁹ but continued efforts are necessary to improve the usability and quality of EHR data. This study outlines some of the challenges of using EHR data and provides practical suggestions for improvement. Initially, it took several years to build partnerships in the community, the clinic, and with academia. As the CBRN was forming, much effort was taken to compile a medical database for research and found that the HIE provided the best foundation for the future of a national database. This study also shows how quickly data can be transferred from the clinic to the HIE, de-identified, and transferred to the researcher (less than 1 year). Finally, a major strength of this study was the opportunity to have a built-in comparison group by having access to non-MSFW patients attending the same clinic.

EHRs have the capacity to collect many different patient indicators, but the best quality (consistency

in coding and entry) was found in those required or directly related to Meaningful Use. Stage 1, Step 5 Meaningful Use requirements include demographics and vital signs assessment during a patient visit. Core measure 5 under Stage 1 involves collecting preferred language, gender, race, ethnicity, and date of birth. Core measure 8 involves collecting vital signs, including height, weight, blood pressure, and BMI calculation for adults.⁷ Variables not required by Meaningful Use at the time, such as family history, had no coding scheme and no systematic entry method, whereas variables that were required by Meaningful Use had the lowest percent of missing values and were the simplest to understand.

Conclusion

The purpose of this study was to assess the feasibility of using EHRs to conduct epidemiologic research in a specific working population, with future studies to include prospective, longitudinal research in MSFWs and other populations that utilize FQHCs. From the FQHC database, we were able to successfully describe the overall patient population and compare known MSFWs to known non-MSFWs, but we also had a significant issue with missing demographic and medical data. Our findings indicated that chronic disease is similar among this group of MSFW patients compared with other patients groups at this clinic. Also, our findings were consistent with the current literature on hypertension risk factors, including age, obesity, and sex. Future studies may include additional data collection and improved validation of data from the clinic to the researcher. Medical chart review may supplement current survey methods such as the NAWS⁴⁰ in describing MSFW health outcomes. This may be an effective way to improve epidemiologic research in populations with a weak knowledge base, specifically MSFWs. By reaching this vulnerable working population, EHR data may be vital for occupational injury and illness research in support of Stage 3 Meaningful Use.

Acknowledgments

This project would not have been possible without the CBRN steering committee, including Bobbi Ryder, Nick Heyer, Eva

Shipp, Deborah Del Junco, and Melissa Valerio, Centex Systems Support Services, and the enthusiastic support of Salud Family Health Center.

Funding

This project was supported by the NIH National Institute for Environmental Health Sciences, (grant no. 1RC4ES019405-01), Southwest Center for Agricultural Safety and Health at The University of Texas HSC at Tyler (CDC National Institute for Occupational Safety and Health [NIOSH] Cooperative Agreement No. U50 OH07541), and Southwest Center for Occupational and Environmental Health, a NIOSH Education and Research Center (grant no. 5T42OH008421). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of NIH, CDC, or the Community-Based Research Network.

References

1. Bureau of Labor Statistics, US Department of Labor. Occupational Outlook Handbook, 2012–13 Edition. Agricultural Workers 2012. Available at: <http://www.bls.gov/ooh/farming-fishing-and-forestry/agricultural-workers.htm>. Accessed April 8, 2012.
2. Larson AC. Migrant and Seasonal Farmworker Enumeration Profiles Study. Available at: <http://www.ncfh.org/enumeration/PDF10%20Texas.pdf>. Published 2000. Accessed April 10, 2010.
3. Carroll D, Samardick RM, Bernard S, Gabbard S, Hernandez T. Findings from the National Agricultural Workers Survey (2001–2002). A Demographic and Employment Profile of United States Farm Workers. Available at: http://www.doleata.gov/agworker/report9/naws_rpt9.pdf. Published 2005. Accessed April 7, 2012.
4. US Department of Health and Human Services. Health Resources and Services Administration (HRSA) Primary Care: The Health Center Program “Special Populations”. Available at: <http://bphc.hrsa.gov/about/specialpopulations/index.html>. Accessed April 11, 2012.
5. Brower MA, Earle-Richardson GB, May JJ, Jenkins PL. Occupational injury and treatment patterns of migrant and seasonal farmworkers. *J Agromedicine*. 2009;14: 172–178.
6. Committee on the Respiratory Protection Curriculum for Occupational Health Nursing Programs, Institute of Medicine. *Incorporating Occupational Information in Electronic Health Records: Letter Report*. Washington, DC: The National Academies Press; 2011.
7. Centers for Medicare and Medicaid Services. 2014 Definition Stage 1 of Meaningful Use. Available at: https://www.cms.gov/Regulations-and-guidance/Legislation/EHRIncentivePrograms/meaningful_use.html. Published 2014. Accessed January 12, 2015.

8. Lurio J, Morrison FP, Pichardo M, et al. Using electronic health record alerts to provide public health situational awareness to clinicians. *J Am Med Inform Assoc.* 2010;17:217–219.

9. Ancker JS, Edwards AM, Miller MC, Kaushal R. Consumer perceptions of electronic health information exchange. *Am J Prev Med.* 2012;43:76–80.

10. Patel VN, Abramson E, Edwards AM, Cheung MA, Dhopeshwarkar RV, Kaushal R. Consumer attitudes toward personal health records in a beacon community. *Am J Manag Care.* 2011;17:e104–e120.

11. Ancker JS, Silver M, Miller MC, Kaushal R. Consumer experience with and attitudes toward health information technology: a nationwide survey. *J Am Med Inform Assoc.* 2013;20:152–156.

12. Ancker JS, Barron Y, Rockoff ML, et al. Use of an electronic patient portal among disadvantaged populations. *J Gen Intern Med.* 2011;26:1117–1123.

13. Kukafka R, Ancker JS, Chan C, et al. Redesigning electronic health record systems to support public health. *J Biomed Inform.* 2007;40:398–409.

14. Calderwood MS, Platt R, Hou X, et al. Real-time surveillance for tuberculosis using electronic health record data from an ambulatory practice in eastern Massachusetts. *Public Health Rep.* 2010;125:843–850.

15. Hripcsak G, Soulakis ND, Li L, et al. Syndromic surveillance using ambulatory electronic health records. *J Am Med Inform Assoc.* 2009;16:354–361.

16. International Society for Disease Surveillance (ISDS). Final Recommendation: Core Processes and EHR Requirements for Public Health Syndromic Surveillance. Available at: http://www.syndromic.org/uploads/files/ISDSRecommendation_FINAL.pdf. Published 2011. Accessed October 2, 2012.

17. Khurshid A, Diana ML, Luce SD. Health information exchange: metrics to address quality of care and return on investment. *Perspect Health Inf Manag.* 2012;9:1e.

18. Pevnick JM, Claver M, Dobalian A, et al. Provider stakeholders' perceived benefit from a nascent health information exchange: a qualitative analysis. *J Med Syst.* 2012;36:601–613.

19. Shapiro JS. Evaluating public health uses of health information exchange. *J Biomed Inform.* 2007;40:S46–S49.

20. Howard J, Clark EC, Friedman A, et al. Electronic health record impact on work burden in small, unaffiliated, community-based primary care practices. *J Gen Intern Med.* 2013;26:107–113.

21. Wiljer D, Urowitz S, Apatu E, et al.; Canadian Committee for Patient Accessible Health Records. Patient accessible electronic health records: exploring recommendations for successful implementation strategies. *J Med Internet Res.* 2008;10:e34.

22. National Center for Farmworker Health (NCFH). Available at: <http://www.ncfh.org>. Published 2011. Accessed October 5, 2012.

23. Garcia D, Hopewell J, Liebman AK, Mountain K. The Migrant Clinicians Network: connecting practice to need and patients to care. *J Agromedicine.* 2012;17:5–14.

24. Cooper SP, Shipp EM, Felknor SA, et al. *Community-Academic Linkages to Build a National Network for Farmworker Research.* Bethesda, MD: National Institute of Environmental Health Sciences (NIEHS); 2010–2013. RFA-OD-09-010(999,999).

25. Salud Family Health Centers. Available at: <http://www.saludclinic.org/>. Published 2009. Accessed January 2, 2012.

26. About Health Level Seven (HL7) International. Available at: <http://www.hl7.org/about/index.cfm?ref=footer>. Published 2012. Accessed December 8, 2012.

27. StataCorp. *Stata Statistical Software.* College Station, TX: StataCorp; 2009:11.

28. National High Blood Pressure Education Program. *The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure.* Bethesda, MD: National Heart, Lung, and Blood Institute; August 2004. Report No.: 04-5230.

29. National Center for Farmworker Health (NCFH). Occupational health findings from the National Agricultural Worker Survey (NAWS). In: *National Center for Farmworker Health 22nd Annual Midwest Stream Forum*; November 8, 2012; Crowne Plaza. San Antonio, TX: National Center for Farmworker Health (NCFH); 2012.

30. Emmi KE, Jurkowski JM, Codru N, Bell EM, Kacica MA, Carter TP. Assessing the health of migrant and seasonal farmworkers in New York State: statewide data 2003–2005. *J Health Care Poor Underserved.* 2010;21:448–463.

31. Hansen E, Donohoe M. Health issues of migrant and seasonal farmworkers. *J Health Care Poor Underserved.* 2003;14:153–164.

32. Dever A. *Migrant Health Status: Profile of a Population With Complex Health Problems.* Migrant Clinicians Network Monograph Series. Austin, TX: Migrant Clinicians Network; 1991.

33. Villarejo D, McCurdy SA, Bade B, Samuels S, Lighthall D, Williams D 3rd. The health of California's immigrant hired farmworkers. *Am J Ind Med.* 2010;53:387–397.

34. Anthony M, Williams JM, Avery AM. Health needs of migrant and seasonal farmworkers. *J Community Health Nurs.* 2008;25:153–160.

35. Ostchega Y, Yoon SS, Hughes J, Louis T. Hypertension awareness, treatment, and control—continued disparities in adults: United States, 2005–2006. *NCHS Data Brief.* 2008;(3):1–8.

36. Rodríguez AAA, Rodilla JMR, Delclos J. Documenting occupation in the medical chart of a public hospital system: 2006–2010. *Arch Prev Riesgos Labor.* 2012;15:124–8.

37. Ditolvi Vera G, Benavides Márquez F, Armengol O, Barrionuevo-Rosas L. Cumplimentación de la ocupación en las historias clínicas de la Atención Primaria: 1992–2007. *Aten Primaria*. 2010;42: 486–487.
38. Szklo M, Nieto FJ, eds. *Epidemiology: Beyond the Basics*. 2nd ed. Sudbury, MA: Jones and Bartlett Publishers; 2007.
39. Buntin MB, Burke MF, Hoaglin MC, Blumenthal D. The benefits of health information technology: a review of the recent literature shows predominantly positive results. *Health Aff (Millwood)*. 2011;30:464–471.
40. US Department of Labor. National Agricultural Workers Survey (NAWS). Available at: <http://www.doleta.gov/agworker/naws.cfm>. Published 2011. Accessed December 1, 2012.