



## Associations between urinary biomarkers of polycyclic aromatic hydrocarbon exposure and reproductive function during menstrual cycles in women<sup>☆</sup>



Ulrike Luderer<sup>a,e,f,\*</sup>, Fletcher Christensen<sup>b</sup>, Wesley O. Johnson<sup>b</sup>, Jianwen She<sup>c,1</sup>, Ho Sai Simon Ip<sup>c</sup>, Junqiang Zhou<sup>c</sup>, Josephine Alvaran<sup>c</sup>, Edward F. Krieg Jr<sup>d</sup>, James S. Kesner<sup>d</sup>

<sup>a</sup> Center for Occupational and Environmental Health and Division of Occupational and Environmental Medicine, Department of Medicine, University of California Irvine, 100 Theory, Suite 100, Irvine, CA 92617, USA

<sup>b</sup> Department of Statistics, University of California Irvine, Irvine, CA 92697, USA

<sup>c</sup> Environmental Health Laboratory Branch, California Department of Public Health, Richmond, CA, USA

<sup>d</sup> Division of Applied Research and Technology, National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention, Public Health Service, US Department of Health and Human Services, 4676 Columbia Parkway, Cincinnati, OH 45226, USA

<sup>e</sup> Dept of Developmental and Cell Biology, University of California Irvine, Irvine, CA 92697, USA

<sup>f</sup> Program in Public Health, University of California Irvine, Irvine, CA 92697, USA

### ARTICLE INFO

#### Article history:

Received 26 August 2016

Received in revised form 1 December 2016

Accepted 21 December 2016

Available online 5 January 2017

#### Keywords:

Polycyclic aromatic hydrocarbon

Ovarian function

Reproduction

Luteinizing hormone

Estradiol

Menstrual cycle

### ABSTRACT

Essentially all women are exposed to polycyclic aromatic hydrocarbons (PAHs), formed during incomplete combustion of organic materials, including fossil fuels, wood, foods, and tobacco. PAHs are ovarian toxicants in rodents, and cigarette smoking is associated with reproductive abnormalities in women. Biomonitoring of hydroxylated PAH (OH-PAH) metabolites in urine provides an integrated measure of exposure to PAHs via multiple routes and has been used to characterize exposure to PAHs in humans. We hypothesized that concentrations of OH-PAHs in urine are associated with reproductive function in women. We recruited women 18–44 years old, living in Orange County, California to conduct daily measurement of urinary luteinizing hormone (LH) and estrone 3-glucuronide (E<sub>1</sub>3G) using a microelectronic fertility monitor for multiple menstrual cycles; these data were used to calculate endocrine endpoints. Participants also collected urine samples on cycle day 10 for measurement of nine OH-PAHs. Models were constructed for eight endpoints using a Bayesian mixed modeling approach with subject-specific random effects allowing each participant to act as a baseline for her set of measurements. We observed associations between individual OH-PAH concentrations and follicular phase length, follicular phase LH and E<sub>1</sub>3G concentrations, preovulatory LH surge concentrations, and periovulatory E<sub>1</sub>3G slope and concentration. We have demonstrated the feasibility of using urinary reproductive hormone data obtained via fertility monitors to calculate endocrine endpoints for epidemiological studies of ovarian function during multiple menstrual cycles. The results show that environmental exposure to PAHs is associated with changes in endocrine markers of ovarian function in women in a PAH-specific manner.

© 2016 Elsevier Ltd. All rights reserved.

<sup>☆</sup> The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the National Institute for Occupational Safety and Health (NIOSH). Mention of any company or product does not constitute endorsement by NIOSH.

\* Corresponding author at: COEH, 100 Theory Drive, Suite 100, Irvine, CA 92617, USA.

E-mail addresses: [uluderer@uci.edu](mailto:uluderer@uci.edu) (U. Luderer), [fchrste@uci.edu](mailto:fchrste@uci.edu) (F. Christensen), [wjohnson@ics.uci.edu](mailto:wjohnson@ics.uci.edu) (W.O. Johnson), [Jianwen.She@cdph.ca.gov](mailto:Jianwen.She@cdph.ca.gov) (J. She), [erk3@cdc.gov](mailto:erk3@cdc.gov) (E.F. Krieg), [jsk4@cdc.gov](mailto:jsk4@cdc.gov) (J.S. Kesner).

<sup>1</sup> This project was a collaboration with the California Environmental Contaminant Biomonitoring Program. The conclusions presented do not necessarily reflect or represent the official views or policies of the California Department of Public Health or the State of California.

### 1. Introduction

Polycyclic aromatic hydrocarbons (PAHs) are ubiquitous environmental pollutants formed during incomplete combustion of organic materials such as wood, tobacco, fossil fuels, and food (ATSDR, 1995). Data from the National Health and Nutrition Examination Survey (NHANES) on concentrations of hydroxylated PAHs (OH-PAHs) in the urine of representative samples of Americans show that essentially all Americans are exposed to PAHs (Li et al., 2008; NHANES, 2009). For non-smokers who do not consume grilled or roasted foods, air pollution is the largest source of exposure. Residents of urban areas have higher inhalation exposure to PAHs than do residents of rural areas (Menzie et al., 1992).

Exposure to tobacco smoke can be the principal source of PAH exposure for smokers. Sidestream smoke has 10-fold higher concentrations

of PAHs than mainstream tobacco smoke, and second hand smoke is another major source of exposure (Lodovici et al., 2004; Lu et al., 2008; Shopland et al., 2001). PAHs have been measured in human serum and ovarian follicular fluid (Neal et al., 2008). The latter study showed that follicular fluid of women smokers had significantly elevated levels of the PAH benzo[*a*]pyrene (BaP) compared to follicular fluid of non-smoking women.

Many PAHs are mutagenic and carcinogenic (ATSDR, 1995; IARC, 1983, 2010) and are potent ovarian toxicants and ovarian tumorigens in rodents. Both single high doses (Mattison, 1979; Mattison and Nightingale, 1982; Mattison and Thorgeirsson, 1979; Mattison et al., 1980; Takizawa et al., 1984) and multiple lower doses (Borman et al., 2000) of BaP, 9,12-dimethyl-1,2-benzanthracene (DMBA), and 3-methylcholanthrene destroy immature primordial and primary follicles in mice and rats. DMBA concentration-dependently increases reactive oxygen species and induces apoptosis in cultured rat antral follicles (Tsai-Turton et al., 2007). BaP inhibits growth, survival, and estradiol and anti-Müllerian hormone secretion of cultured mouse secondary follicles (Neal et al., 2007; Sadeu and Foster, 2011).

Epidemiological studies have linked smoking to decreased fecundability (probability of pregnancy) (Alderete et al., 1995; Baird and Wilcox, 1985; Hassan and Killick, 2004; Jensen et al., 1998) and earlier onset of menopause (Harlow and Signorello, 2000; Mattison et al., 1989; van Noord et al., 1997). In a prospective study in which urinary estrogen and progesterone metabolites were measured daily during multiple menstrual cycles, exposure to environmental tobacco smoke in nonsmoking women was associated with lower concentrations of estrone conjugates during the 20-day window around ovulation (Chen et al., 2005).

PAHs are readily absorbed through the lungs, gut, and skin (ATSDR, 1995). PAHs generally require metabolic activation to exert toxicity. They are first oxidized by cytochrome P450 (CYP) enzymes to epoxides, which undergo hydrolysis by epoxide hydrolase to diols, which can be oxidized by CYPs to diol epoxides or converted to catechols by aldo-keto reductases (Penning, 2004; Verma et al., 2012; Xue and Warshawsky, 2005). The diol epoxide metabolites are DNA damaging mutagens, and the catechols undergo redox cycling, generating reactive oxygen species that can damage cellular macromolecules (Denissenko et al., 1996; Penning, 2004; Verma et al., 2012; Xue and Warshawsky, 2005). The epoxides can also spontaneously rearrange to monohydroxylated intermediates, which are excreted as conjugates in the urine or bile (Verma et al., 2012). Biomonitoring of these hydroxylated PAH metabolites in urine provides an integrated measure of PAH exposure via multiple exposure routes (Li et al., 2008).

Although the altered reproductive function observed in women who smoke or who are exposed to environmental tobacco smoke has been postulated to be due to exposure to PAHs in tobacco smoke, no studies have directly examined the associations between PAH exposure biomarkers and measures of hypothalamic-pituitary-ovarian axis function in women. We hypothesized that PAH exposure causes ovarian dysfunction manifested as altered urinary luteinizing hormone (LH) and estrogen metabolite profiles and even anovulatory menstrual cycles.

## 2. Methods

### 2.1. Study participants

Study participants were recruited for a pilot study to lay the groundwork for a subsequent larger, adjunct study to the planned National Children's Study (NCS). The purpose of the pilot study was to test the feasibility of recruiting women who were not intending to become pregnant and not using hormonal contraception for a study of the association between urinary PAH metabolites measured once per menstrual cycle and urinary reproductive hormone concentrations measured daily using a microelectronic dipstick monitor for six menstrual cycles.

Initially, women were recruited based on NCS Vanguard Study protocols (Baker et al., 2014; Montaquila et al., 2010). For the Vanguard Study, Orange County, California was divided into 15 geographical strata, which were each then further divided into 11 geographical units (GUs). Each GU was then divided into 10 segments or 'neighborhoods'. In a two-stage sampling process, 15 of those segments, totaling 15,000 households, were sampled by the NCS Coordinating Center for screening. The remaining segments were not further studied under the NCS Vanguard Study. Women for the current study were recruited from four segments that were not selected for the NCS Vanguard Study. Two segments from predominantly non-Hispanic white GUs (in Irvine, CA) and two segments from predominantly Hispanic GUs (in Santa Ana and Costa Mesa, CA) were selected to assure representation of the two largest ethnic groups in Orange County. Initially eligible women were identified and recruited by door-to-door contact in the home, with follow-up telephone and email contact by study staff. Subsequently, when the NCS shifted to other recruitment strategies, eligible women for the present study were recruited at public events such as health fairs at universities and colleges, work places and events sponsored by community groups. The current study population is thus a convenience sample. Baseline study visits and sample collection occurred between October 2010 and July 2012.

### 2.2. Inclusion and exclusion criteria

Eligible women were between the ages of 18–44, residing in Orange County California, who were not pregnant, currently not planning to conceive, not using hormonal contraception, did not have a history of surgical sterilization, treatment with antineoplastic drugs or radiation therapy to the pelvis and did not have conditions known to cause infertility by mechanisms other than ovarian failure (pelvic inflammatory disease, endometriosis). Women who had recently been pregnant or breastfeeding were asked to delay starting the study until they had one full menstrual cycle after the birth if not breastfeeding or after they stopped breastfeeding.

### 2.3. Participation and compensation

Recruits were told that they would be responsible for testing their first morning void urine daily for reproductive hormones using a dipstick fertility monitor, collecting one urine sample during each of the study menstrual cycles, completing a daily diary and a monthly questionnaire, and potentially taking one or more home pregnancy tests. Participants were given the option of keeping the fertility monitors upon completion of the study or of receiving a payment of \$100 upon returning the monitor. Written, informed consent was obtained from all participants in person by a member of the study staff. All participants were fluent in English or Spanish. The study was approved by the University of California Irvine Institutional Review Board (Study Number 2009-7034).

### 2.4. Baseline study visit

After completing informed consent, participants were instructed to go to one of the two Orange County locations of the UC Irvine Institute for Clinical and Translational Science (ICTS) for their baseline visit 5–9 days prior to their next menses onset. Study staff administered the standardized NCS preconception questionnaire (NCS, 2007) to obtain information about the participant's demographics, medical history, reproductive history, tobacco smoke exposure, exercise history, occupational history, residential history, housing characteristics, use of chemicals (e.g. cleaning agents, pesticides) in and around the home and yard, and pets. Questions relating to tobacco, alcohol, and illicit drug use were taken from the NCS First Trimester Maternal In-Person Questionnaire (NCS, 2007). We refer to these variables hereafter as baseline covariates.

Participants were instructed to fill out a daily diary during the study period to provide information about days and amount of menstrual bleeding (guided by a menstrual pictogram) (Wyatt et al., 2001), contraception use, number of cigarettes smoked, illness, use of medications, number of alcoholic drinks, and sexual intercourse.

ICTS nursing staff measured height, weight, and blood pressure and collected a blood sample. Study staff gave participants urinary hormone monitors, urine collection kits, and home pregnancy tests and provided instruction in how to use them.

### 2.5. Urinary reproductive endocrine testing

Participants were given a Clearblue Easy Fertility Monitor (Swiss Precision Diagnostics, Bedford, UK) and instructed to perform daily urinary dipstick tests to measure estrone 3-glucuronide (E<sub>1</sub>3G) and LH beginning on the first day of their next menstrual cycle. The Clearblue Easy Fertility Monitor measures daily urinary E<sub>1</sub>3G and LH without any need for collection or storage of urine samples. For each menstrual cycle, the participant pushes the monitor “M button” if she had onset of menstrual bleeding during the preceding 24 h (cycle day one). The time of day that she pushes the “M” button becomes the center of the 6 h testing window for subsequent days of that cycle. If the monitor is turned on outside the testing window or is not turned on at all on a given day, no test is requested and the result is recorded as a “fail”. The monitors request daily test sticks starting on cycle day 6 and continuing until an LH surge has been identified or until 20 days have passed, whichever is first. In subsequent cycles, the monitor requests tests based on the timing of the LH surges in prior cycles. To conduct a test, the participant holds a disposable test stick in the urine stream of the first morning void for 3 s. The test stick is then inserted into the monitor to measure LH using a two-site non-competitive immunoassay and E<sub>1</sub>3G by a 1-site competitive immunoassay (Robinson et al., 2007). The monitor displays a fertility status (low, high, or peak), derived by proprietary algorithmic interpretations of the LH and E<sub>1</sub>3G concentrations. The monitor readings do not include any adjustments for urine concentration.

The monitors are capable of storing up to two months of data, including date and time the monitor was turned on, days a test stick was requested and completed, and LH and E<sub>1</sub>3G readings. Study staff downloaded data from the monitors to a data card and then to a computer file, which was sent to the manufacturer for conversion to unitless LH and E<sub>1</sub>3G values.

Study staff visited participants at 1–2 month intervals to collect each participant's monthly urine samples and download data from the monitor to data cards. Participants were also instructed to perform a human chorionic gonadotropin Clearblue Easy Pregnancy Test (Swiss Precision Diagnostics) if their menses onset did not begin within 10 days of their expected date. If the pregnancy test was positive, the participants were removed from the current study.

### 2.6. Calibration of monitor LH and E<sub>1</sub>3G values

Standards were created to calibrate the Clearblue Easy Fertility Monitor LH and E<sub>1</sub>3G unitless values sent by the manufacturer. A stock solution of WHO 2nd International Standard for Pituitary LH for Immunoassay (80/552) with a concentration of 3500 mIU/mL in water was serially diluted in a urine pool containing 1 mIU LH/mL to generate standards with calculated concentrations of 0.4, 1.2, 1.7, 3.7, 11.9, 44.7, 175.9, and 700.8 mIU LH/mL. E<sub>1</sub>3G purified by recrystallization (Kesner et al., 1992) was used to generate E<sub>1</sub>3G standards: 0, 1.6, 3.1, 12.5, 50, 200 ng E<sub>1</sub>3G/mL. These LH and E<sub>1</sub>3G standards were submitted to the Clearblue Easy Fertility Monitors and dipsticks in triplicate (LH) and duplicate (E<sub>1</sub>3G) using three and two separate Clearblue Easy monitors, respectively. The means of the monitor readings were used to construct LH and E<sub>1</sub>3G standard curves (Supplemental Figure S1).

The monitor LH and E<sub>1</sub>3G values were calibrated using the standard curves. In consultation with the manufacturer on the bases of monitor

readings, rules were established to deal with LH and E<sub>1</sub>3G readings that were off the low or high ends of the standard curves or “0” readings. If the LH and E<sub>1</sub>3G readings for a single sample were both “0,” the test was considered flawed and set to missing. If LH reading was “0” or off the low end of the standard curve, the reading was accepted as low and set to half the lowest LH standard level. If an E<sub>1</sub>3G reading was off the low end of the standard curve the reading was accepted as low and set to half the lower limit of the lowest E<sub>1</sub>3G standard level. E<sub>1</sub>3G readings of “0” or > 100 ng/mL were considered flawed and set to missing. If an LH peak reading coincided with an E<sub>1</sub>3G level > 100 ng/mL, the E<sub>1</sub>3G value was set to missing and LH was evaluated by a panel of four experts on a case-by-case basis to determine whether the LH peak was a valid peak based on the details of that cycle and the participant's other cycles.

### 2.7. Calculation of summary endocrine endpoints

Algorithms for menstrual cycle function endpoints were modified from those previously described (Luderer et al., 2013) due to the limited sampling window in this study. Endpoints describing the cycle and phase lengths and LH and E<sub>1</sub>3G secretion patterns are defined in Table 1. LH Surge Onset and Day of E<sub>1</sub>3G Peak were used only to calculate other endpoints and were not further analyzed. Menses onset for each cycle was determined from the monitor output and/or the daily diary. Analysis indicated that almost all quantitative endocrine endpoints had approximately normal distributions, with the sole exception of menstrual cycle length, which was heavily right-skewed.

### 2.8. Measurement of urinary hydroxylated PAH metabolites

Participants were instructed to collect one urine sample on the 10th day after menses onset into a polypropylene beaker and to pour 10 mL of that sample from the beaker into each of 4 cryogenic polypropylene

**Table 1**  
Endocrine endpoint algorithms.

Endpoint (abbreviation)	Algorithm
Cycle length	Number of days from first day of menstrual bleeding through the day before next onset of menstrual bleeding.
Follicular phase length	Number of days from first day of menstrual bleeding to the day of the LH surge onset.
Luteal phase length	Cycle length minus follicular phase length.
LH surge onset	Day of the first LH rise $\geq 2.5$ -fold above the mean of the previous 3–7 day LH baseline.
Highest LH surge level (Highest LH)	Highest LH value of the cycle that is $\geq 2.5$ -fold above the LH surge onset baseline and $\leq 4$ days after the LH surge onset.
Peak LH surge level (Peak LH)	Same as highest LH surge level except it is omitted if there is a missing value on an adjacent day.
Follicular LH level (Follicular LH)	Mean for all days before LH surge onset, or cycle days 6 through 10 for cycles without an LH surge. Omit if $< 3$ values.
Ovulatory status	Ovulatory cycle = Has a defined LH surge onset; Anovulatory cycle = No LH surge onset for cycles with no missing LH values from cycle day 11 through the 9th day before the next menses onset. Must have start and end menses Indeterminate cycle = Cycles that are neither ovulatory nor anovulatory
Mid-follicular E <sub>1</sub> 3G level (Follicular E <sub>1</sub> 3G)	Mean for cycle day 1 through day $-2$ from LH surge onset, or for E <sub>1</sub> 3G values on days 6 through 10 for cycles with no LH surge onset. Omit if $< 3$ values.
Periovulatory E <sub>1</sub> 3G level (Periovulatory E <sub>1</sub> 3G)	Mean for 7 days centered around the day of the LH surge onset. Omit if $< 3$ values.
Preovulatory E <sub>1</sub> 3G rise (E <sub>1</sub> 3G slope)	Slope for 3 days prior to day of E <sub>1</sub> 3G peak. Omit if the first or third value is missing.
Day of E <sub>1</sub> 3G peak	Day of the highest E <sub>1</sub> 3G value $\pm 3$ days from the day of the LH surge onset. For cycles with no surge, select highest cycle E <sub>1</sub> 3G value of the cycle.

vials to be stored in their home freezer until picked up by study staff every month. Hydroxylated PAH (OH-PAH) metabolites were measured in the urine samples as biomarkers of PAH exposure. We chose cycle day 10 because it is approximately in the middle of the follicular phase days when the monitor requests test sticks.

Urinary OH-PAH metabolites were measured for each subject for two cycles (3 participants) or three cycles (48 participants) during the testing period by the California Department of Public Health Environmental Health Laboratory using procedures developed by the Centers for Disease Control and Prevention NHANES study (Li et al., 2008; Romanoff et al., 2006). If a participant only had collected two or three cycles of urinary hormone data, then OH-PAHs were measured in all cycles. If they had collected urinary hormone data for more than three cycles, then the first three cycles with the fewest missing days of urinary hormone data were selected for OH-PAH measurements using isotope dilution gas chromatography/high resolution mass spectrometry (GC-HR-MS). Urine samples were first enzymatically deconjugated and subjected to liquid phase extraction and silylation by *N*-Methyl-*N*-trimethylsilyltrifluoroacetamide (MSTFA), analyzed by GC-HR-MS for nine PAH metabolites, 1- and 2-hydroxy naphthalene (1-NAP, 2-NAP), 1-hydroxy pyrene (1-PYR), 1-, 2-, and 3-hydroxy phenanthrene (1-PHEN, 2-PHEN, 3-PHEN), and 2-, 3-, and 9-hydroxy fluorene (2-FLUO, 3-FLUO, 9-FLUO) (Li et al., 2008; Romanoff et al., 2006). Additional details regarding the OH-PAH measurement methods can be found in Supplemental materials.

A small percentage of analytes were below the limits of detection (LOD) of the assay. These were set to the LOD/ $\sqrt{2}$  (Ogden, 2010).

### 2.9. Statistical analyses

Descriptive statistics (arithmetic means and standard deviations and geometric means for continuous variables and percentages in each group for categorical variables) were calculated for demographic variables (Table 2), endocrine endpoints (Table 3) and OH-PAH concentrations (Table 4). Intra-class correlation coefficients (ICCs) were calculated using mean squares from one-way random effects ANOVA models with participant as the independent variable (Bartko, 1966) to assess the correlation of individual OH-PAH metabolite concentrations among repeated samples within participants. To assess the correlations among the OH-PAH metabolite concentrations within the same urine sample, pairwise Pearson correlations were calculated (Table 5).

Observed metabolite concentrations were right-skewed, but were approximately normal after a log-transformation was applied. These log transformed values were then standardized by subtracting the corresponding sample mean and dividing by the corresponding standard deviation so that each new transformed variable has sample mean zero and sample standard deviation one. Then, in part because metabolites of phenanthrene (1-PHEN, 2-PHEN, 3-PHEN) and of fluorene (2-FLUO, 3-FLUO, 9-FLUO) can be collinear, additional transformations were considered for each. For each woman and time at which these measurements were taken, an overall average concentration was computed for PHEN, and another for FLUO. These variables were multiplied by a constant for technical reasons and were included in models as PHENS and FLUOS. Additional details are provided in Supplemental Methods.

For phenanthrene and fluorene, metabolite-specific variables were calculated as differences between the specified (standardized log) metabolite and the corresponding overall average concentrations (PHENS and FLUOS), for each woman-time combination. Each of these differences was multiplied by a suitable constant for technical reasons. These transformed variables were termed the same as the original variables for simplicity. So the variable we analyzed subsequently termed 1-PHEN involved the difference between the originally standardized log of 1-PHEN minus the average of the corresponding 1-, 2-, and 3-PHENS, for each of the woman-time combinations, and multiplied by a constant. This was all done to make clearer the difference between an

**Table 2**  
Characteristics of study participants.

Characteristic	Mean $\pm$ SD	Range	N
Age at baseline	29.9 $\pm$ 6.5	18, 44	51
Height (m)	1.63 $\pm$ 0.07	1.50, 1.83	48
Weight (kg)	66.6 $\pm$ 15.7	49.2, 124.9	49
BMI	24.9 $\pm$ 5.3	18.6, 42.6	48
Characteristic	Number	Percent	
Ethnicity/Race			
Hispanic	11	21.6	
Non-Hispanic White	22	43.1	
Non-Hispanic Asian	14	27.5	
Non-Hispanic Black	4	7.8	
Education – highest level attained			
High school diploma or less	3	5.9	
Some college or vocational	20	39.2	
Bachelor's degree	16	31.4	
Graduate degree	11	21.6	
Family income/year			
<\$20,000	4	7.8	
$\geq$ \$20,000 < \$50,000	12	23.5	
$\geq$ \$50,000 < \$75,000	10	19.6	
$\geq$ \$75,000 < \$100,000	9	17.6	
$\geq$ \$100,000	9	17.6	
Current smoking			
Yes	6	11.8	
No	45	88.2	
Alcohol			
$\geq$ 2 days/wk	12	23.5	
1–4 days/month	16	31.4	
<1 day/month or never	21	41.2	
Caffeinated beverages (drink regularly)			
Coffee	28	54.9	
Tea	21	41.2	
Soda	10	19.6	
Energy drinks	2	3.9	
Vigorous exercise (minutes in past 7 days)			
0	12	23.5	
>0 $\leq$ 120	22	43.1	
>120 $\leq$ 390	10	19.6	
>390	7	13.7	
Walking (hours in past 7 days)			
$\leq$ 1	9	17.6	
>1 $\leq$ 4	21	41.2	
>4 $\leq$ 7	8	15.7	
>7	13	25.5	

Percentages may not total to 100 because of missing data or because groups not mutually exclusive.

overall effect for a compound like phenanthrene and the effect for having unusually high (or low) concentration of one metabolite relative to the average concentration of phenanthrene metabolites observed in the individual.

The two naphthalene metabolites were not combined. 1-NAP, in addition to being a metabolite of naphthalene, is generated by metabolism of the pesticides carbaryl and napropamide and the beta-blocking medication propranolol (Hill et al., 1995; Shealy et al., 1997; Walle and Gaffney, 1972). 2-NAP, in addition to being a metabolite of naphthalene, is also a metabolite of naphthoanilide and (2-naphthylxy)acetic acid (Hill et al., 1995).

To investigate the role of OH-PAH concentrations in predicting the aforementioned endocrine endpoints, we performed a two-stage procedure. Models for all endpoints were constructed in the same way—through a Bayesian mixed modeling approach with subject-specific random effects allowing each participant to act as a baseline for her set of measurements. In the first stage we built a model involving only non-PAH baseline covariates, using a backwards stepwise algorithm to select a parsimonious model that fits the data. The starting baseline model included the covariates age, race, educational attainment, stress, body-mass index (BMI), alcohol use, caffeine intake, and measures of how many minutes the participant walked and engaged

**Table 3**

Means, standard deviations, and geometric means of menstrual cycle endpoint variables averaged over repeated measures within participant.

Endpoint variable	Mean (SD)	Geometric mean	N*	n**
Cycle length (days)	33.2 (10.7)	32.0	51	297
Follicular phase length (days)	16.3 (3.3)	16.0	47	199
Luteal phase length (days)	14.3 (5.3)	13.6	47	193
Mean follicular phase E <sub>1</sub> 3G (ng/mL)	20.2 (7.5)	18.7	50	219
Mean follicular phase LH (mIU/mL)	9.1 (2.0)	8.9	50	233
Preovulatory E <sub>1</sub> 3G rise (slope)	3.2 (3.7)	3.2	49	178
Periovulatory E <sub>1</sub> 3G (ng/mL)	31.1 (10.3)	29.4	47	191
Peak LH (mIU/mL)	37.9 (12.1)	36.1	45	132
Highest LH (mIU/mL)	36.6 (10.7)	35.3	47	199
Ovulatory status of cycles				N (%)
Ovulatory				199 (67.0)
Anovulatory				12 (4.0)
Indeterminate				86 (29.0)

Generated using all cycles available for each woman, regardless of whether that cycle had OH-PAH measurements. Calculated by taking the mean for each participant for each endpoint, then calculating the means, SDs, and geometric means of those new variables. \*N refers to number of women for whom there was at least one cycle of data for that variable. \*\*n refers to the total number of cycles for which each endpoint could be calculated, out of the 305 cycles observed.

in vigorous physical activity each week. Estimates of regression coefficients are their posterior means. These were numerically approximated using Markov Chain Monte Carlo (MCMC) simulation (Christensen et al., 2011), Chapter 6). This allowed us to directly examine the posterior probability that each covariate in the model was associated with a non-zero regression coefficient by determining the proportion of simulated coefficient values that were above zero.

The second stage model then began from the selected baseline covariate model, adding all of the OH-PAH variables and an indicator variable denoting whether the participant was a smoker to that model. Because OH-PAH information was only available for 2–3 cycles per participant, we used a shared-parameter modeling strategy to take full advantage of our available endpoint data: mean-centering the OH-PAH covariates, fitting models for the cycles without PAH information using only baseline covariates, fitting models for the cycles with PAH information using both baseline and OH-PAH covariates, and constraining the baseline model coefficients to be equal between models for the same endpoint. As in the first stage, we used a backwards selection procedure to again pick a parsimonious model that fit the data well with any additional terms that had been selected. Our primary research interest was to assess the effect of PAHs on endocrine endpoints, and so our goal in these analyses was to determine whether OH-PAHs could provide any incremental benefit in modeling those endpoints beyond the fit we obtained with the baseline model.

Finally, we compared the deviance information criterion (DIC, a common Bayesian measure of model fit) for the baseline and final models. DIC for these models was calculated manually from the MCMC iterates to ensure that only covariates of interest were considered in assessing model complexity. Generally, a decrease of 3 or more

in DIC is suggestive that a lower-scoring model is preferred to a higher-scoring model (Spiegelhalter et al., 2002). In this way, we addressed a primary goal of this paper by assessing the added benefit of OH-PAH terms over and above the model involving only baseline characteristics. Analyses were performed using R and WinBUGS, two publicly available statistical analysis programs. Additional details about the statistical methods are provided in Supplemental Materials.

### 3. Results

#### 3.1. Demographics and participant characteristics

Three participants were eliminated from the analyses because they did not collect any urine for PAH measurements (one woman) or they had performed so few urine hormone test sticks that the no endocrine endpoints could be calculated. Demographic and other characteristics of the 51 remaining participants are shown in Table 2. The mean age of the participants was 29.9 years. Non-Hispanic white women made up the largest racial/ethnic group among the participants, followed by Asian and Hispanic, in that order. More than 92% of the participants had some education beyond high school graduation, with 21.6% having a graduate degree. The mean BMI was 24.9. More than a third of the participants had engaged in more than 2 h of vigorous exercise, while nearly 24% had not engaged in any vigorous exercise, during the week prior to the baseline interview. Forty-one percent of the participants reported walking >4 h during the week prior to the baseline interview. Fewer than 12% of the participants smoked, and 23.5% reported drinking alcoholic beverages on 2 or more days per week.

#### 3.2. Menstrual cycle endpoints

We had monitor data for 305 menstrual cycles from the 51 participants. Of these, 150 cycles had OH-PAH measurements. Representative menstrual cycle LH and E<sub>1</sub>3G concentrations for a participant with regular cycles with clearly defined LH peaks are shown in Fig. 1A, while representative data for a participant with some anovulatory cycles are shown in Fig. 1B. Both of these participants missed very few days of sampling. In contrast, some participants had many missing days. Of 4726 potential test days recorded by the monitors, tests were not performed on 1168 days. Most of these were because the participant did not turn the monitor on at all that day or turned it on outside of the test window; 10 participants had one to several cycles when they took a hiatus for travel or other reasons and resumed thereafter.

We applied the algorithms summarized in Table 1 to the urinary LH and E<sub>1</sub>3G concentrations and menses onset data to generate endocrine endpoints for each cycle. These endocrine endpoints are summarized

**Table 4**

Means, standard deviations, and geometric means of OH-PAH concentrations.

OH-PAH	Unadjusted concentration (pg/mL)			Creatinine adjusted concentration (ng/g creat)	
	Mean (SD)	Geometric mean	LOD	Mean (SD)	Geometric mean
1-NAP	2252 (2005)	1632	<25	1785 (1245)	1387
2-NAP	8738 (10085)	5797	<20	6192 (5037)	4965
1-PYR	142 (124)	110	<20	107 (64)	95
1-PHEN	150 (89)	125	<10	119 (60)	107
2-PHEN	41 (22)	35	<10	33 (14)	31
3-PHEN	75 (42)	64	<10	59 (23)	55
2-FLUO	272 (204)	213	<20	209 (121)	184
3-FLUO	140 (139)	105	<20	106 (64)	91
9-FLUO	361 (316)	278	<37	296 (259)	240

Calculated by taking the mean concentration for each participant for each OH-PAH, then calculating the means, SDs, and geometric means of those new variables.

**Table 5**  
Correlations among concurrent measurements of OH-PAHs<sup>a</sup>.

	1-NAP	2-NAP	1-PYR	1-PHEN	2-PHEN	3-PHEN	2-FLUO	3-FLUO	9-FLUO
1-NAP	1	0.105	0.234	0.429	0.285	0.352	0.374	0.446	0.424
2-NAP		1	0.163	-0.038	0.030	-0.011	0.072	0.067	-0.113
1-PYR			1	0.597	0.574	0.654	0.385	0.465	0.379
1-PHEN				1	0.738	0.738	0.635	0.565	0.655
2-PHEN					1	0.764	0.573	0.536	0.615
3-PHEN						1	0.638	0.683	0.631
2-FLUO							1	0.887	0.470
3-FLUO								1	0.459
9-FLUO									1

<sup>a</sup> Pearson correlations for all 151 urine samples for which OH-PAHs were measured.

in Table 3. This table also shows for how many cycles each endpoint could be calculated. Cycle length was calculable for the largest number of cycles, 297 of 305 cycles. The mean follicular phase LH and E<sub>1</sub>3G concentrations were calculable for the next largest numbers of cycles because these variables do not depend on having identified a mid-cycle LH surge. Because ovulation status was indeterminate for 29% of cycles due to missing urinary LH data during mid-cycle, we had low statistical power to detect effects of OH-PAHs on ovulation and therefore do not present models for this endpoint.


3.3. PAH exposures

OH-PAHs were measured around menstrual cycle day 10 for three cycles (48 participants) or two cycles (3 participants) per participant. Table 4 shows the arithmetic and geometric mean concentrations and LODs of urinary OH-PAH metabolites. All participants had detectable

concentrations of each of the measured OH-PAHs in at least one urine sample, which is consistent with ubiquitous exposure to PAHs. Overall, very few urine samples had concentrations of the measured PAH metabolites below the LOD. 1- and 2-NAP concentrations were above the LOD in all samples assayed. 2-FLUO, 9-FLUO, 1-PHEN, and 3-PHEN were below the LOD in one urine sample each (0.7% of the samples). 1-PYR was below the LOD in 6 samples (4%); 2-PHEN in 12 samples (7.9%) and 3-FLUO in 5 samples (3.3%).

ICCs ranged from 0.23 for 3-PHEN to 0.67 for 2-FLUO (creatinine adjusted), indicating that concentrations of the urinary OH-PAH metabolites were minimally to moderately correlated among repeated samples within participants. This provides justification for analyzing multiple cycles within participants for associations between OH-PAHs and endocrine endpoints.

Pairwise correlations among the different OH-PAHs within urine samples are shown in Table 5. All of the measured metabolites tended



**Fig. 1.** Representative urinary LH and E<sub>1</sub>3G concentrations for two participants across multiple menstrual cycles. A) A participant with regular cycles with clearly defined LH peaks. B) A participant with several ostensibly anovulatory cycles without clear LH peaks. Blue diamonds indicate the onset of menses. Days for which urinary OH-PAHs were measured are indicated by a purple X.

to have moderate correlation with each other, with higher correlations for metabolites of the same parent compound. 2-NAP was the exception; it had relatively low correlation with the other metabolites including 1-NAP. The moderately high collinearity among metabolites of fluorene and among metabolites of phenanthrene demonstrated in Table 5 provides some justification for the combination and difference-score forms we used for these metabolites in the endocrine endpoint analyses.

**Table 6**  
Baseline and PAH models for endocrine endpoints.

Endpoint	DIC (w/o PAHs)	DIC (with PAHs)	
Cycle length	1211.5	1203.3	
Baseline	Race + Age + Walking + Vigorous Activity		
PAHs	9-FLUO	$\beta = -0.507$	$\Pr(\beta < 0)^a = 0.960$
	2-PHEN	$\beta = 0.323$	$\Pr(\beta > 0) = 0.889$
	1-PYR	$\beta = 0.728$	$\Pr(\beta > 0) = 0.994$
Follicular phase length	977.9	961.6	
Baseline	Race + Age + Stress + Vigorous Activity		
PAHs	3-FLUO	$\beta = 1.040$	$\Pr(\beta > 0) = 0.999$
	1-PYR	$\beta = 0.578$	$\Pr(\beta > 0) = 0.982$
Follicular LH	1072.6	1068.3	
Baseline	Race + Age + Caffeine <sup>b,c</sup> + BMI + Alcohol + Vigorous Activity		
PAHs	2-NAP	$\beta = 0.446$	$\Pr(\beta > 0) = 0.962$
	PHENs	$\beta = -0.963$	$\Pr(\beta < 0) = 0.996$
	1-PYR	$\beta = 0.548$	$\Pr(\beta > 0) = 0.945$
Highest LH	1563.2	1557.3	
Baseline	Race + Age + Caffeine <sup>b,c</sup> + BMI <sup>c</sup> + Alcohol <sup>b</sup> + Walking <sup>b</sup> + Education <sup>b</sup>		
PAHs	2-FLUO	$\beta = 2.057$	$\Pr(\beta > 0) = 0.932$
	3-FLUO	$\beta = -2.000$	$\Pr(\beta < 0) = 0.941$
	FLUOs	$\beta = 5.308$	$\Pr(\beta > 0) = 0.983$
	1-NAP	$\beta = -1.977$	$\Pr(\beta < 0) = 0.924$
	2-NAP	$\beta = -3.143$	$\Pr(\beta < 0) = 0.936$
Peak LH	1055.7	1035.8	
Baseline	Race + Age + Caffeine <sup>b,c</sup> + BMI <sup>b,c</sup> + Alcohol <sup>b</sup> + Walking <sup>b</sup> + Education <sup>c</sup>		
PAHs	2-FLUO	$\beta = 2.755$	$\Pr(\beta > 0) = 0.949$
	3-FLUO	$\beta = -2.988$	$\Pr(\beta < 0) = 0.981$
	FLUOs	$\beta = 7.262$	$\Pr(\beta > 0) = 0.992$
	1-NAP	$\beta = -3.735$	$\Pr(\beta < 0) = 0.981$
	2-NAP	$\beta = -1.823$	$\Pr(\beta < 0) = 0.893$
	2-PHEN	$\beta = -2.371$	$\Pr(\beta < 0) = 0.950$
	PHENs	$\beta = -4.280$	$\Pr(\beta < 0) = 0.967$
Endpoint	DIC (w/o PAHs)	DIC (with PAHs)	
Follicular E13G	1428.7	1420.0	
Baseline	Race + Age + Caffeine <sup>b</sup> + Walking		
PAHs	2-FLUO	$\beta = 1.325$	$\Pr(\beta > 0) = 0.979$
	3-FLUO	$\beta = -1.854$	$\Pr(\beta < 0) = 0.999$
	1-NAP	$\beta = 0.830$	$\Pr(\beta > 0) = 0.924$
	2-NAP	$\beta = -0.680$	$\Pr(\beta < 0) = 0.878$
	2-PHEN	$\beta = -0.613$	$\Pr(\beta < 0) = 0.870$
	Smoking	$\beta = 4.975$	$\Pr(\beta < 0) = 0.940$
Periovulatory E13G	1322.9	1318.8	
Baseline	Race + Age + Caffeine + Alcohol + Vigorous Activity + Education		
PAHs	FLUOs	$\beta = 1.595$	$\Pr(\beta > 0) = 0.920$
	2-NAP	$\beta = -1.381$	$\Pr(\beta < 0) = 0.954$
	1-PYR	$\beta = -0.985$	$\Pr(\beta < 0) = 0.853$
	Smoking	$\beta = 8.585$	$\Pr(\beta > 0) = 0.969$
E13G Slope	1097.7	1094.4	
Baseline	Race + Age + Stress <sup>b</sup> + Alcohol <sup>b</sup> + Walking + Vigorous Activity		
PAHs	1-NAP	$\beta = 0.967$	$\Pr(\beta > 0) = 0.954$
	2-NAP	$\beta = 0.815$	$\Pr(\beta > 0) = 0.944$
	1-PHEN	$\beta = -0.952$	$\Pr(\beta < 0) = 0.939$
	Smoking	$\beta = 1.809$	$\Pr(\beta > 0) = 0.878$

<sup>a</sup> Posterior probability that covariate is associated with a non-zero regression coefficient (proportion of simulated coefficient values that were above zero).

<sup>b</sup> These variables were included in the baseline model but dropped in the PAH model under our selection criteria.

<sup>c</sup> Because Highest LH and Peak LH use the same data whenever a peak was detected (see Table 1), we chose to force BMI and caffeine into the Peak LH baseline model to make it agree with the Highest LH baseline model.

#### 3.4. Associations between OH-PAH metabolites and endocrine endpoints


Table 6 presents the final OH-PAH models for each endocrine endpoint, PAH coefficient estimates, the proportion of positive coefficients from the MCMC, a list of baseline covariates identified by the stepwise modeling procedure for which the models have been adjusted, and DIC scores for comparing the baseline and final models. Creatinine-adjusted OH-PAH measurements were used for all models displayed. The models for all endpoints were refit with unadjusted OH-PAH measurements, and the results did not differ in any appreciable respect, and thus are not presented here.

For all eight endpoints considered, addition of OH-PAHs or smoking status improved the model fit (decreased DIC by >3) beyond that provided by the covariates in the baseline models. Smoking status played a role in modeling the three E<sub>1</sub>3G endpoints, but directly measured OH-PAHs were selected into all eight models even when smoking status was accounted for.

Because understanding what these models are saying can be difficult, we provide graphical interpretations using the follicular phase length (Fig. 2) and highest LH (Supplemental Figure S3) endpoints. Let us consider a hypothetical participant with urinary 1-PYR concentration of 85 ng/g Cr for her first menstrual cycle, and 280 ng/g Cr for her second cycle. A 1-PYR concentration of 85 ng/g Cr is approximately the median concentration in our dataset, and a concentration of 280 ng/g Cr reflects a value two standard deviations higher on the log scale. Then the expected increase in follicular phase length is  $2\beta_{1PYR} = 1.156$  days longer in the second cycle than in the first (Fig. 2). In addition, the graphs in Fig. 2 show that a shift in 2-FLUO or 9-FLUO from their median to their 75th percentile concentrations is associated with approximately a half day decrease in follicular phase length, while a similar shift in 3-FLUO is associated with about a one day increase. For a sense of how large these effects are, based on our data we find that given identical covariate values, about half the time a woman's follicular phase length will vary by 3.3 days or less. Similarly, for about half of the participants the woman-to-woman difference in average follicular phase length is about 3.7 days or less. A predicted difference of 1–2 days based on OH-PAH concentration levels is less than the range of expected cycle-to-cycle variability, but it is still a noteworthy effect.

Highest LH and Peak LH were calculated from the same data when a surge was identified with no missing data. Peak LH was not calculated when the peak of the LH surge could not be confirmed due to missing data. Results for both endpoints are similar, as we would hope, and show roles for all naphthalene and fluorene metabolites studied, as well as some naphthalene and phenanthrene metabolites. Of particular note, the coefficients corresponding to 2-FLUO and 3-FLUO have opposite signs and similar magnitudes. Recall that these covariates represent the difference between their specific metabolite and the overall shared level of the fluorene compound across all metabolites. What we appear to see here, then, is that highest LH increases when the 2-FLUO metabolite accounts for a greater proportion of the total fluorene in a subject's urine than the 3-FLUO metabolite. We also see that increasing levels of fluorene overall are associated with an increase in the highest LH level. Supplemental Figure S3 provides a graphical representation of the effects of changes in 1-NAP, 2-FLUO, and 3-FLUO on highest LH. We see from the graphs that highest LH decreases with increasing 1-NAP. A shift in 2-FLUO or 9-FLUO from their medians to their 75th percentile concentrations results in approximately a 5 and 1 mIU/mL increase in highest LH, respectively, while a similar shift in 3-FLUO is associated with about a 3 mIU/mL decrease in highest LH.

For the three follicular phase E<sub>1</sub>3G endpoints, we observed positive associations with smoking. On average, the slope of the estradiol rise is steeper in smokers than the slope in non-smokers. In practical terms, for a white woman of average age (in the study) who doesn't walk or exercise much, the slope for nonsmokers shows an increase of about 2 ng/mL per day, and the slope for shows a daily increase of about 4 ng/mL. E<sub>1</sub>3G slope also increases with 1- and 2-NAP and



**Fig. 2.** Graphical representation of changes in follicular phase length with changes in 1-PYR and 2-, 3-, and 9-FLUO derived from the final model. All four graphs show the changes in follicular phase length with increasing urinary concentration of 1-PYR at two different concentrations of 9-FLUO, the study median concentration (solid line) and the 75th percentile concentration (dotted line). 2-FLUO was held constant at its study median concentration in the two graphs on the left and at its 75th percentile in the two graphs on the right. 3-FLUO was held constant at its study median concentration in the two upper graphs and at its 75th percentile in the two lower graphs. The graphs show that follicular phase length increases with increasing 1-PYR. A shift in 2-FLUO or 9-FLUO from the median to the 75th percentile results in approximately a half day decrease in follicular phase length, while a similar shift in 3-FLUO is associated with a one day increase.

decreases when 1-PHEN is higher than the other phenanthrene metabolites. On average, the periovulatory E<sub>1</sub>3G concentrations are higher for women who smoke by nearly 9 ng/mL. A white woman of average age who doesn't exercise much, doesn't have a college degree, doesn't drink caffeinated beverages, and doesn't drink alcohol will on average have a periovulatory E<sub>1</sub>3G of about 20 ng/mL if she doesn't smoke and 28.5 ng/mL if she does smoke. Periovulatory E<sub>1</sub>3G also increases with fluorene metabolites and decreases with 2-NAP and 1-PYR. On average, follicular phase estradiol is higher for smokers, with our model predicting a 5-unit increase in E<sub>1</sub>3G for smokers compared to non-smokers. A white woman of average age who doesn't walk much would have an average follicular phase E<sub>1</sub>3G of 15 ng/mL if she doesn't smoke, and 20 ng/mL if she does. Follicular E<sub>1</sub>3G also increases with 1-NAP or 2-FLUO that is higher than the other fluorene metabolites and decreases with 2-NAP or 3-FLUO higher than the other fluorene metabolites.

The final model for menstrual cycle length did not include any OH-PAHs when all cycles were included in the models. When only cycles with length within the normative range (21–35 days) were analyzed,

cycle length increased with 1-PYR and 2-PHEN concentrations and decreased with 9-FLUO concentration.

#### 4. Discussion

Animal studies have clearly demonstrated that several higher molecular weight PAHs are potent ovotoxicants that destroy ovarian follicles, causing ovarian failure (Borman et al., 2000; Mattison, 1979; Mattison and Nightingale, 1982; Mattison and Thorgeirsson, 1979; Mattison et al., 1980; Takizawa et al., 1984). Cigarette smoking is associated with decreased fecundity and earlier menopause in women, which may be due to the dozens of PAHs found in tobacco smoke (Alderete et al., 1995; Baird and Wilcox, 1985; Harlow and Signorello, 2000; Hassan and Killick, 2004; Jensen et al., 1998; Mattison et al., 1989; van Noord et al., 1997). However, this is the first study that has directly examined the associations of specific biomarkers of PAH exposure with measures of ovarian function in women. We found overall positive associations of fluorene metabolites and negative associations of naphthalene metabolites with two measures of LH surge amplitude. We

found positive associations of 2-NAP and 1-PYR and negative association of phenanthrene metabolites with average follicular LH. Follicular phase length was positively associated with 3-FLUO and 1-PYR. Smoking was positively associated with follicular and periovulatory E<sub>1</sub>3G concentrations and E<sub>1</sub>3G slope, and PAH metabolites were retained in these models.

A few prior studies have examined the effects of cigarette smoking on similar endpoints. Smoking has been associated with shorter cycle and follicular phase lengths (Windham et al., 1999), increased early follicular phase urinary E<sub>1</sub>3G, non-significantly decreased luteal phase urinary pregnanediol-3-glucuronide, and higher urinary FSH concentrations during the luteal to follicular phase transition (Windham et al., 2005). Similarly, increased early follicular phase serum estradiol and follicular phase progesterone concentrations have been reported in smokers compared to non-smokers (Zumoff et al., 1990). The same study reported decreased follicular phase serum LH concentrations and no differences in peak preovulatory LH concentrations. In contrast, another study reported that exposure to environmental tobacco smoke was associated with decreased mid to late follicular phase urinary E<sub>1</sub>3G concentrations during non-conception cycles only (Chen et al., 2005).

It is difficult to compare the associations between OH-PAH metabolites and endocrine endpoints in the present study with the reported associations between exposure to tobacco smoke and similar endocrine endpoints in prior studies. Active smoking is associated with significantly increased urinary excretion of all nine OH-PAHs measured in the present study, while exposure to environmental tobacco smoke is associated with lesser or no increase in excretion of these metabolites (Aquilina et al., 2010; Suwan-Ampai et al., 2009). Tobacco smoke contains many high molecular weight PAHs, which were not directly biomonitoring in the present study because their metabolites are primarily excreted in the bile and are below the limit of detection in urine (Li et al., 2008). Tobacco smoke also contains thousands of other compounds (Shopland et al., 2001), and therefore observed associations with smoking may or may not be due to the PAH content of tobacco smoke. Smoking was retained in the second stage of modeling along with specific PAH metabolites only for the three E<sub>1</sub>3G endpoints in the present study. This indicates that the OH-PAH metabolites contribute additional information to the models above and beyond that contributed by smoking. We observe increased follicular E<sub>1</sub>3G concentrations with smoking, which is consistent with increased early follicular phase E<sub>1</sub>3G in smokers reported by Windham and coworkers (Windham et al., 2005). We also observed associations of follicular E<sub>1</sub>3G with fluorene, naphthalene, and phenanthrene metabolite concentrations. In contrast, our finding that periovulatory E<sub>1</sub>3G increases with smoking is opposite the findings of Chen et al. regarding the association between environmental tobacco smoke exposure and mid to late follicular phase E<sub>1</sub>3G in non-conception cycles (Chen et al., 2005). We also observed that periovulatory E<sub>1</sub>3G decreases with 2-NAP and 1-PYR and increases with fluorene. While we find that peak and highest LH decreased with naphthalene and phenanthrene and increased with fluorene, we see no effect for smoking on these endpoints, concordant with prior findings of no effect of smoking on LH surge amplitude (Zumoff et al., 1990). Higher preovulatory LH concentrations have been associated with conception cycles compared to nonconception cycles (Baird et al., 1999). While we find that both pyrene and the fluorene metabolite profile associate with follicular phase length, we see no effect for smoking, which contrasts with prior findings associating shorter follicular phase length with smoking (Windham et al., 1999).

For women who do not smoke, food and exposure to particulate matter air pollution are the major sources of PAH exposures. Consumption of grilled meat is associated with greater increases in urinary 1- and 2-NAP than the other PAH metabolites in the NHANES data (Suwan-Ampai et al., 2009). More detailed studies in which OH-PAHs were monitored repeatedly before and after consumption of grilled chicken (Li et al., 2012) or smoked salmon (Motorykin et al., 2015) showed that

urinary levels of all nine OH-PAHs increase rapidly reaching peak concentrations within 8.5 h and decrease rapidly with median half-lives ranging from 1.7 to 7.0 h. The OH-PAHs monitored in the present study have also been measured in relation to exposure to wood smoke. Urinary excretion of all nine increased after exposure, but 2-NAP showed the strongest association with personal air monitoring of wood smoke components, and 1-PYR showed no association (Li et al., 2015). Inhalation exposure to lower molecular weight PAHs, including the parent compounds of the metabolites measured in the present study, is more strongly associated with indoor sources, while higher molecular weight PAH exposures are associated with outdoor air pollution sources (Nethery et al., 2012). Compared to other studies that examined the effects of smoking on reproductive endpoints in women, the measurement of biomarkers that integrate PAH exposures from all routes is a strength of our study, and our models demonstrate that individual OH-PAH metabolites may have divergent associations with the same endpoint.

Our study is the first to convert output from urinary hormone monitors to concentrations and to utilize these data to calculate endocrine endpoints independently of the monitor algorithms. Other studies have demonstrated that use of the Clearblue Easy Fertility Monitors can increase pregnancy rates by identification of the potentially fertile window (Robinson et al., 2007; Tanabe et al., 2001). The monitors, along with daily diaries, have also been used to establish menses dates in a cohort study of the effects of environmental exposures on time to pregnancy (Buck Louis et al., 2011; Buck Louis et al., 2014a; Buck Louis et al., 2012; Buck Louis et al., 2014b). We generated standard curves to convert monitor readings to LH and E<sub>1</sub>3G concentrations and used these to calculate summary endocrine variables for each menstrual cycle. The dynamic range of the monitor LH assay spanned more than two orders of magnitude, while that of the monitor E<sub>1</sub>3G assay spanned about one order of magnitude (Supplemental Figure S1). As a result, there were many more E<sub>1</sub>3G than LH monitor readings above or below the limits of the respective standard curves (282 versus 99). Because the commercially available monitors used for this study are designed to minimize testing days, we only had data for a maximum of 20 days of each cycle, starting no earlier than cycle day 6 and ending soon after the LH surge ended. Therefore, we were not able to calculate any early follicular phase or luteal phase endpoints. In addition, because the monitors do not measure progesterone metabolite concentrations we were not able to confirm that ovulation occurred by identifying a significant luteal phase rise in progesterone or fall in the ratio of E<sub>1</sub>3G to progesterone metabolites (Baird et al., 1995; Baird et al., 1991; Kassam et al., 1996). Finally, the monitor output did not include outputs (i.e., creatinine, osmolality, specific gravity) to adjust for urine dilution when converting the monitor readings to endocrine concentrations.

Our observations of associations of some OH-PAHs with follicular and periovulatory E<sub>1</sub>3G concentrations are consistent with experimental studies that have reported effects of the OH-PAHs measured in the present study or their parent compounds on ovarian steroid synthesis in several species. Dietary phenanthrene exposure dose-dependently decreased plasma estradiol concentrations in female flounder and decreased androstenedione, testosterone, and estradiol synthesis in cultured flounder ovaries via inhibition of the enzymatic activities of key steroidogenic enzymes (Monteiro et al., 2000a, 2000b). 9-PHEN, 1-NAP, 1-PYR all inhibited steroidogenic cytochrome P450 enzyme activities in carp ovary and testis subcellular fractions (Fernandes and Porte, 2013). Phenanthrene exposure also inhibited ovarian development and down-regulated ovarian aromatase, estrogen and androgen receptor expression in another fish species, the marine medaka (Sun et al., 2015). In cultured rat ovarian luteal cells, phenanthrenequinone, a photooxidation product of phenanthrene, inhibited LH-stimulated progesterone secretion and increased reactive oxygen species (Nykamp et al., 2001).

While the different associations we observed between various OH-PAH metabolites and several endocrine endpoints are interesting, we

must be cautious in interpreting these associations because most of the OH-PAHs were highly correlated with one another. Only 2-NAP was not highly correlated with the other OH-PAHs. Interestingly, 2-NAP was a significant predictor for five of the eight endocrine endpoints. For the other OH-PAHs retained in seven of the nine models, their combined effects may be considered to be representative of the combined effects of all of the highly correlated OH-PAHs. In addition, because we examined a large number of endpoints and PAH metabolites, some of the associations may be spurious and should be confirmed in larger studies. It is also possible that associations were missed because we were only able to measure biomarkers of PAH exposure once per cycle during the follicular phase, reflecting exposures during the preceding day to a few days. In the future, it would be ideal to conduct a similar study in which PAH metabolites are measured more frequently during each cycle.

Another limitation of the present study is that 29% of the cycles were indeterminate for ovulatory status due to missed testing days. As a result, we had reduced power to detect effects of OH-PAHs on ovulatory status. Missed test days also decreased the number of cycles for which some of the other endpoints could be calculated. A larger study will be required to confirm the associations between OH-PAHs and endocrine endpoints we observed.

In summary, we have demonstrated the feasibility of using urinary reproductive hormone data obtained via microelectronic fertility monitors to calculate endocrine endpoints for epidemiological studies of ovarian function during multiple menstrual cycles. We observed associations between biomarkers of environmental PAH exposure and follicular phase length, follicular phase LH and E<sub>1</sub>3G concentrations, preovulatory LH surge amplitude, and periovulatory E<sub>1</sub>3G concentration and slope. The results show that environmental exposure to PAHs is associated with endocrine markers of ovarian function in women.

## Acknowledgments

The authors thank the WHE study participants and the dedicated WHE staff, Claudia Shambaugh, Opalina Valencia, and Stacey Kojaku. The authors also thank the staff of the UC Irvine Institute for Clinical and Translational Science for assistance with data and specimen collection; Scott Littlejohn of Alere, Inc., Waltham, MA and Sarah Weddell of Swiss Precision Diagnostics Spark, Bedford, UK for converting fertility monitor data and assistance with interpretation of monitor data; and Dr. Lynn Flowers for exploratory analyses performed as part of her MS thesis in Environmental Health Sciences. The study was funded by the National Institutes of Health (NIH), National Institute of Environmental Health Sciences grant numbers ES016846 and ES020454 (to UL); grant number UL1 RR031985 from the National Center for Research Resources (NCR), a component of the NIH and the NIH Roadmap for Medical Research; a California Environmental Contaminant Biomonitoring Program RFI Laboratory Pilot Study; the UC Irvine Center for Occupational and Environmental Health.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.envint.2016.12.021>.

## References

- Alderete, E., Eskenazi, B., Sholtz, R., 1995. Effect of cigarette smoking and coffee drinking on time to conception. *Epidemiology* 6, 403–408.
- Aquilina, N.J., Delgado-Saborit, J.M., Meddings, C., Baker, S., Harrison, R.M., Jacob, P.I., Wilson, M., Yu, L., Duan, M., Benowitz, N.L., 2010. Environmental and biological monitoring of exposures to PAHs and ETS in the general population. *Environ. Int.* 36, 763–771.
- ATSDR, 1995. *Toxicological Profile for Polycyclic Aromatic Hydrocarbons*. US Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry, Atlanta, GA.
- Baird, D.D., Wilcox, A.J., 1985. Cigarette smoking associated with delayed conception. *JAMA* 253, 2979–2983.
- Baird, D.D., Weinberg, C.R., Wilcox, A.J., McConaughy, D.R., 1991. Using the ratio of urinary oestrogen and progesterone metabolites to estimate the day of ovulation. *Stat. Med.* 10, 255–266.
- Baird, D.D., McConaughy, D.R., Weinberg, C.R., Musey, P.I., Collins, D.C., Kesner, J.S., Knecht, E.A., Wilcox, A.J., 1995. Application of a method for estimating day of ovulation using urinary estrogen and progesterone metabolites. *Epidemiology* 6, 547–550.
- Baird, D.D., Weinberg, C.R., Zhou, H., Kamel, F., McConaughy, D.R., Kesner, J.S., Wilcox, A.J., 1999. Preimplantation urinary hormone profiles and the probability of conception in healthy women. *Fertil. Steril.* 71, 40–49.
- Baker, D., Park, C., Sweeney, C., McCormack, L., Durkin, M., Brenner, R., Dabelea, D., Entwisle, B., 2014. Recruitment of women in the National Children's Study initial vanguard study. *Am. J. Epidemiol.* 179, 1366–1374.
- Bartko, J.J., 1966. The intraclass correlation coefficient as a measure of reliability. *Psychol. Rep.* 19, 3–11.
- Borman, S.M., Christian, P.J., Sipes, I.G., Hoyer, P.B., 2000. Ovotoxicity in female Fischer rats and B6 mice induced by low-dose exposure to three polycyclic aromatic hydrocarbons: comparison through calculation of an ovotoxic index. *Toxicol. Appl. Pharmacol.* 167, 191–198.
- Buck Louis, G.M., Schisterman, E.F., Sweeney, A.M., Wilcosky, T.C., Gore-Langton, R.E., Lynch, C.D., Boyd Barr, D., Schrader, S.M., Kim, S., Chen, Z., Sundaram, R., 2011. Designing prospective cohort studies for assessing reproductive and developmental toxicity during sensitive windows of human reproduction and development – the LIFE study. *Paediatr. Perinat. Epidemiol.* 25, 413–424.
- Buck Louis, G.M., Sundaram, R., Schisterman, E.F., Sweeney, A.M., Lynch, C.D., Gore-Langton, R.E., Chen, Z., Kim, S., Caldwell, K.L., Barr, D.B., 2012. Heavy metals and couple fecundity, the LIFE study. *Chemosphere* 87, 1201–1207.
- Buck Louis, G.M., Sundaram, R., Schisterman, E.F., Sweeney, A., Lynch, C.D., Kim, S., Maisog, J.M., Gore-Langton, R., Eisenberg, M.L., Chen, Z., 2014a. Semen quality and time to pregnancy: the Longitudinal Investigation of Fertility and the Environment Study. *Fertil. Steril.* 101, 453–462.
- Buck Louis, G.M., Sundaram, R., Sweeney, A., Schisterman, E.F., Maisog, J., Kannan, K., 2014b. Urinary bisphenol a, phthalates, and couple fecundity: the Longitudinal Investigation of Fertility and the Environment (LIFE) study. *Fertil. Steril.* 101, 1359–1366.
- Chen, C., Wang, X., Wang, L., Yang, F., Tang, G., Xing, H., Ryan, L., Lasley, B., Overstreet, J.W., Stanford, J.B., Xu, X., 2005. Effect of environmental tobacco smoke on levels of urinary hormone markers. *Environ. Health Perspect.* 113, 412–417.
- Christensen, R., Johnson, W., Branscum, A., Hanson, T.E., 2011. *Bayesian Ideas and Data Analysis*. CRC Press, Boca Raton.
- Denissenko, M.F., Pao, A., Tang, M.-S., Pfeifer, G.P., 1996. Preferential formation of benzo[a]pyrene adducts at lung cancer mutational hotspots in P53. *Science* 274, 430–432.
- Fernandes, D., Porte, C., 2013. Hydroxylated PAHs alter the synthesis of androgens and estrogens in subcellular fractions of carp gonads. *Sci. Total Environ.* 447, 152–159.
- Harlow, B.L., Signorello, L.B., 2000. Factors associated with early menopause. *Maturitas* 35, 3–9.
- Hassan, M.A.M., Killick, S.R., 2004. Negative lifestyle is associated with a significant reduction in fecundity. *Fertil. Steril.* 81, 384–392.
- Hill, R.H.J., Head, S.L., Gregg, M., Shealy, D.B., Bailey, S.L., Williams, C.C., Sampson, E.J., Needham, L.L., 1995. Pesticide residues in urine of adults living in the United States: reference range concentrations. *Environ. Res.* 71, 99–108.
- IARC, 1983. *Polynuclear Aromatic Compounds, Part 1, Chemical, Environmental and Experimental Data*. World Health Organization, International Agency for Research on Cancer, Lyons, France.
- IARC, 2010. *Some Non-heterocyclic Polycyclic Aromatic Hydrocarbons and Some Related Exposures* ed<sup>s</sup>. World Health Organization, International Agency for Research on Cancer, Lyons, France.
- Jensen, T.K., Henriksen, T.B., Hjollund, N.H., Scheike, T., Kolstad, H., Giwercman, A., Ernst, E., Bonde, J.P., Skakkebaek, N.E., Olsen, J., 1998. Adult and prenatal exposures to tobacco smoke as risk indicators of fertility among 430 Danish couples. *Am. J. Epidemiol.* 148, 992–997.
- Kassam, A., Overstreet, J.W., Snow-Harter, C., De Souza, M.J., Gold, E.B., Lasley, B.L., 1996. Identification of anovulation and transient luteal function using a urinary pregnanediol-3-glucuronide ratio algorithm. *Environ. Health Perspect.* 104, 408–413.
- Kesner, J.S., Krieg, E.F., Knecht, E.A., Wright, D.M., 1992. Power analyses and immunoassays for measuring reproductive hormones in urine to assess female reproductive potential in field studies. *Scand. J. Work Environ. Health* 18 (Suppl. 2), 33–36.
- Li, Z., Sandau, C.D., Romanoff, L.C., Caudill, S.P., Sjödin, A., Needham, L.L., Patterson Jr., D.G., 2008. Concentration and profile of 22 urinary polycyclic aromatic hydrocarbon metabolites in the US population. *Environ. Res.* 107, 320–331.
- Li, Z., Romanoff, L., Bartell, S., Pittman, E.N., Trinidad, D.A., McClean, M., Webster, T.F., Sjödin, A., 2012. Excretion profiles and half-lives of ten urinary polycyclic aromatic hydrocarbon metabolites after dietary exposure. *Chem. Res. Toxicol.* 25, 1452–1461.
- Li, Z., Trinidad, D., Pittman, E.N., Riley, E.A., Sjödin, A., Dills, R.L., Paulsen, M., Simpson, C.D., 2015. Urinary polycyclic aromatic hydrocarbon metabolites as biomarkers to woodsmoke exposure – results from a controlled exposure study. *J. Expo. Sci. Environ. Epidemiol.* 1–8.
- Lodovici, M., Akpan, V., Evangelisti, C., Dolara, P., 2004. Sidestream tobacco smoke as the main predictor of exposure to polycyclic aromatic hydrocarbons. *J. Appl. Toxicol.* 24, 277–281.
- Lu, X., Kambe, F., Cao, X., Kozaki, Y., Kaji, T., Ishii, T., Seo, H., 2008. 3β-Hydroxysteroid-Δ24 reductase is a hydrogen peroxide-scavenger, protecting cells from oxidative stress-induced apoptosis. *Endocrinology* 149, 3267–3273.
- Luderer, U., Kesner, J.S., Fuller, J.M., Krieg, E.F.J., Meadours, J.W., Tramma, S.L., Yang, H., Baker, D., 2013. Effects of gestational and lactational exposure to heptachlor epoxide on age at puberty and reproductive function in men and women. *Environ. Res.* 121, 84–94.

- Mattison, D.R., 1979. Difference in sensitivity of rat and mouse primordial oocytes to destruction by polycyclic aromatic hydrocarbons. *Chem. Biol. Interact.* 28, 133–137.
- Mattison, D.R., Nightingale, M.S., 1982. Oocyte destruction by polycyclic aromatic hydrocarbons is not linked to the Inducibility of ovarian aryl hydrocarbon (benzo(a)pyrene) hydroxylase activity in (DBA/2N × C57BL/6N) F1 × DBA/2N backcross mice. *Pediatr. Pharmacol.* 2, 11–21.
- Mattison, D.R., Thorgeirsson, S.S., 1979. Ovarian aryl hydrocarbon hydroxylase activity and primordial oocyte toxicity of polycyclic aromatic hydrocarbons in mice. *Cancer Res.* 39, 3471–3475.
- Mattison, D.R., White, N.B., Nightingale, M.R., 1980. The effect of benzo(a)pyrene on fertility, primordial oocyte number, and ovarian response to pregnant Mare's serum gonadotropin. *Pediatr. Pharmacol.* 1, 143–151.
- Mattison, D.R., Plowchalk, D.R., Meadows, M.J., Miller, M.M., Malek, A., London, S., 1989. The effect of smoking on oogenesis, fertilization, and implantation. *Semin. Reprod. Endocrinol.* 7, 291–304.
- Menzie, C.A., Potocki, B.B., Santodonato, J., 1992. Ambient concentrations and exposure to carcinogenic PAHs in the environment. *Environ. Sci. Technol.* 26, 1278–1284.
- Montaquila, J.M., Brick, J.M., Curtin, L.R., 2010. Statistical and practical issues in the design of a national probability sample of births for the Vanguard Study of the National Children's Study. *Stat. Med.* 29, 1368–1376.
- Monteiro, P.R.R., Reis-Henriques, M.A., Coimbra, J., 2000a. Plasma steroid levels in female flounder (*Platichthys flesus*) after chronic dietary exposure to single polycyclic aromatic hydrocarbons. *Mar. Environ. Res.* 49, 453–467.
- Monteiro, P.R.R., Reis-Henriques, M.A., Coimbra, J., 2000b. Polycyclic aromatic hydrocarbons inhibit in vitro ovarian steroidogenesis in the flounder (*Platichthys flesus* L.). *Aquat. Toxicol.* 48, 549–559.
- Motorykin, O., Santiago-Delgado, L., Rohlman, D., Schrlau, J.E., Harper, B., Harris, S., Harding, A., Kile, M.L., Massey Simonich, S.L., 2015. Metabolism and excretion rates of parent and hydroxy-PAHs in urine collected after consumption of traditionally smoked Salmon for native American volunteers. *Sci. Total Environ.* 514, 170–177.
- NCS, 2007. The National Children's Study Research Plan. [https://ncsarchive.s-3.net/bioshare/study\\_documents/](https://ncsarchive.s-3.net/bioshare/study_documents/).
- Neal, M.S., Zhu, J., Holloway, A.C., Foster, W.G., 2007. Follicle growth is inhibited by benzo(a)pyrene, at concentrations representative of human exposure, in an isolated rat follicle culture assay. *Hum. Reprod.* 22, 961–967.
- Neal, M.S., Zhu, J., Foster, W.G., 2008. Quantification of benzo(a)pyrene and other PAHs in the serum and follicular fluid of smokers versus non-smokers. *Reprod. Toxicol.* 25, 100–106.
- Nethery, E., Wheeler, A.J., Fisher, M., Sjodin, A., Li, Z., Romanoff, L.C., Foster, W., Arbuckle, T.E., 2012. Urinary polycyclic aromatic hydrocarbons as a biomarker of exposure to PAHs in air: a pilot study among pregnant women. *J. Expo. Sci. Environ. Epidemiol.* 22, 70–80.
- NHANES, 2009. Fourth National Report on Human Exposure to Environmental Chemicals. Department of Health and Human Services, Centers for Disease Control and Prevention.
- Nykamp, J.A., Bols, N.C., Carlson, J.C., 2001. Phenanthrenequinone disrupts progesterone production in rat luteal cells. *Reprod. Toxicol.* 15, 393–398.
- Ogden, T.L., 2010. Handling results below the level of detection. *Ann. Occup. Hyg.* 54, 255–256.
- Penning, T.M., 2004. Aldo-keto reductases and formation of polycyclic aromatic hydrocarbon o-Quinones. *Methods Enzymol.* 378, 31–67.
- Robinson, J.E., Wakelin, M., Ellis, J.E., 2007. Increased pregnancy rate with use of the Clearblue Easy fertility monitor. *Fertil. Steril.* 87, 329–334.
- Romanoff, L.C., Li, Z., Young, K.J., Blakely, N.C.I., Patterson Jr., D.G., Sandau, C.D., 2006. Automated solid-phase extraction method for measuring urinary polycyclic aromatic hydrocarbon metabolites in human biomonitoring using isotope-dilution gas chromatography high-resolution mass spectrometry. *J. Chromatogr. B* 835, 47–54.
- Sadeu, J.C., Foster, W.G., 2011. Effect of in Vitro exposure to benzo(a)pyrene, a component of cigarette smoke, on folliculogenesis, steroidogenesis, and oocyte nuclear maturation. *Reprod. Toxicol.* 31, 402–408.
- Shealy, D.B., Barr, J.R., Ashley, D.L., Patterson Jr., D.G., Camann, D., Bond, A.E., 1997. Correlation of environmental carbaryl measurements with serum and urinary 1-naphthol measurements in a farmer applicator and his family. *Environ. Health Perspect.* 105, 510–513.
- Shopland, D.R., Burns, D.M., Benowitz, N.L., Amacher, R.H., 2001. Risks Associated with Smoking Cigarettes with Low Machine-Measured Yields of Tar and Nicotine: U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health, National Cancer Institute.
- Spiegelhalter, D.J., Best, N.G., Carlin, B.P., van der Linde, A., 2002. Bayesian measures of model complexity and fit. *J. Royal Stat. Soc. B* 64, 583–639.
- Sun, L., Zuo, Z., Chen, M., Chen, Y., Wang, C., 2015. Reproductive and transgenerational toxicities of phenanthrene on female marine Medaka (*Oryzias melastigma*). *Aquat. Toxicol.* 162, 109–116.
- Suwan-Ampai, P., Navas-Acien, A., Strickland, P.T., Agnew, J., 2009. Involuntary tobacco smoke exposure and urinary levels of polycyclic aromatic hydrocarbons in the United States, 1999–2002. *Cancer Epidemiol. Biomark. Prev.* 18, 884–893.
- Takizawa, K., Yagi, H., Jerina, D.M., Mattison, D.R., 1984. Murine strain differences in ovotoxicity following intraovarian injection with benzo(a)pyrene, (+)-(7R,8S)-oxide, (–)-(7R,8S)-Dihydrodiol, or (+)-(7R,8S)-diol-(9S,10R)-epoxide-2. *Cancer Res.* 44, 2571–2576.
- Tanabe, K., Susumu, N., Hand, K., Nishii, K., Ishikawa, I., Nozawa, S., 2001. Prediction of the potentially fertile period by urinary hormone measurements using a new home-use monitor: comparison with laboratory hormone analyses. *Hum. Reprod.* 16, 1619–1624.
- Tsai-Turton, M., Nakamura, B.N., Luderer, U., 2007. Induction of apoptosis by 9,10-dimethyl-1,2-benzanthracene (DMBA) in cultured preovulatory rat follicles is preceded by a rise in reactive oxygen species and is prevented by glutathione. *Biol. Reprod.* 77, 442–451.
- van Noord, P.A.H., Boersma, H., Dubas, J.S., te Velde, E.R., Dorland, M., 1997. Age at natural menopause in a population-based screening cohort: the role of menarche, fecundity, and lifestyle factors. *Fertil. Steril.* 68, 95–102.
- Verma, N., Pink, M., Retteneier, A.W., Schmitz-Spanke, S., 2012. Review on proteomic analyses of benzo(a)pyrene toxicity. *Proteomics* 12, 1731–1755.
- Walle, T., Gaffney, T.E., 1972. Propranolol metabolism in man and dog: mass spectrometric identification of six new metabolites. *J. Pharmacol. Exp. Ther.* 182, 83–92.
- Windham, G.C., Elkin, E.P., Swan, S.H., Waller, K.O., Fenster, L., 1999. Cigarette smoking and effects on menstrual function. *Obstet. Gynecol.* 93, 59–65.
- Windham, G.C., Mitchell, P., Anderson, M., Lasley, B.L., 2005. Cigarette smoking and effects on hormonal function in premenopausal women. *Environ. Health Perspect.* 113, 1285–1290.
- Wyatt, K.M., Dimmock, P.W., Walker, T.J., O'Brien, P.M.S., 2001. Determination of total menstrual blood loss. *Fertil. Steril.* 76, 125–131.
- Xue, W., Warshawsky, D., 2005. Metabolic activation of polycyclic aromatic hydrocarbon and heterocyclic aromatic hydrocarbons and DNA damage: a review. *Toxicol. Appl. Pharmacol.* 206, 73–93.
- Zumoff, B., Miller, L., Levit, C.D., Miller, E.H., Heinz, U., Kalin, M., Denman, H., Jandorek, R., Rosenfeld, R.S., 1990. The effect of smoking on serum progesterone, estradiol, and luteinizing hormone levels over a menstrual cycle in normal women. *Steroids* 55, 507–511.