

Available online at www.sciencedirect.com

Metabolism

www.metabolismjournal.com

New horizons for focused ultrasound (FUS) – therapeutic applications in neurodegenerative diseases



Diane B. Miller*, James P. O'Callaghan**

Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, Morgantown, WV 26505

ARTICLE INFO

Keywords:

Microbubbles
Obicodilation
Sonothrombolysis
Transcranial FUS
Transcranial

ABSTRACT

Access to the CNS and delivery of therapeutics across the blood–brain barrier remains a challenge for most treatments of major neurological diseases such as AD or PD. Focused ultrasound represents a potential approach for overcoming these barriers to treating AD and PD and perhaps other neurological diseases. Ultrasound (US) is best known for its imaging capabilities of organs in the periphery, but various arrangements of the transducers producing the acoustic signal allow the energy to be precisely focused (F) within the skull. Using FUS in combination with MRI and contrast agents further enhances accuracy by providing clear information on location. Varying the acoustic power allows FUS to be used in applications ranging from imaging, stimulation of brain circuits, to ablation of tissue. In several transgenic mouse models of AD, the use of FUS with microbubbles reduces plaque load and improves cognition and suggests the need to investigate this technology for plaque removal in AD. In PD, FUS is being explored as a way to non-invasively ablate the brain areas responsible for the tremor and dyskinesia associated with the disease, but has yet to be utilized for non-invasive delivery of putative therapeutics. The FUS approach also greatly increases the range of possible CNS therapeutics as it overcomes the issues of BBB penetration. In this review we discuss how the characteristics and various applications of FUS may advance the therapeutics available for treating or preventing neurodegenerative disorders with an emphasis on treating AD and PD.

© 2017 Published by Elsevier Inc.

1. Introduction

The strikingly detailed images produced by diagnostic ultrasound (US) have made this technology and its applications familiar to physicians and the lay public alike. In this brief article we will review a modification/extension of US technology, focused ultrasound (FUS), that may find application in the treatment of AD and PD, neurodegenerative diseases that have proven resistant to current therapies. Unfortunately, to date, the available treatments for these devastating diseases are merely palliative and directed toward ameliorating symptoms (e.g., acetylcholinesterase inhibitors in AD, levo-

dopa in PD) but do not prevent, cure, or slow disease progression [1,2]. FUS would allow precise targeting of the diseased brain areas to more effectively deliver potential therapeutics than systemic delivery methods. As most neurodegenerative diseases do not affect the brain in a global fashion, the current means of making a precise delivery to the target site (e.g. direct injection) are highly invasive and can involve a craniotomy or placement of a cannula, etc. Although other techniques used to deliver therapeutics to brain (e.g., intranasal) are not invasive, they are inefficient at precise delivery to the desired target site; high potency agents are required to offset the dilution caused by travel distance to the

* Publication of this article was supported by the Collège International de Recherche Servier (CIRS).

** Correspondence to: Tel.: +1 304 285 5732.

E-mail addresses: dum6@cdc.gov (D.B. Miller), jdo5@cdc.gov (J.P. O'Callaghan).

site and brain wide delivery remains an unwanted outcome. In contrast, FUS may be able to overcome many of the technical issues involved in direct delivery of therapeutics to a target brain area in an efficient and safe fashion (e.g., [3]). Here, we highlight the relevant properties of FUS that make this technology suitable in a number of ways for treating neurodegenerative diseases such as AD and PD. Of note, others have described the beneficial properties of both US and FUS that make them suitable for neurological applications and treating a variety of CNS diseases [4–7].

2. US and FUS Defined

US technology may seem “magical” to those awed by the detail of its imaging capability, but in truth it is a mechanically produced pressure/sound wave of 20 kHz or greater with physical properties like that of any other sound wave. In simple terms, a transducer or piezoelectric crystal is electrically stimulated and produces an oscillation/vibration (i.e., sound wave) that moves through the transmission media. The 20 kHz wave is a higher frequency than most humans can hear; hence the label ultra. The US can be moved directly through, or be refracted, absorbed or reflected by the media which can be air, water or tissue. Of most importance for therapeutic applications, these waves possess the remarkable ability to penetrate deep into various solid substances, including brain tissue. US produces biological effects in living tissue that are considered to be due to thermal or mechanical actions and consequently there are low and high power applications of US with heating likely dominating when the power is high [8]. Low power uses include enhanced drug uptake through the skin, i.e., sonophoresis, and momentary disruption of the cell membrane to facilitate delivery of therapeutic agents, i.e., sonoporation. In FUS, an array of transducers is used to produce multiple beams of US which are focused on the target via an acoustic lens, resulting in a therapeutic effect only where the beams intersect, but not where the individual beams pass through tissue. FUS is frequently used in conjunction with real-time imaging by US or MRI to monitor the target site. This technology has been used in the treatment of uterine fibroids, kidney stones, blood clots, excessive bleeding in traumatic injury, and in cosmetic medicine for fat removal.

3. Microbubbles – Application in FUS

These micron-scale gas bubbles are frequently used in conjunction with FUS and US. They are about the size of a red blood cell (~10 or less μm diameter) and are stabilized by a shell of albumin, lipid, protein or polymer – if not they would quickly disperse into blood once they enter the circulation. Microbubbles have seen wide usage as they can both image and treat the target and for these reasons are considered theranostic agents [9]. FUS activates the microbubbles when they pass through the focal point of the converging beams causing an expansion and contraction of the gas core. In brain this can affect cellular structure and result in an opening of

the tight junctions between the endothelial cells of the BBB at the focus site, allowing delivery of the substances comprising the shell or carried by the shell. There appear to be no long term consequences of the opening and closing of the BBB in this fashion [e.g., 10]. Both imaging and therapeutic agents have been delivered in this manner.

4. Barriers to Effective Treatment of the most Common Neurodegenerative Disorders, AD and PD - Using FUS to Improve Treatment

Although AD and PD are among the most prevalent neurodegenerative disorders, efforts at developing treatments of these diseases have been hampered by a lack of understanding of their causes, as well as a paucity of biomarkers with which to diagnose and follow disease progression in living patients. Both diseases involve the death of neurons that begins long before the classic symptoms that define each disease are seen. The protracted course of neuronal death long before symptoms occur makes it difficult to prevent either disease. So does the lack of suitable biomarkers for prodromal detection and for evaluating the therapeutic efficacy of various treatments, especially in early stage disease. To date, efforts to slow or prevent either of these diseases have been unsuccessful [1,2,11,12].

Therapeutic efforts in AD largely have focused on β -amyloid. Aberrant turnover, metabolism and deposition of amyloid precursor protein (APP) and amyloid- β peptide, along with possible aberrant phosphorylation of tau, lead to large aggregations of these proteins and formation of amyloid plaques. Amyloid plaques are considered by many to be a risk factor and an underlying cause in the development of AD. Thus, the amyloid hypothesis [13] drives treatment strategies for AD because elimination or reduction of amyloid plaques may reduce the neurodegeneration that ultimately serves as the basis for this disease. Despite positive results from preclinical studies utilizing various mouse models of AD, effective therapies have been harder to achieve than first expected [11]. The mouse studies demonstrated the ability of passive and active immune strategies to reduce or eliminate plaque and restore cognitive abilities. However, these immunotherapy strategies have not been as effective in humans and/or clinical trials were halted prematurely due to unexpected side-effects. The human trial failures may be due to the necessity of delivering antibodies systemically, resulting in an insufficient amount of antibody at the appropriate brain target or a failure of the aged AD patient to mount a sufficient antibody response to the administered antigen. Although A β antibodies cross the BBB, they do so at a very low level (~0.1–2.0% of the administered dose) making it difficult to achieve an efficacious amount of the antibody in brain which is crucial [14]. Due to this therapeutic obstacle, cost can be an issue because of the large volume of monoclonal antibodies needed for efficacy with systemic treatment. FUS offers a non-invasive and potentially inexpensive approach to achieving targeted therapeutically effective levels of A β immunotherapy. In an AD mouse model, FUS with microbubbles delivered [15] antibodies directly to brain at levels able to

rapidly (within 4 days) reduce pathology. Also, it appears that FUS with microbubbles may allow endogenous IgG and IgM A β antibodies to enter brain through the opened BBB as both types were found bound to plaques in treated cortex [16].

As immunological strategies for AD treatment have not yet been successful, non-immunological or pharmacological approaches to prevent, reduce or eliminate amyloid plaques are being sought. Recent studies with AD mouse models and FUS or scanning (the focus of the beam is moved in small increments to treat a defined area) FUS, employing a limited number of exposures (~1/week for 5–6 weeks) in conjunction with microbubbles, show opening of the BBB allowing entry of albumin and immunoglobulins. Significant reductions of amyloid plaques and the activation of microglia involved in plaque clearance were obtained with this non-immune approach. Furthermore, the plaque reduction was accompanied by an improvement in cognition and evidence of neurogenesis, findings suggestive of increased neuronal plasticity [16–18]. It has not been determined if the plaque reduction, microglia activation and clinical improvement can be induced by just FUS without the use of microbubbles or entry of albumin into brain. Albumin is known to activate microglia and almost all circulating A β is bound to it [19,20]. AD 3xTg mice develop significant A β pathology and direct infusion of human serum albumin (HAS) for ~ a month significantly reduced the plaque burden as well as the level of the toxic A β oligomers [21]. In a study with a small number of AD patients, exchange of endogenous plasma with plasma containing 5% albumin (prepared from whole human blood and available at different percentages as a therapeutic) to increase the patient's level, over a several month period, resulted in reduction of A β 40 and A β 42 in CSF. Reduced A β peptide levels were associated with stability in cognitive scores apparent at 1-year follow-up [22]. It is clear that albumin can bind A β and aid in its clearance. However, direct delivery of albumin to brain in AD patients has not been tested with FUS-delivered microbubbles containing albumin, an approach that may be an effective treatment strategy for increasing A β clearance from brain.

PD, like AD, is characterized by a pathologic hallmark involving a misfolded/aggregated protein – namely, α -synuclein found in the extracellular space, and in Lewy bodies and neurites [14]. Alpha-synuclein also is present in CSF and plasma of PD patients. Thus, a reduction of extracellular α -synuclein, as well as preventing its aggregation into large fibrillary clusters, are clear therapeutic strategies for various immunotherapies [23]. In animal models of PD, active immunotherapies were effective at reducing the accumulated protein in neurons, they reduce α -synuclein inclusions in substantia nigra, and they reduce the neurodegeneration caused by accumulation of the misfolded protein. Passive immunization studies also have been effective in reducing the α -synuclein pathology and associated behavioral problems. Limited clinical information is available concerning immunotherapy, with little activity being undertaken in implementing this potential therapeutic in PD. However, it would be expected that many of the problems found in developing immunotherapies for AD might be expected to occur in PD applications. Accordingly, FUS

applications would be expected to reduce or mitigate the problems of PD immunotherapies in humans.

In PD, unlike AD, where immunotherapies are in the forefront, the main therapeutic focus has been on finding ways to replace or stave off the continuing degeneration of the affected DA neurons of the substantia nigra. Considerable research with various animal models of neurodegeneration has shown that many different NTFs can protect and repair damaged neurons [24,25]. DA containing neurons, in particular, appear very responsive to and dependent for survival on various NTFs, especially those of the transforming growth factor- β (TGF β) family. This family includes glial cell-line derived neurotrophic factor (GDNF), neurturin, and growth/differentiation factor 5 (GDF5). Of note, several members of this family have been evaluated in clinical trials. Other NTFs have been identified as well that display beneficial effects in the compromised nigral striatal pathway. Due to promising preclinical work in animal models of PD, there have been a number of human trials to determine if NTFs are efficacious in the treatment of PD. There has been no successful translation of these treatments to humans. The failure in translation may be directly related to technical issues concerning delivery of the factors to the target area in sufficient quantities.

NTFs are too large to cross the BBB and are rapidly metabolized in the periphery which necessitates their direct delivery to brain or a means to generate them locally within the brain. NTFs have been delivered to or produced in selected brain sites by direct infusion, implants of stem cells genetically modified to overexpress a particular NTF, or introduction of recombinant viral vectors able to effect long-term expression in the target cells. Intranasal administration of NTFs can result in their entry into the CNS but not to specific targets. These drawbacks in current therapies with NTFs may be overcome with the application of FUS. Indeed, preclinical work has demonstrated the ability of FUS to deliver sufficient levels of neurturin to the nigral striatal area of mouse brain. FUS delivered a greater amount of the factor than direct injection and engendered the expected bioactivity of this NTF [26]. To get the same coverage of the target structure of caudate-putamen in humans would require placement of multiple cannulas. Interestingly, intranasal administration of substances in conjunction with FUS and microbubbles was able to effect targeted delivery to the caudate putamen and may be another means of delivery of neurotrophic factors [3].

Also of note for potential treatment of AD and PD are the temporary and reversible neuromodulating and neurogenic properties of FUS. These properties may be beneficial in slowing or repairing neuronal damage in neurodegenerative disorders. Low-intensity FUS without microbubbles can alter neurotransmitter levels and stimulate or suppress brain activity in multiple areas (e.g., visual, motor, etc.), without causing apparent damage to brain. Furthermore, FUS, when applied along with microbubbles, can open the BBB and stimulate cell proliferation and neurogenesis in hippocampus [27,28]. The observed neurogenesis in this brain area may be due to increases in trophic or growth factors (BDNF, VEGF, bFGF). That FUS can increase these factors may be of benefit in both AD and PD.

5. Possible Safety Issues

The acoustic energy of FUS can be converted to thermal or mechanical energy and used as a tool for ablating brain tumors, epileptic foci, and brain areas causing essential tremor and the dyskinesia of PD [6]. Despite the potential for FUS at this level to cause damage, at lower energy levels this technology appears relatively safe. At lower acoustic energies in conjunction with microbubbles, long-term repeated exposure (4–20 months) causes focused opening of the BBB in the cortex or caudate-putamen area with no apparent signs of tissue damage (e.g., edema, hemorrhage) or deficits in a variety of behavioral tests [29]. It also should be noted that transcranial continuous US has been used for many years at low levels to visualize the human brain for diagnostic purposes in echoencephalography; no safety concerns have arisen despite the potential to cause elevated brain temperatures. Use of MRI for accurate targeting along with imaging of the outcome, as well as use of real-time acoustic feedback controllers, is expected to result in an improved safety profile for FUS [30]. Animal research also shows FUS is able to stimulate brain motor pathways and neurotransmitter levels, induce cortical excitation, and modulate brain region-specific activity without BBB opening. These effects do not appear to be due to thermal or mechanical mechanisms and are suggestive of a margin of safety.

Of course, wider transition of the technology to clinical application will require more extensive testing prior to using it routinely. A more thorough examination of the possible safety issues that may occur due to repeated BBB opening, or other aspects of the technology in therapeutic applications, will need to be undertaken. Depending on the acoustic pressure and pulse applied, the BBB can remain open for minutes to hours [31]. Although FUS carries little risk of infection when used to deliver therapeutics, there may be unintended consequences. For example, unsafe levels of cavitation and subsequent heating or ablation of tissue occurring at temperatures above 55°C may lead to damage and prolonged BBB opening. This condition may promote the unwanted brain entry of cells (e.g., immune cells like macrophages, red blood cells, bacteria, viruses, etc.), proteins and peptides (albumin, chemokines, cytokines, etc.) from the periphery. Other unwanted consequences may include failure of the BBB to reclose, injury due to unexpected mechanical effects, unexpected interactions with the anesthesia needed in certain applications, or problematic properties (toxicity, microvascular effects, etc.) of microbubbles [32]. Careful attention to controlling the degree of BBB opening may mitigate some of these concerns [16].

Finally, to move forward with FUS for treatment of AD and PD will require continued development of equipment, as the shape and positioning of the transducers determines the brain area that can be targeted. Many efficacy and safety studies have been conducted in experimental animals whose skulls are much thinner and differently shaped than that of humans. Even though some studies have involved larger animals [10,29,33] with thicker skulls (e.g., non-human primates and pigs), the human skull is quite variable in shape, density, and thickness. Bone is much more effective at mitigating acoustic energy than soft tissue. The characteristics of the human skull

makes focusing the multiple beams and delivery of effective US levels more difficult, as the individual beams may get out of phase. Care must be taken to mitigate focal heating and reduce the chance of injury along the beam path. Thus, the thickness and shape of the skull are more of a barrier for the delivery of FUS in humans than in experimental animals. Also, BBB opening duration may depend on individual patient characteristics as well as the degree of anesthesia [29,31]. Consequently, exposure parameters may need to be adjusted for each individual to insure delivery of the desired amount of acoustic energy.

6. Summary

As we have noted, recent animal studies suggest FUS alone or in conjunction with microbubbles may provide a quick, efficient, and non-invasive way to deliver therapeutics to brain for treatment of AD and PD. Many disease modifying therapies do not effectively cross the BBB, including antibodies, genes, growth factors, stem cells, etc. However, the successful preclinical applications of FUS technology suggest that its introduction in clinical settings may provide new avenues for treatment of neurodegenerative diseases.

Acknowledgements

Publication of this article was supported by the College International de Recherche Servier (CIRS).

Conflicts of Interest

None.

REFERENCES

- [1] Cacabelos R, Torrellas C, Carrera I, Cacabelos P, Corzo L, Fernández-Novoa L, et al. Novel therapeutic strategies for dementia. *CNS Neurol Disord Drug Targets* 2016;15: 141–241.
- [2] Miller DB, O’Callaghan JP. Biomarkers of Parkinson’s disease (PD): present and future. *Metabolism* 2015;64(Suppl. 1):S40–6.
- [3] Chen H, Chen CC, Acosta C, Wu SY, Sun T, Konofagou EE. A new brain drug delivery strategy: focused ultrasound-enhanced intranasal drug delivery. *PLoS One* 2014;9(10): e108880. <http://dx.doi.org/10.1371/journal.pone.0108880>.
- [4] Aryal M, Arvanitis CD, Alexander PM, McDannold N. Ultrasound-mediated blood–brain barrier disruption for targeted drug delivery in the central nervous system. *Adv Drug Deliv Rev* 2014;72:94–109.
- [5] Ghanouni P, Pauly KB, Elias WJ. Transcranial MR-guided focused ultrasound: a review of the technology and neuro applications. *AJR Am J Roentgenol* 2015;205:150–9.
- [6] Piper RJ, Hughes MA, Moran CM, Kandasamy J. Focused ultrasound as a non-invasive intervention for neurological disease: a review. *Br J Neurosurg* 2016;30:286–93.
- [7] Leinenga G, Langton C, Nisbet R, Götz J. Ultrasound treatment of neurological diseases – current and emerging applications. *Nat Rev Neurol* 2016;12:161–74.

- [8] ter Haar G. Therapeutic applications of ultrasound. *Prog Biophys Mol Biol* 2007;93:111–29.
- [9] Sirsi S, Borden M. Microbubble compositions, properties and biomedical applications. *Bubble Sci Eng Technol* 2009;1:3–17.
- [10] McDannold N, Aarvanitis CD, Vykhodtseva N, Livingstone MS. Temporary disruption of the blood–brain barrier by use of ultrasound and microbubbles: safety and efficacy evaluation in rhesus macaques. *Cancer Res* 2012;72:3653–63.
- [11] Herrup K. The case for rejecting the amyloid cascade hypothesis. *Nat Neurosci* 2015;18:794–9.
- [12] Toyn J. What lessons can be learned from failed Alzheimer's disease trials? *Expert Rev Clin Pharmacol* 2015;8:267–9.
- [13] Selkoe DJ, Hardy J. The amyloid hypothesis of Alzheimer's disease at 25 years. *EMBO Mol Med* 2016. <http://dx.doi.org/10.15252/emmm.201606210>.
- [14] George S, Brundin P. Immunotherapy in Parkinson's disease: micromanaging alpha-synuclein aggregation. *J Parkinsons Dis* 2015;5:413–24.
- [15] Jordão JF, Ayala-Grosso CA, Markham K, Huang Y, Chopra R, McLaurin J, et al. Antibodies targeted to the brain with image-guided focused ultrasound reduces amyloid-beta plaque load in the TgCRND8 mouse model of Alzheimer's disease. *PLoS One* 2010;5(5):e10549.
- [16] Jordão JF, Thévenot E, Markham-Coultes K, Scarcelli T, Weng YQ, Khima K, et al. Amyloid- β plaque reduction, endogenous antibody delivery and glial activation by brain-targeted, transcranial focused ultrasound. *Exp Neurol* 2013;248:16–29.
- [17] Burgess A, Dubey S, Yeung S, Hough O, Eterman N, Aubert I, et al. Alzheimer disease in a mouse model: MR imaging-guided focused ultrasound targeted to the hippocampus opens the blood–brain barrier and improves pathologic abnormalities and behavior. *Radiology* 2014;273:736–45.
- [18] Leinenga G, Gotz J. Scanning ultrasound removes amyloid- β and restores memory in an Alzheimer's disease mouse model. *Sci Transl Med* 2015;7(278):278ra33.
- [19] Prajapati KD, Sharma SS, Roy N. Current perspectives on potential role of albumin in neuroprotection. *Rev Neurosci* 2011;22:355–63.
- [20] Ranaivo HR, Wainwright MS. Albumin activates astrocytes and microglia through mitogen-activated protein kinase pathways. *Brain Res* 2010;1313C:222–31.
- [21] Ezra A, Rabinovich-Nikitin R, Rabinovich-Toidman P, Solomon B. Multifunctional effect of human serum albumin reduces Alzheimer's disease related pathologies in the 3xTg mouse model. *J Alzheimers Dis* 2016;50:175–88.
- [22] Boada M, Ortiz P, Anaya F, Hernández I, Muñoz J, Núñez L, et al. Amyloid-targeted therapeutics in Alzheimer's disease: use of human albumin in plasma exchange as a novel approach for A β mobilization. *Drug News Perspect* 2009;22:325–39.
- [23] Lindstrom V, Lindström V, Ihse E, Fagerqvist T, Bergström J, Nordström E, et al. Immunotherapy targeting A-synuclein, with relevance for future treatment of Parkinson's disease and other Lewy body disorders. *Immunotherapy* 2014;6:141–53.
- [24] Cummins G, Barker RA. What is the most promising treatment for Parkinson's disease: genes, cells, growth factors or none of the above? *Regen Med* 2012;7:617–21.
- [25] Kelly MD, O'Keeffe GW, Sullivan AM. Viral vector delivery of neurotrophic factors for Parkinson's disease therapy. *Expert Rev Mol Med* 2015;17:1–14.
- [26] Samiotaki G, Acosta C, Wang S, Konofagou EE. Enhanced delivery and bioactivity of the neurturin neurotrophic factor through focused ultrasound-mediated blood–brain barrier opening *in vivo*. *J Cereb Blood Flow Metab* 2015;35:611–22.
- [27] Scarcelli T, Jordão JF, O'Reilly MA, Ellens N, Hynynen K, Aubert I. Stimulation of hippocampal neurogenesis by transcranial focused ultrasound and microbubbles in adult mice. *Brain Stimul* 2014;7:304–7.
- [28] Yoo S-S, Bystritsky A, Lee J-H, Zhang Y, Fischer K, Min BK, et al. Focused ultrasound modulates region-specific brain activity. *Neuroimage* 2011;56:1267–75.
- [29] Downs ME, Buch A, Sierra C, Karakatsani ME, Teichert T, Chen S, et al. Long-term safety of repeated blood–brain barrier opening via focused ultrasound with microbubbles in non-human primates performing a cognitive task. *PLoS One* 2015;10(5):e0125911. <http://dx.doi.org/10.1371/journal.pone.0125911>.
- [30] Burgess A, Shah K, Hough O, Hynynen K. Focused ultrasound-mediated drug delivery through the blood–brain barrier. *Expert Rev Neurother* 2015;15:477–91.
- [31] Samiotaki G, Konofagou EE. Dependence of the reversibility of focused-ultrasound-induced blood–brain barrier opening on pressure and pulse length *In Vivo*. *IEEE Trans Ultrason Ferroelectr Freq Control* 2013;60:2257–65.
- [32] ter Haar G. Safety and bio-effects of ultrasound contrast agents. *Med Biol Eng Comput* 2009;47:893–900.
- [33] Xie F, Boska MD, Lof J, Uberti MG, Tsutsui JM, Porter TR. Effects of transcranial ultrasound and intravenous microbubbles on blood brain barrier permeability in a large animal model. *Ultrasound Med Biol* 2008;34:2028–34.