

Association of exposure to polycyclic aromatic hydrocarbons (estimated from job category) with concentration of 1-hydroxypyrene glucuronide in urine from workers at a steel plant

Daehee Kang, Nathaniel Rothman, Soo-Hun Cho, Hyun Sul Lim, Ho-Jang Kwon, Sun-Min Kim, Brian Schwartz, Paul T Strickland

Abstract

Objectives—Increased risk of lung cancer has been associated with employment in the steel industry. This association is thought to be due in part to increased concentrations of polycyclic aromatic hydrocarbons (PAHs) in air found in this work environment. Measurement of PAH metabolites in human urine provides a means of assessing individual internal dose of PAHs. This study examined the relative contribution of occupation and smoking to urinary concentration of 1-hydroxypyrene glucuronide (1-OHPG) among a group of workers at a steel plant.

Methods—Concentrations of 1-OHPG in urine from 44 workers with jobs associated with increased air concentrations of PAHs and 40 workers with jobs with low or no exposure to PAHs were measured. 20 workers in each group were not current smokers. Urinary 1-OHPG was measured by synchronous fluorescence spectroscopy after immunoaffinity chromatography specific for PAH metabolites.

Results—Mean (SEM) urinary 1-OHPG concentration was 2.16 (0.42) pmol/ml urine among the 44 occupationally exposed workers compared with 0.38 (0.05) among the 40 workers with no or low exposure ($P < 0.0001$). Mean urinary 1-OHPG concentration was 1.82 (0.41) pmol/ml urine among the 44 current smokers compared with 0.75 (0.20) among the 40 non-smokers ($P < 0.005$). Mean 1-OHPG concentrations in non-smokers were 0.26 ($n = 20$), 0.70 ($n = 15$), and 2.84 pmol/ml urine ($n = 5$) for strata of exposure to PAHs (no or low, mid, and high) based on job category; the corresponding values in smokers were 0.55 ($n = 20$), 0.94 ($n = 12$), and 4.91 pmol/ml ($n = 12$), respectively. Multiple linear regression showed significant differences between subjects in different PAH exposure strata and between smokers and non-smokers. Both smoking and occupational exposure to PAHs were associated with increased concentrations of 1-OHPG in urine. Amounts of foods containing PAHs ingested by this group of workers were relatively low and did not

contribute significantly to urinary 1-OHPG concentrations.

Conclusions—These results indicate that 1-OHPG is a common urinary metabolite in people with recent occupational exposure to PAHs and is associated with both job category and estimated stratum of PAH exposure.

(*Occup Environ Med* 1995;52:593-599)

Keywords: polycyclic aromatic hydrocarbons; 1-hydroxypyrene glucuronide; urinary biomonitoring

Polycyclic aromatic hydrocarbons (PAHs) are produced during incomplete combustion of organic materials, and humans are exposed to these compounds from a variety of occupational, environmental, personal, and dietary sources.^{1,2} In the steel industry, increased air concentrations of PAHs are found in several work areas, particularly near blast furnaces and coke ovens.³ Urinary biomarkers of PAH exposure or internal dose have been proposed as a means of assessing recent exposure to these compounds. Urinary metabolites of PAHs are useful because of their ease of collection and extensive chemical characterisation. Becher and Bjørseth⁴ developed an analytical procedure to measure PAHs in human urine by chemical reduction of metabolites to parent PAHs with subsequent analysis by high performance liquid chromatography (HPLC) with fluorescence detection. Application of this method to urine samples from workers at aluminum plants⁵ and coke ovens⁶ did not show differences in urinary PAH concentrations between exposed workers and non-exposed controls.

After Keimig *et al* reported that 1-hydroxypyrene was a major metabolite of the carcinogen pyrene in pig urine,⁷ Jongeneelen *et al*⁸⁻¹⁵ conducted a series of studies evaluating the excretion of 1-hydroxypyrene in rodent and human urine. They measured this metabolite in urine of people occupationally exposed to PAHs (coke-oven workers, petroleum coke handlers, creosote impregnating plant workers) or therapeutically exposed (psoriatic patients treated with mineral coal tar and patients with eczematous dermatitis undergoing coal tar treatment).⁹⁻¹⁶ Excretion

Division of Occupational Health, Department of Environmental Health Sciences, Johns Hopkins School of Hygiene and Public Health, Baltimore, MD, USA
D Kang
B Schwartz
P T Strickland

Occupational Studies Section, Epidemiology and Biostatistics Program, National Cancer Institute, Bethesda, MD, USA
N Rothman

Department of Preventive Medicine, Seoul National University, Seoul, Korea
S-H Cho
H-J Kwon
S-M Kim

Department of Preventive Medicine, Dongguk University, Pohang, Korea
H S Lim

Present Address: Hazard Evaluation and Technical Assistance Branch, NIOSH, Cincinnati, OH 45226, USA
D Kang

Correspondence to: Dr Paul Strickland, Department of Environmental Health Sciences, Johns Hopkins School of Public Health, 615 North Wolfe Street, Baltimore, MD 21205, USA.

Accepted 25 May 1995

of 1-hydroxypyrene occurred fairly rapidly (6–35 h) after exposure to coke oven emissions.⁹ These studies analysed urine after enzymatic hydrolysis treatment with β -glucuronidase or aryl-sulphatase to convert conjugated metabolites to 1-hydroxypyrene. We have recently developed a sensitive and specific method to measure 1-hydroxypyrene-glucuronide (1-OHPG), the most abundant water soluble metabolite of pyrene in unhydrolysed urine.¹⁷ To investigate the usefulness of this metabolite as a potential biomarker of occupational exposure to PAHs, we examined the relative contribution of occupation and smoking on urinary concentration of 1-OHPG among a group of steel plant workers in a modern steel production factory. As different job categories within the steel industry are associated with exposure to varying concentrations of combustion product and risk of lung cancer,¹⁸ we examined urinary 1-OHPG concentration in workers with various jobs within the steel plant.

Methods

STUDY POPULATION

Study participants were employees at a modern steel plant located in South Korea. Urine samples were collected in conjunction with an annual physical examination after informed consent was provided to participate in the study. Urine samples (40 ml) were frozen at -70°C , shipped to the United States on dry ice, and stored at -70°C for four months before analysis. The 44 exposed workers had jobs that dealt with the processing of raw materials and steel including coal, coke, iron ore, molten iron, and molten steel. The 40 control subjects worked mainly in office areas that are located about 2 km from the iron and steel making plant. Also included in this control group were non-office support workers in the plant that included construction and transportation workers. Concurrent personal air sampling information was not available; however, personal air measurements were available from one year before the urine collection in a group of workers at the coke plant only.

Job categories were used to assign to the workers relative PAH exposures — that is, no, low, mid, or high PAH exposure based on the scheme published by Bjørseth and Becher³ (table 1). There were 20 non-smokers in the combined mid and high exposure groups and

20 non-smokers in the combined no and low exposure groups. A self-administered questionnaire was used to obtain subject information that included smoking habits, job history, major diseases, dietary habits, drug use, and possible use of personal protective equipment. Although personal protective equipment was available in some of the mid and high exposure groups, its use was difficult to confirm by questionnaire.

ASSAY FOR 1-HYDROXYPYRENE GLUCURONIDE

The 1-OHPG was measured in urine with a recently developed assay.¹⁷ Urine samples (2 ml) were treated with 0.1N HCl (90°C , 60 min), neutralised, and loaded onto Sep-pak C18 cartridges (Waters). After washing with 30% methanol, the relatively non-polar metabolites were eluted with 4 ml of 80% methanol and the volume of eluate was reduced to 0.5 ml by evaporation. The concentrated samples were diluted to 4 ml with 15 mM phosphate buffered saline (PBS) and loaded on to immunoaffinity columns prepared with 0.8 ml cyanogen bromide activated sepharose 4B (Sigma) coupled with monoclonal antibody 8E11 that recognises several PAH-DNA adducts and metabolites.¹⁹ We have previously shown that 1-OHPG binds to these columns.¹⁷ After washing the columns three times with 4 ml of 15 mM PBS, compounds bound to the monoclonal antibody 8E11 were eluted with 2 ml of 40% methanol in three fractions. Eluate fractions were analysed by synchronous fluorescence spectroscopy with a Perkin-Elmer LS50 fluorescence spectrometer. The excitation-emission monochromators were driven synchronously with a wavelength difference of 34 nm. Under these conditions 1-OHPG produces a characteristic fluorescence emission maximum at 381 nm (347 nm excitation).¹⁷ Fluorescence intensity was used to quantify 1-OHPG; the limit of detection was 0.03 pmol/ml urine. The coefficient of variation of the assay was 8%–10% during the period of sample analysis. A subset of samples was further purified by HPLC then by synchronous fluorescence spectroscopic analysis of individual fractions to confirm the identity of the fluorophore as already described.¹⁷

DATA ANALYSIS

Log transformation of urinary 1-OHPG concentrations normalised the frequency distribution in the exposed and control study groups. Log of 1-OHPG concentration was the outcome variable in all analyses. Summary data are presented as mean (SEM) and median (range). Group differences between exposed and controls, or smokers and non-smokers were tested with Student's *t* test. Pearson correlation was used to test the relation between urinary 1-OHPG concentration and stratum of PAH exposure or number of cigarettes smoked a day. Analysis of variance was used to test for overall differences in 1-OHPG concentrations among job categories. Duncan's multiple range test was used to compare differences between job categories. Multiple

Table 1 Polycyclic aromatic hydrocarbon job exposure categories

Job category	n	Exposure category*
Office	33	None
Construction or transport	7	Low
Continuous casting	5	Mid
Steel production	10	Mid
Iron production	12	Mid
Blast furnace	8	High
Coke oven	9	High

*Based on Bjørseth and Becher categorisation.³ Low = $< 0.01 \mu\text{g BaP}/\text{m}^3$; Mid = $0.01\text{--}1.0 \mu\text{g BaP}/\text{m}^3$; High = $> 1.0 \mu\text{g BaP}/\text{m}^3$.

linear regression was used to test for group differences between strata of PAH exposure or between smoker groups, to test for linear trends of each exposure variable, and to evaluate potential interaction and confounding. Two sided statistical tests were used throughout; P values < 0.05 were considered to be significant.

Results

Before analysing the samples, job categories were used to assign exposure to PAHs based on the published scheme of Bjørseth and Beche³ (table 1). The high PAH exposure stratum included blast furnace ($n = 8$) and coke plant ($n = 9$) workers; the mid PAH exposure stratum included continuous casting workers ($n = 5$), steel production furnace workers ($n = 10$), and iron production workers ($n = 12$) who mainly transported pig iron from the blast furnace to the steel making furnace; and the no or low PAH exposure stratum included office workers ($n = 33$) and construction or transportation workers ($n = 7$). Distribution of smoking was similar in the exposed and no or low exposure groups with

Table 2 Demographic characteristics of steel plant workers

Characteristic	No or low exposure group (n(%)) n = 40	Exposed group (n(%)) n = 44
Age:		
19–29	17 (42.5)	5 (11.4)
30–39	17 (42.5)	25 (56.8)
≥40	6 (15.0)	13 (29.5)
Smoking:		
Never or former	20 (50.0)	20 (45.5)
Current		
5–19/day	13 (32.5)	13 (29.5)
≥20/day	7 (17.5)	11 (25.0)
Job category		
Office	33 (82.5)	
Construction or transport	7 (17.5)	
Casting		5 (11.4)
Steel production		10 (22.7)
Iron production		12 (27.3)
Blast furnace		8 (18.2)
Coke plant		9 (20.4)

Based on job category.




Figure 1 Representative synchronous fluorescence spectrum of immunoaffinity-purified urine samples from an iron production worker and an office worker.

about 50% non-smokers in each group and about 20% one pack a day smokers (table 2). The age distribution between the exposed and no or low exposure group was different ($P = 0.008$ by ANOVA). Age was not found to be significantly associated with 1-OHPG in multiple regression analysis (see results).

Synchronous fluorescence spectroscopy analysis of immunoaffinity purified urine samples indicated a fluorescence excitation maximum (347 nm) characteristic of the pyrene moiety in 1-OHPG in 82 of the 84 samples analysed (fig 1 shows an example of detectable and non-detectable samples). The HPLC separation of a subset of 20 urine samples followed by synchronous fluorescence spectroscopic analysis of fluorescent fractions indicated that 98%–100% of the pyrene moiety fluorescence was due to 1-OHPG (data not shown). Non-detectable samples were assigned a value of 0.015 pmol/ml urine which is half the limit of detection of the assay.

Urinary 1-OHPG concentrations of subjects with jobs associated with exposure to combustion products were compared with the no or low exposure group (fig 2). The mean (SEM) 1-OHPG concentration in exposed workers (2.16 (0.42) pmol/ml, $n = 44$) was significantly greater than that of no or low exposure group (0.38 (0.05) pmol/ml, $n = 40$) ($P < 0.0001$ by t test). Also, the mean (SEM) 1-OHPG concentration in current smokers (1.82 (0.41) pmol/ml, $n = 44$) was significantly greater than that of non-smokers (0.75 (0.20) pmol/ml, $n = 40$, $P < 0.005$ by t test). We also compared urinary 1-OHPG concentrations of workers in each of the seven different job categories within the plant (fig 3). All but two workers in the blast furnace or coke plant categories had 1-OHPG concentrations higher than 2 pmol/ml, whereas all office





Figure 2 Box and whisker plot of urinary 1-OHPG concentrations by occupational PAH exposure group. Lowest whisker = 10th percentile; lower edge of box = 25th percentile; middle line of box = 50th percentile; upper edge of box = 75th percentile; highest whisker = 90th percentile of the data.

Figure 3 Box and whisker plot of urinary 1-OHPG concentrations by job category.



workers had concentrations lower than 1 pmol/ml. Overall, there was a significant difference in 1-OHPG concentrations between job categories ($P < 0.0001$, by ANOVA). Mean 1-OHPG concentrations in either blast furnace or coke oven workers were higher than mean concentrations in each of the other five job categories ($P < 0.05$ by Duncan's multiple range). Similarly, mean 1-OHPG concentration in the 33 office workers was lower than in each of the other six job categories, including the seven construction or transportation workers ($P < 0.05$ by Duncan's multiple range test).

To examine the dose-response relation between 1-OHPG and PAH exposure, we tested the correlation between strata of PAH exposure and 1-OHPG. A highly significant correlation between exposure strata (no or low, mid, or high) and 1-OHPG was found (Pearson $r = 0.65$, $P < 0.0001$). The combined effect of smoking and occupation on 1-OHPG concentrations was examined by

categorising the three occupational PAH exposure strata by smoking group (fig 4). Mean 1-OHPG concentrations in smokers were greater than for non-smokers in the same PAH exposure stratum (table 3). The absolute difference in 1-OHPG concentrations (2.07 pmol/ml) between smokers and non-smokers in the high PAH exposure group was greater than the difference in 1-OHPG concentrations between smokers and non-smokers in either the mid PAH exposure group (0.24 pmol/ml) or the no or low exposure group (0.25 pmol/ml). Although this is consistent with an interactive effect between smoking and occupational exposure to PAHs, the test for interaction by linear regression was not significant ($P = 0.30$). We had limited power to evaluate this interaction because 12 of 17 (71%) workers in the high PAH exposure stratum were current smokers, leaving only five non-smokers for comparison.

Multiple regression analysis indicated that both occupational exposure to PAHs and smoking were significant predictors of urinary 1-OHPG concentrations (table 4). The overall model r^2 was 0.55, indicating that 55% of the variation of 1-OHPG concentration was explained by the variables smoking and occupational exposure to PAHs. Similar results were obtained after adjustment for age or urinary creatinine.

Table 3 Urinary 1-OHPG concentration by PAH exposure strata

	PAH exposure strata		
	No or low (n = 40)	Mid (n = 27)	High (n = 17)
Smoking:			
Never or ex:			
n	20	15	5
mean (SEM) (pmol/ml)	0.26 (0.06)	0.70 (0.28)	2.84 (0.90)
Current:			
n	20	12	12
mean (SEM) (pmol/ml)	0.55 (0.07)	0.94 (0.42)	4.91 (1.02)

Figure 4 Box and whisker plot of urinary 1-OHPG concentrations stratified by smoking and PAH exposure strata.




Table 4 Predictors of urinary 1-OHPG by multiple linear regression

Variable	β (SE β)*	P value	Overall model (r^2)
Model A:			
Intercept	-1.29		0.51
PAH exposure1†	0.49 (0.23)	0.036	
PAH exposure2‡	2.43 (0.27)	<0.0001	
Model B:			
Intercept	-1.07		0.098
Smoking (yes or no)	0.81 (0.27)	0.0037	
Model C:			
Intercept	-1.55		0.55
PAH exposure1†	0.52 (0.22)	0.022	
PAH exposure2‡	2.32 (0.26)	<0.0001	
Smoking (Yes or no)	0.52 (0.20)	0.01	
Model C (creatinine adjusted):			
Intercept	-3.38		0.44
PAH exposure1†	0.62 (0.25)	0.015	
PAH exposure2‡	2.13 (0.29)	<0.0001	
Smoking (Yes or no)	0.46 (0.22)	0.04	

*SE β = standard error of β coefficient.

†PAH exposure1 = mid v no or low PAH exposure strata.

‡PAH exposure2 = high v no or low PAH exposure strata.

The dose-response relation of cigarette smoking with concentration of 1-OHPG was examined by testing the association between number of cigarettes smoked a day and 1-OHPG concentrations. Overall, there was a significant association between 1-OHPG concentration and number of cigarettes smoked (Pearson's $r = 0.28$, $P < 0.01$, $n = 82$; two missing, data not shown). When examined by PAH exposure strata, there was a significant association between concentration of 1-OHPG and number of cigarettes smoked in the no or low PAH exposure stratum (Pearson's $r = 0.44$, $P < 0.004$, $n = 40$) but not the combined mid and high PAH exposure stratum (Pearson's $r = 0.26$, $P = 0.09$, $n = 42$). To evaluate the sensitivity of the assay to detect moderate concentrations of smoking, we compared 1-OHPG in non-smokers and those who smoked ≤ 20 cigarettes a day (mean cigarettes a day = 15; range = 7–20) in the no PAH exposure stratum (office workers only). A significant difference in 1-OHPG concentrations was found between the 17 non-smokers (0.20 (0.04) pmol/ml) and the 15 moderate smokers (0.41 (0.05) pmol/ml, $P = 0.003$, by t test).

The association between 1-OHPG concentrations and dietary PAH exposure was assessed with data on consumption of foods containing high concentrations of PAHs from the dietary questionnaire. Among this population of workers, relatively small quantities of foods known to contain high PAH concentrations were consumed. Fifteen of 84 study subjects had consumed small amounts of grilled or broiled meat within the two weeks before sample collection; however, only five of these subjects had consumed it within 48 hours of collection. Thus we could only compare 1-OHPG concentrations between those workers with low dietary PAH exposure ($n = 15$) and those with no dietary PAH exposure ($n = 69$). No significant difference was found between these groups; therefore, diet did not seem to significantly contribute to urinary 1-OHPG in this population.

Discussion

We examined 1-OHPG concentration in urine from 44 workers with jobs in areas of increased PAH concentrations in air and 40 workers with jobs at the same site with low or no exposure to PAHs. Mean 1-OHPG concentration in exposed workers was significantly increased compared with unexposed controls. We also compared urinary 1-OHPG concentrations of workers in each of the seven different job categories within the plant. There was a significant difference in log transformed 1-OHPG concentrations between job categories. Mean 1-OHPG concentrations in the blast furnace and coke oven workers were significantly increased compared with workers in other job categories.

Assignment of exposure strata for PAHs based on job category was used to investigate the association of urinary 1-OHPG with PAH exposure. A clear dose-response relation was

found between mean 1-OHPG concentrations and no or low, mid, and high PAH exposure strata in both the current smokers and the non-smokers. The correlation between PAH exposure strata and 1-OHPG was highly significant (Pearson $r = 0.65$, $P < 0.0001$). Also, 1-OHPG concentration was correlated with quantity of cigarettes smoked among all workers or among no or low exposed workers. Thus, both smoking and occupational exposure to PAHs was associated with increased concentrations of 1-OHPG in urine from steel plant workers. This was verified by regression analysis.

We have chosen not to adjust our 1-OHPG values for urinary creatinine as this had little effect on the analyses or our conclusions. Also, deletion of samples with creatinine values < 4.0 $\mu\text{mol/ml}$ urine as suggested,⁹ left associations with occupational exposures unchanged and only slightly strengthened most associations with smoking. In previous studies of urinary 1-OHPG after dietary PAH exposure²⁴ or smoking (unpublished), we have found no improvement of associations between 1-OHPG and PAH exposure after adjustment for creatinine.

Although concurrent PAH exposure data were not available, job category was used to construct relative exposure strata. Measurements of benzene soluble fractions in personal air samples were available from one year before urine collection in the coke oven workers only. Mean concentrations of benzene soluble fractions were 0.24–0.49 mg/m^3 for three high exposure job titles in the coke oven battery. This range is below the published mean value of 2.08 mg/m^3 from personal samples that represents 11 job titles at 10 coke plants measured 25–30 years ago.²⁰ A more recent study reported mean values of 0.3 and 0.5 mg/m^3 (individual range: < 0.1 –5.7) among 16 highly exposed workers at two coke ovens.^{9, 21} The corresponding mean benzo(a)pyrene concentrations measured in those samples were 0.7 and 1.2 (individual range: 0.1–6.5) $\mu\text{g/m}^3$.

Assessment of urinary 1-hydroxypyrene has been applied in populations exposed to PAHs from different sources (occupation, therapy, and food). Recent reports of 1-hydroxypyrene in human urine indicate that early studies may have overestimated the amount of 1-hydroxypyrene.^{22, 23} Concentrations of 1-hydroxypyrene in human urine from people with similar occupational exposures vary widely in different studies. These results indicate that considerable variability occurs in these measurements when performed by different investigators at different worksites. The basis for this variability (laboratory, exposure, or biological) is unclear at the present time.

Although it is difficult to directly compare urinary 1-OHPG concentrations in this study with the values of other occupational exposure studies because 1-hydroxypyrene has been measured in most studies, the mean (range) urinary concentration of 1-OHPG adjusted for creatinine among non-smoking office workers in this study (0.05 (0.004–0.16)

$\mu\text{mol/mol}$ creatinine, $n = 17$) was comparable with 1-hydroxypyrene concentrations of non-smoking controls reported in several other studies: 0.17 (0.01–0.93) $\mu\text{mol/ml}$ creatinine, $n = 14^9$; 0.16 (0.10–0.22) $\mu\text{mol/mol}$ creatinine, $n = 20^{22}$; and 0.26 (0.02–0.66) $\mu\text{mol/mol}$ creatinine, $n = 52^{10}$. A more recent study reported lower concentrations of urinary 1-hydroxypyrene in 43 non-smoking controls: 0–0.01 $\mu\text{mol/mol}$ creatinine (10–90 percentiles).²³ The similarly low urinary 1-OHPG concentrations among non-smoking controls in our study might be due to low dietary consumption of PAHs. Amounts of foods ingested by this group of workers that contained PAHs were relatively low and did not contribute significantly to urinary 1-OHPG concentrations. Infrequent consumption of foods with a high PAH content, such as smoked or broiled foods, is characteristic of the Korean diet. Dietary information collected by questionnaire indicated that relatively little broiled, smoked, or grilled meat was consumed by the subjects in the present study. Furthermore, the mean urinary 1-OHPG concentration among non-smoking office workers in this study was comparable to the mean concentration in urine samples from workers on a controlled diet that contained very low concentrations of PAHs.²⁴ Mean 1-OHPG concentrations in non-smokers adjusted for creatinine were 0.05, 0.15, 0.37 (range of high PAH exposure group, 0.02–0.89) $\mu\text{mol/mol}$ creatinine for the no or low, mid, and high PAH exposure groups based on job category. The corresponding values in smokers were 0.06, 0.19, 0.82 (range of high PAH exposure group, 0.15–3.57) $\mu\text{mol/mol}$ creatinine, respectively. Mean 1-OHPG concentration in our group of smokers with high exposure to PAHs was similar to that found previously in bench side coke oven workers,²⁵ in smokers exposed to diesel exhaust,¹² in road pavers,¹² and in foundry workers.²⁶ The results in this study are, however, generally lower than the values reported in studies with moderate to high PAH exposure: in highly exposed coke oven workers,²⁵ in aluminum reduction plant workers,^{27,28} and in coal tar distillation plant workers.¹⁰ These differences may be due to a variety of factors including (a) differences in exposure concentrations between steel plants, (b) metabolic differences between the subjects studied, (c) differences in assay conditions and accuracy, or (d) bias associated with creatinine normalisation. Relative differences in 1-OHPG concentrations between controls and coke oven workers are comparable with those found in other studies. The 11-fold difference in urinary 1-OHPG concentrations found between office workers and coke plant workers is consistent with the results of Jongeneelen *et al*⁹ who showed a 10-fold difference between non-smoking controls and coke oven workers, and a sixfold difference between controls and coke oven workers who smoked.

In this study, smokers had consistently higher urinary 1-OHPG concentrations than

non-smokers. The amount of pyrene in one cigarette is 50–200 ng (230–917 pmol). Therefore, smoking 10 cigarettes a day would yield a maximum daily inhaled dose of about 2.3–9.2 nmol. The estimated increase in urinary 1-OHPG concentration from such a dose would be about 0.1 nmol/l urine, calculated from the linear regression model. This suggests that on average about 1%–4% of the pyrene inhaled from cigarette smoke is excreted as 1-OHPG in urine assuming a daily urinary volume of 1 litre.

In summary, there are several potential applications of an assay for urinary 1-OHPG. Firstly, although the proportion of pyrene in different working environments varies from 2% to 20% of the total PAHs, several studies have reported fairly constant ratios of pyrene to other PAHs in specific work environments such as coke plants^{9,25} or aluminum plants.²⁷ Therefore the measurement of 1-OHPG may be useful in assessment of exposure to total PAHs in these types of plants. Secondly, although excretion of 1-hydroxypyrene occurs fairly rapidly after exposure (6–35 h^{9,25}) and excretion of 1-OHPG declines to near baseline concentrations within 48 hours of the end of dietary exposure,²⁴ this metabolite could prove useful as a component of medical biomonitoring in the workplace. The advantages of simplicity and rapidity of the assay and the ready availability and non-invasiveness of specimen collection make the assay amenable to routine sampling. Thirdly, the assay could also be used to complement air monitoring data when evaluating the effectiveness of new interventions, engineering control devices, or personal protective equipment in the working environment.²⁹ Fourthly, these and other urinary PAH metabolites may prove to be useful in assessing human PAH exposure from multiple sources when different kinetics for inhaled or ingested PAHs must be considered in exposure assessment or when estimation of external exposure concentrations is difficult. The biological implications of increased urinary 1-OHPG concentrations should be explored in the future in prospective cohort studies that have banked urine samples.

We gratefully acknowledge the participation of the study subjects. This research was supported in part by DHHS grants P30-ES03819 and P01-ES06052.

- Grimmer G. Sources and occurrence of polycyclic aromatic hydrocarbons. In: Environmental carcinogens selected methods of analysis. Vol 3. Analysis of polycyclic aromatic hydrocarbons in environmental samples. *IARC Sci Publ* 1979;29:31–54.
- International Agency for Research on Cancer. *IARC monographs on the evaluation of the carcinogenic risk of chemicals to humans: polynuclear aromatic compounds, part 1. Volume 32*. Lyon: IARC, 1983.
- Bjørseth A, Becher G. PAH in work atmosphere: occurrence and determination. Boca Raton: CRC Press, 1986:137.
- Becher G, Bjørseth A. Determination of exposure to polycyclic aromatic hydrocarbons by analysis of human urine. *Cancer Lett* 1983;17:301–11.
- Venier F, Clonfero E, Cottica D, Gava C, Zordan M, Pozzoli L, *et al*. Mutagenic activity and polycyclic aromatic hydrocarbon levels in urine of workers exposed to coal tar pitch volatile in an anode plant. *Carcinogenesis* 1985;6:749–52.
- Haugen A, Becher G, Benestad C, Vahakangas K, Trivers GE, Newman MJ, *et al*. Determination of polycyclic

- aromatic hydrocarbons in the urine, benzo(a)pyrene diol epoxide-DNA adducts in lymphocyte DNA, and antibodies to the adducts in sera from coke oven workers exposed to measured amounts of polycyclic aromatic hydrocarbons in the work atmosphere. *Cancer Res* 1986; 46:4178-83.
- 7 Keimig SD, Kirby KW, Morgan DP, Keiser JE, Hubert TD. Identification of 1-hydroxypyrene as a major metabolite of pyrene in pig urine. *Xenobiotica* 1983;13: 415-20.
 - 8 Jongeneelen FJ, Bos RP, Grimmer G. Excretion of pyrene and hydroxypyrene in urine [letter]. *Cancer Lett* 1990;51: 175-9.
 - 9 Jongeneelen FJ, van Leeuwen FE, Oosterink S, Anzion RBM, van der Loop F, Bos RP, et al. Ambient and biological monitoring of coke oven workers: determinants of the internal dose of polycyclic aromatic hydrocarbons. *Br J Ind Med* 1990;47:454-61.
 - 10 Jongeneelen FJ, Anzion RBM, Scheepers PTJ, Bos RP, Henderson PTh, Nijenhuis EH, et al. 1-Hydroxypyrene in urine a biological indicator of exposure to polycyclic aromatic hydrocarbons in several work environments. *Ann Occup Hyg* 1988;32:35-43.
 - 11 Jongeneelen FJ, Scheepers PTJ, Groenendijk A, Van Aerts L, Anzion RBM, Bos RP, et al. Airborne concentration of polycyclic aromatic hydrocarbons among paving workers exposed to coal tar derived road tars. *Am Ind Hyg Assoc J* 1988;49:600-7.
 - 12 Jongeneelen FJ, Anzion RBM, Henderson PT. Determination of hydroxylated metabolites of polycyclic aromatic hydrocarbons in urine. *J Chromatogr* 1987;413:227-32.
 - 13 Jongeneelen FJ, Bos RP, Anzion RBM, Thews JLG, Henderson PT. Biological monitoring of polycyclic aromatic hydrocarbons: metabolites in urine. *Scand J Work Environ Health* 1986;12:137-43.
 - 14 Jongeneelen FJ, Leijdekkers CM, Bos RP, Theuvs LG, Henderson PT. Excretion of 3-hydroxy-benzo[a]pyrene and mutagenicity in rat urine after exposure to benzo[a]pyrene. *J Appl Toxicol* 1985;5:277-82.
 - 15 Jongeneelen FJ, Anzion RBM, Leijdekkers CM, Bos RP, Henderson PT. 1-Hydroxypyrene in human urine after exposure to coal tar and a coal tar derived product. *Int Arch Occup Environ Health* 1985;57:47-55.
 - 16 Clonfero E, Zordan M, Venier P, Paleologo M, Levis AG, Cottica D, et al. Biological monitoring of human exposure to coal tar. Urinary excretion of total polycyclic aromatic hydrocarbons, 1-hydroxypyrene and mutagens in psoriatic patients. *Int Arch Occup Environ Health* 1989; 61:363-8.
 - 17 Strickland PT, Kang DH, Bowman ED, Fitzwilliam A, Downing TE, Rothman N, et al. Identification of 1-hydroxypyrene-glucuronide as a major pyrene metabolite in human urine by synchronous fluorescence spectroscopy and gas chromatography/mass spectrometry. *Carcinogenesis* 1994;15:483-7.
 - 18 International Agency for Research on Cancer. *IARC monographs on the evaluation of the carcinogenic risk of chemicals to humans: polynuclear aromatic compounds, part 3. Vol 34.* Lyon: IARC, 1984:101-90.
 - 19 Santella RM, Lin CD, Cleveland WL, Weinstein IB. Monoclonal antibodies to DNA modified by a benzo[a]pyrene diol epoxide. *Carcinogenesis* 1984;5: 373-7.
 - 20 Fannick N, Gonshor LT, Shockley J Jr. Exposure to coal tar pitch volatiles at coke ovens. *Am Ind Hyg Assoc J* 1972;33:461-8.
 - 21 Jongeneelen FJ. Biological exposure limit for occupational exposure to coal tar pitch volatiles at coke ovens. *Int Arch Occup Environ Health* 1992; 63:511-6.
 - 22 Sherson D, Sigsgaard T, Overgaard E, Loft S, Poulsen HE, Jongeneelen FJ. Interaction of smoking, uptake of polycyclic aromatic hydrocarbons, and cytochrome P450IA2 activity among foundry workers. *Br J Ind Med* 1992;49:197-202.
 - 23 Omland O, Sherson D, Hansen AM, Sigsgaard T, Autrup H, Overgaard E. Exposure of iron foundry workers to polycyclic aromatic hydrocarbons: benzo(a)pyrene-albumin adducts and 1-hydroxypyrene as biomarkers for exposure. *Occup Environ Med* 1994;51:513-8.
 - 24 Kang DH, Rothman N, Poirier MC, Greenberg A, Hsu CH, Schwartz BS, et al. Inter-individual differences in the concentration of 1-hydroxypyrene-glucuronide in urine and polycyclic aromatic hydrocarbon-DNA adducts in peripheral white blood cells after charbroiled beef consumption. *Carcinogenesis* 1995;16:1079-85.
 - 25 Buchet JP, Gennart JP, Mercado-Calderon F, Delavignette JP, Cupers L, Lauwerys R. Evaluation of exposure to polycyclic aromatic hydrocarbons in a coke production and graphite electrode manufacturing plant: assessment of urinary excretion of 1-hydroxypyrene as a biological indicator of exposure. *Br J Ind Med* 1992;49: 761-8.
 - 26 Santella RM, Hemminki K, Tang D, Paik M, Ottman R, Young TL, et al. PAH-DNA adducts in white blood cells and urinary 1-hydroxypyrene in foundry workers. *Cancer Epidemiol Biomarkers Prev* 1993;2:59-62.
 - 27 Tolos WP, Shaw PB, Lowry LK, Mackenzie BA, Deng JF, Markel HL. 1-Pyrenol: a biomarker for occupational exposure to polycyclic aromatic hydrocarbons. *Appl Occup Environ Hyg* 1990;5:303-9.
 - 28 Van Schooten FJ, Jongeneelen FJ, Hillebrand MJX, van Leeuwen FE, de Looft AJA, Dijkmans APG, et al. Polycyclic aromatic hydrocarbon-DNA adducts in white blood cell DNA and 1-hydroxypyrene in the urine from aluminum workers: relation with job category and synergistic effect of smoking. *Cancer Epidemiol Biomarkers Prev* 1995;4:69-77.
 - 29 Van Rooij JGM, Van Lieshout MAV, Bodelier-Bade MM, Jongeneelen FJ. Effect of the reduction of skin contamination on the internal dose of creosote workers exposed to polycyclic aromatic hydrocarbons. *Scand J Work Environ Health* 1993;19:200-7.