

Cancer Incidence Among Massachusetts Firefighters, 1982-1986

Susan R. Sama, BSN, Terry R. Martin, MPH, Letitia K. Davis, ScD, and David Kriebel, ScD

Previous investigations of cancer among firefighters have been limited to mortality data and have yielded inconsistent results. Case-control analyses were conducted in the present surveillance study in order to examine associations between firefighting and cancer incidence in Massachusetts. Subjects were identified through the Massachusetts Cancer Registry files for 1982-1986. Exposure status (firefighting) was determined from the usual occupation reported to the Registry. Nine different cancer types were examined among the 315 reported white male firefighters. Two "unexposed" reference populations were used: policemen and statewide males. Standardized morbidity odds ratios (SMORs) were statistically significantly elevated for melanoma (SMOR = 292; 95% C.I. = 170-503) and bladder cancer (SMOR = 159; 95% C.I. = 102-250) among firefighters compared with the state as a whole. When policemen were used as the reference group, the bladder cancer excess persisted (SMOR = 211; 95% C.I. = 107-414) and non-Hodgkin's lymphoma was elevated (SMOR = 327; 95% C.I. = 119-898); the melanoma excess was largely reduced (SMOR = 138; 95% C.I. = 60-319) but remained elevated among those aged 55-74 years (SMOR = 513; 95% C.I. = 150-1,750). Small number excesses (not significant) were also observed for pancreatic cancer and leukemia compared with police.

Key words: occupational exposures, melanoma, bladder cancer, lymphoma, leukemia, pancreatic cancer

INTRODUCTION

Firefighting is a strenuous and often dangerous occupation. In addition to the obvious safety hazards, such as smoke inhalation, falls, and burns, firefighters are exposed to a variety of toxic substances. These include various carcinogens such as asbestos, benzene, and polycyclic aromatic hydrocarbons [Bendix, 1979; Brandt-Rauf et al., 1988]. Whether firefighters are at excess risk of cancer due to these exposures has yet to be determined.

Department of Work Environment, University of Lowell, Lowell, MA (S.R.S., D.K.).
Occupational Health Surveillance Program, Bureau of Health Statistics, Research and Evaluation, Massachusetts Department of Public Health, Boston (T.R.M., L.K.D.).
Work performed at the Massachusetts Department of Public Health, Boston.
Address reprint requests to Terry R. Martin, Bureau of Health Statistics, Research and Evaluation, Massachusetts Department of Public Health, 150 Tremont Street, 5th floor, Boston, MA 02111.
Accepted for publication January 15, 1990.

Previous investigations of cancer mortality of firefighters have yielded inconsistent results. Studies based on the occupational information reported on death certificates have demonstrated increases in several different types of cancer among firefighters: non-Hodgkin's lymphoma excluding lymphosarcoma [Dewar, 1987]; brain and nervous system and bladder cancer [Milham, 1983]; lymphosarcoma and lung cancer [Petersen and Milham, 1980]; digestive cancer [Kizer, 1987]; colorectal cancer [Berg and Howell, 1975]; and pancreatic, brain and nervous system cancers, and leukemia [Gallagher et al., 1986].

In several cohort mortality studies in which firefighter employment records were linked to death certificates, no excess cancer was found [Musk et al., 1978; Mastromatteo, 1959; Eliopoulos et al., 1984]. Others, however, have demonstrated increased mortality due to colon, bladder, and brain cancer [Vena and Fiedler, 1987]; leukemia and skin cancer [Feuer and Rosenman, 1986]; and male breast cancer (Friedman-Jimenez, personal communication).

To examine cancer patterns among firefighters in Massachusetts, the present surveillance study was undertaken using cancer incidence data collected by the Massachusetts Cancer Registry. Case-control analyses, using occupational information obtained by the Registry, were conducted to examine potential associations between firefighting and the occurrence of nine different cancers shown to be elevated in previous studies. Whereas earlier investigations of firefighters have examined cancer mortality, this surveillance study is based on cancer incidence data.

MATERIALS AND METHODS

Data

Subjects were identified through the Massachusetts Cancer Registry (MCR) maintained by the Massachusetts Department of Public Health (MDPH). The MCR is a statewide, mandatory tumor-reporting system, which began collecting data from hospitals and licensed clinics in 1982. Reportable diagnoses include all malignant neoplasms (with the exception of in situ tumors and basal and squamous cell carcinomas of the skin), benign tumors of the central nervous system, and neoplasms of uncertain behavior. Completeness of reporting has been estimated to be at least 95% based on comparison with national cancer rates [ACS, 1986].

For each newly diagnosed cancer case, hospitals are required to report name, date of birth, sex, smoking status, primary site, histology, and usual occupation and industry. Primary site and histology are coded at the hospitals according to the International Classification of Diseases for Oncology (ICD-O) system [World Health Organization, 1976]. Smoking status information is limited to cigarettes only, and cases are identified as current or former smokers or as having never smoked. Usual occupation and industry is defined as the longest job held and is reported as found in the medical record. Occupation and industry are coded at the MDPH according to the 1980 U.S. Bureau of the Census (BC) system [U.S. Bureau of the Census, 1982].

Analysis

The study period was defined as 1982–1986, the first 5 years for which complete MCR data were available. Male cancer cases with a reported usual occupation of firefighter (BC code 417) or fire chief (BC code 413) were included in the exposed

firefighter group. (There were no female cases reported as firefighters during the study period).

Two unexposed reference groups were identified from the male cases reported to the MCR during 1982–1986: 1) state—cases with any occupational information reported other than firefighter or fire chief; and 2) police—cases reported as policeman (BC code 418), police chief (BC code 414), sheriff (BC code 423), or correctional officer (BC code 424). Police were selected as a reference group because of their probable similarity to firefighters with respect to sociodemographic factors. Study subjects were limited to those aged 18 years or over at the time of diagnosis.

About one-third of the firefighters (100 cases) were reported as “fireman,” a job title that can refer to either a firefighter or a furnace operator. A review of secondary sources of information (hospital and union records, death certificates, and funeral directors) was conducted to determine which of these were firefighters. Six of the 82 cases for whom further information was available were found to be furnace operators and were excluded from the exposed group. The 18 cases who could not be verified were assumed to be firefighters and were included in the exposed group.

Standardized morbidity odds ratios (SMORs) were computed to measure associations between firefighting and specific cancer types. The SMOR in this study is calculated as the ratio of the odds of having the cancer of interest among firefighters to the odds among the “unexposed” reference population. The SMOR can be interpreted as the number of firefighters with the cancer of interest to the number expected based on the cancer odds of the reference group [Miettinen and Wang, 1981].

Age-adjusted SMORs for each of nine cancer types were calculated comparing firefighters to both reference groups—police and state—using six age categories (18–44, 45–54, 55–64, 65–74, 75–84, and 85+ years). The selection of cancer types for examination was based on previous evidence of associations with firefighting and sufficient numbers of observed cases ($n \geq 5$). These included cancers of the colon; rectum; pancreas; lung, bronchus, and trachea; melanoma of the skin; bladder; brain and other nervous system; non-Hodgkin’s lymphoma; and leukemia. “Control” cancers included all other cancers, except those of the organ systems of concern: digestive, respiratory, and lymphatic/hematopoietic, because of their potential association with firefighting.

Analyses were performed using the microcomputer program dEPID [Sullivan and Foster, 1988]. SMORs were calculated by standardizing to the age distribution of the exposed group (firefighters). Approximate 95% confidence intervals for the SMORs were calculated using a variance proposed by Rothman [1986].

RESULTS

There were 321 male cancer cases diagnosed in Massachusetts between 1982 and 1986 and reported to the MCR as firefighters. The distribution of firefighters, police, and statewide males, by age at diagnosis, race, and smoking status, are presented in Table I. Firefighters were similar to both reference groups with respect to age and smoking status. All firefighters with known race were white. All subsequent analyses were conducted limiting both the exposed and reference populations to whites.

SMORs for the nine cancer types examined among the 315 white male firefighters are presented in Table II. Statistically significant elevations were observed,

TABLE I. Distribution of Cancer Cases by Age, Race, and Smoking Status, Male Firefighters, Police, and State, Massachusetts, 1982–1986

Characteristic	Firefighters		Police		State ^a	
	No.	Percent	No.	Percent	No.	Percent
Total cases	321	100.0	392	100.0	29,277	100.0
Age						
18–64 years	166	51.7	230	58.7	14,310	48.9
65+ years	155	48.3	162	41.3	14,967	51.1
Race ^b						
White	315	100.0	377	99.0	27,827	97.1
Nonwhite	0	0.0	4	1.0	835	2.9
Unknown	6		11		615	
Smoking status ^c						
Never	71	24.9	98	29.3	6,782	26.8
Former	82	28.8	102	30.5	7,990	31.6
Current	132	46.3	134	40.1	10,499	41.6
Unknown	36		58		4,006	

^aOnly cases with known occupation are included.

^bCases with unknown race are not included in percent distribution.

^cCases with unknown smoking status are not included in percent distribution.

using the state reference group, for melanoma (SMOR = 292; 95% CI = 170–503) and bladder cancer (SMOR = 159; 95% CI = 102–250). When the police were used as the reference group, the bladder cancer excess persisted (SMOR = 211; 95% CI = 107–414), and non-Hodgkin's lymphoma was also elevated (SMOR = 327; 95% CI = 119–898); however, the melanoma excess was reduced (SMOR = 138; 95% CI = 60–319). SMORs greater than 200 were also seen for pancreatic cancer and leukemia but these findings were based on small numbers and were not statistically significant.

Table III presents age-specific SMORs for melanoma, bladder cancer, and lymphoma, computed using the police reference group. Although melanoma was only slightly elevated among all firefighters compared to police, there was a notable excess among those aged 55–74 years (SMOR = 513; 95% CI = 150–1,750).

DISCUSSION

In this study, we used Massachusetts Cancer Registry data for 1982–1986 to examine cancer incidence in firefighters compared with police and the state as a whole. This type of study is useful as a surveillance tool in order to generate leads about potential associations that can be followed-up with more in-depth etiologic research. Surveillance findings consistent with those of other studies are most worthy of further investigation.

Incidence data have several advantages over mortality data for studying cancer. Cancer registry information provides better diagnostic information than death certificates. Over 96% of the MCR cases are pathologically confirmed. Incidence data are

TABLE II. Standardized Morbidity Odds Ratios (SMORs) for Nine Cancer Types, White Male Firefighters Age 18+ Years, Massachusetts, 1982-1986

Cancer type (ICD-O code) ^a	Observed no.	Reference group			
		State		Police	
		SMOR	95% confidence interval	SMOR	95% confidence interval
Colon (153)	33	120	80-182	104	59-182
Rectum (154)	22	135	84-219	97	50-188
Pancreas (157)	6	98	42-226	319	72-1,415
Lung, bronchus, and trachea (162)	71	122	87-169	130	84-203
Melanoma of skin (173, H 872-879)	18	292	170-503	138	60-319
Bladder (188)	26	159	102-250	211	107-414
Brain and other nervous system (191-192)	5	86	34-215	152	39-592
Non-Hodgkin's lymphoma (H 959-964, 967-970, 972, 975-976)	14	159	89-284	327	119-898
Leukemia (H 980-994, excl. 984)	6	112	48-259	267	62-1,154

^aCode refers to primary site, except when preceded by H (for histology).

TABLE III. Age-Specific Standardized Morbidity Odds Ratios (SMORs) for Melanoma, Bladder Cancer, and Lymphoma, White Male Firefighters, Massachusetts, 1982-1986, Using White Police as the Reference Group

Age group (years)	Melanoma			Bladder cancer			Lymphoma		
	N	SMOR	95% confidence interval	N	SMOR	95% confidence interval	N	SMOR	95% confidence interval
18-54	5	55	16-196	4	125	26-588	2	132	16-1,072
55-74	11	513	150-1,750	18	219	99-484	10	538	150-1,924
75+	2	110	13-934	4	440	42-4,626	2	220	17-2,814

also more valuable for studying nonfatal cancers, such as bladder cancer, which are often not listed on death certificates of cases who die of other causes.

One disadvantage of this study is the underreporting of occupational information to the MCR. Because occupational information is available for only approximately 50% of all MCR cases, the actual number of cancer cases among firefighters may be up to twice as high as is reported here. The numerator-based analysis conducted in the present study could produce biased risk estimates if the reporting of firefighter varied by cancer site. It was not possible to evaluate reporting patterns without an additional source of information on the usual occupation of all cancer cases. However, there is little reason to suspect that reporting of firefighters would vary by cancer site.

A second limitation is that the MCR occupational information is obtained from medical records, and some misclassification of usual occupation is likely. Risk estimates based on this information may be biased in either direction if there is systematic

misclassification by cancer site. However, a recent study of Missouri Cancer Registry data [Brownson et al., 1989] found that reporting of usual occupation was fairly accurate (70%) and that misclassification was random, which would bias risk estimates towards the null.

Another limitation is that the MCR does not provide information about actual workplace exposures. Grouping all firefighters together, as in the current analysis, would tend to dilute the effects of exposure and bias risk estimates towards the null.

The present findings are consistent with previous reports of excess melanoma, bladder cancer, and lymphoma among firefighters. Excess melanoma mortality has been observed among New Jersey [Feuer and Rosenman, 1986] and Boston (Friedman-Jimenez, personal communication) firefighters.¹ As in our study, the excess among New Jersey firefighters was not notably elevated when police were used as the reference population. Police have been found to have excess skin cancer in several investigations [Feuer and Rosenman, 1986; Petersen and Milham, 1980]. One possible explanation for these findings is that the firefighter and police populations have relatively similar ethnic compositions. Ethnicity, or more specifically skin color, is a well-known risk factor for melanoma; individuals with fair skin are at increased risk [Elwood et al., 1984].

It is important to note, however, that firefighters were found to have a slight excess of melanoma in both the New Jersey study and ours, even when compared with police. In the present analysis, there was a marked excess in the 55–74-year-old group, in whom the observed incidence was five times greater than expected. This finding suggests that occupational exposures may contribute to melanoma incidence in firefighters. Carcinogens, including polycyclic aromatic hydrocarbons (PAHs) in soot, are potential causal agents. PAHs are produced from the combustion of organic materials [Bjorseth et al., 1978] and, hence, are present at all fires. Firefighters often have dermal contact with soot that penetrates their clothing.

Whatever the contributing factors are, firefighters do appear to be at increased risk. Establishment of melanoma screening for firefighters seems advisable, especially since melanoma is easy to detect and has a very high survival rate when detected early. Firefighter populations are readily identifiable and accessible, making them suitable for targeted screening [Koh et al., 1989].

Bladder cancer mortality has been shown to be increased among firefighters in Buffalo and Washington State [Vena and Fiedler, 1987; Milham, 1983]. One potential cause is exposure to PAHs, which have been suggested as a possible cause of bladder cancer [Silverman et al., 1986]. Excess bladder cancer has been observed previously among other workers exposed to combustion products, including chimney sweeps [Gustavsson et al., 1988] and aluminum smelter workers [Armstrong et al., 1986].

It is possible that the observed bladder cancer excess is related to cigarette smoking, if firefighters smoke more than the police or state reference groups. Based on the limited smoking data available, the proportions of current and former smokers in the three groups differ only slightly (Table I). Also, lung cancer, which is much more strongly correlated with smoking, is not notably elevated (SMOR = 130; 95%

¹None of the melanoma deaths in the Boston study overlapped with the melanoma cases in the present investigation.

CI = 84–203) among firefighters compared with police who do have a slightly lower reported smoking prevalence. Calculation of smoking-adjusted SMORs [Axelson, 1978], using the MCR smoking information, reduced the observed risk estimates for bladder cancer by only a small amount. Although the smoking data may not be completely accurate, we would not expect misclassification to differ for firefighters, police, and other occupational groups. Hence it is unlikely that the observed bladder cancer excess in firefighters is completely due to smoking.

Excess mortality from lymphoma has also been previously reported among firefighters [Dewar, 1987; Petersen and Milham, 1980]. There is mounting evidence that lymphoma, like leukemia, may be induced by exposure to benzene [Young, 1989]. Firefighters may be exposed to appreciable concentrations of benzene, which is present at most fires [Brandt-Rauf et al., 1988]. Benzene is vaporized from products such as gasoline, solvents, degreasing agents, and paint strippers and is a combustion product of polyvinyl chloride, polystyrene, and other plastics [Bendix, 1979]. Excess leukemia mortality [Feuer and Rosenman, 1986; Gallagher et al., 1986] and incidence [Morton and Marjanovic, 1984] among firefighters have been observed previously; we found excess incidence based on six cases (not significant). Further study examining the possible association between benzene exposure during fires and lymphoma and leukemia is necessary.

We did not observe excesses of brain, lung, colon, or rectal cancers as were seen in some other studies. This may in part be explained by differences between incidence and mortality data, methodological differences between the various studies, low power to detect weak associations, or random occurrence. Our findings of associations between firefighting and melanoma, lymphoma, and bladder cancer support previous observations and warrant further investigation to determine the potential causes. Melanoma screening for firefighters should be considered. Its advisability does not rest on proof that the observed excess risk is occupational in origin.

ACKNOWLEDGMENTS

This work was supported in part by a cooperative agreement with the National Institute for Occupational Safety and Health (NIOSH) (grant U60/CCU100816).

REFERENCES

- American Cancer Society (1986): "1986 Cancer Facts and Figures." New York: ACS.
- Armstrong BG, Tremblay CG, Cyr D, Thériault GP (1986): Estimating the relationship between exposure to tar volatiles and the incidence of bladder cancer in aluminum smelter workers. *Scand J Work Environ Health* 12:486–493.
- Axelson O (1978): Aspects on confounding in occupational health. *Scand J Work Environ Health* 4: 85–89.
- Bendix S (1979): Firefighter exposure to environmental carcinogens. *J Combust Toxicol* 6:127–135.
- Berg JW, Howell MA (1975): Occupation and bowel cancer. *J Toxicol Environ Health* 1:75–89.
- Bjorseth A, Bjorseth O, Fjeldstad PE (1978): Polycyclic aromatic hydrocarbons in the work atmosphere: I. Determination in an aluminum reduction plant. *Scand J Work Environ Health* 4:212–223.
- Brandt-Rauf PW, Fallon LF Jr, Tarantini T, Idema C, Andrews L (1988): Health hazards of fire fighters: exposure assessment. *Br J Ind Med* 45:606–612.
- Brownson RC, Davis JR, Chang JC, DiLorenzo TM, Keefe TJ, Bagby JR (1989): A study of the accuracy of cancer risk factor information reported to a central registry compared with that obtained by interview. *Am J Epidemiol* 129:616–624.

- Dewar RF (1987): "Mortality Experience of Pennsylvania Workers 1983–1985." Harrisburg, PA: Pennsylvania Department of Health.
- Eliopoulos E, Armstrong BK, Spickett JT, Heyworth F (1984): Mortality of firefighters in Western Australia. *Br J Ind Med* 41:183–187.
- Elwood JM, Gallagher RP, Hill GB, Spinelli JJ, Pearson JC, Threlfall WJ (1984): Pigmentation and skin reaction to sun as risk factors for cutaneous melanoma: Western Canada Melanoma Study. *Br Med J* 288:99–102.
- Feuer E, Rosenman K (1986): Mortality in police and firefighters in New Jersey. *Am J Ind Med* 9:517–527.
- Gallagher RP, Threlfall WJ, Band PR, Spinelli JJ, Coldman AJ (1986): "Occupational Mortality in British Columbia 1950–1978." Ottawa: Statistics Canada.
- Gustavsson P, Gustavsson A, Hogstedt C (1988): Excess of cancer in Swedish chimney sweeps. *Br J Ind Med* 45:777–781.
- Kizer KW (1987): "California Occupational Mortality, 1979–1981." Sacramento, CA: California Department of Health Services.
- Koh HK, Lew RA, Prout MN (1989): Screening for melanoma/skin cancer. Theoretic and practical considerations. *J Am Acad Dermatol* 20:159–172.
- Mastromatteo E (1959): Mortality in city firemen, II. A study of mortality in firemen of a city fire department. *AMA Arch Ind Health* 20:227–233.
- Miettinen OS, Wang J-D (1981): An alternative to the proportionate mortality ratio. *Am J Epidemiol* 114:144–148.
- Milham S Jr (1983): "Occupational Mortality in Washington State, 1950–1979." DHHS (NIOSH) Publication No. 83-116. Washington, DC: U.S. Government Printing Office.
- Morton W, Marjanovic D (1984): Leukemia incidence by occupation in the Portland-Vancouver metropolitan area. *Am J Ind Med* 6:185–205.
- Musk AW, Monson RR, Peters JM, Peters RK (1978): Mortality among Boston firefighters, 1915–1975. *Br J Ind Med* 35:104–108.
- Petersen GR, Milham S Jr (1980): "Occupational Mortality in the State of California, 1959–1961." Cincinnati: NIOSH.
- Rothman KJ (1986): "Modern Epidemiology." Boston: Little, Brown and Company, p 232.
- Silverman DT, Hoover RN, Mason TJ, Swanson GM (1986): Motor exhaust-related occupations and bladder cancer. *Cancer Res* 46:2113–2116.
- Sullivan KM, Foster DA (1988): "dEPID: A Program for Stratified and Standardized Analysis. Version 2.12." Atlanta: Centers for Disease Control.
- U.S. Bureau of the Census (1982): "1980 Census of Population Alphabetical Index of Industries and Occupations." Washington, DC: U.S. Government Printing Office.
- Vena JE, Fiedler RC (1987): Mortality of a municipal-worker cohort: IV. Firefighters. *Am J Ind Med* 11:671–684.
- World Health Organization (1976): "International Classification of Diseases for Oncology." Geneva: WHO.
- Young N (1989): Benzene and lymphoma. *Am J Ind Med* 15:495–498.