

# Subclinical Measures of Cardiovascular Health Among Wildland Firefighters

Md Zahid Hasan, MS, Erin O. Semmens, PhD, Kathleen Navarro DuBose, PhD,  
L. Kaili McCray, PhD, and Curtis W. Noonan, PhD

**Objective:** The aim of the study is to compare subclinical measures of cardiovascular health among wildland firefighters (WFFs) to the US general population. **Methods:** Our cross-sectional study compared body mass index, total cholesterol, and blood pressure in 11,051 WFFs aged 17 to 64 years using Department of the Interior Medical Screening Program clinical screening examinations between 2014–2018 to National Health and Nutrition Examination Survey of 2015–2016 cycle using adjusted logistic regression analyses. **Results:** The logistic regression model shows significantly higher odds of hypertension and prehypertension in WFFs (2.84 times more with 95% CI: 2.28–3.53) than US general population. There were no consistent differences in body mass index or total cholesterol between the two population. **Conclusions:** Hypertension and prehypertension were more prevalent in WFFs compared with the US general population, which suggests the need for actions for protecting against cardiovascular disease among WFFs.

**Keywords:** wildland firefighters, hypertension, body mass index, total cholesterol, blood pressure

The area burned by wildfires each year has grown more than five-fold in the United States (US) from 1983 to 2021.<sup>1</sup> In 2021 alone, approximately 59,000 wildfires burned a total area of 7.1 million acres.<sup>2</sup> As the climate continues to warm due to global warming, and due to widespread drought across the western US,<sup>3</sup> the frequency and impact of wildfires will continue to grow.<sup>4,5</sup> This increase in fire affects the people living in communities near where wildfires occur and wildland firefighters (WFFs) who are directly or indirectly involved in suppressing wildland fires.<sup>6,7</sup> The common workplace hazard of WFFs include smoke, arduous duty, high physical stress, sleep disturbance and deprivation, shift work, heat exposure, and dehydration.<sup>8–11</sup> Approximately 19,000 federal WFFs worked under the Department of Agriculture's Forest service and four agencies of Department of Interior (DOI).<sup>12</sup>

From the Center for Population Health Research, School of Public and Community Health Sciences University of Montana, Missoula, MT (M.Z.H., E.O.S., C.W.N.); Department of the Interior, Office of Wildland Fire, Boise, ID (K.N.D., L.K.M.).  
Md. Zahid Hasan, 0009-0002-5751-2050  
Erin O. Semmens, 0000-0003-0531-7659  
Kathleen Navarro DuBose, 0000-0002-2866-2087  
L. Kaili McCray, 0009-0005-8168-1921  
Curtis W. Noonan, 0000-0003-1050-4205

Funding sources: The study was supported by the National Institute for Occupational Safety and Health (R21OH011385 to E.O.S. and C.W.N.) and National Institutes of Health, National Institute for General Medical Sciences (P20GM130418 to C.W.N.).

Conflict of interest: None declared.

Disclaimer: The findings and conclusions in this report are those of the author(s) and should not be construed to represent any official Department of the Interior or US Government determination or policy.

Ethical considerations: This study was approved by the University of Montana's Institutional Review Board.

Authors' contribution: All authors contributed to manuscript writing. Study design was led by E.O.S., C.W.N., and M.Z.H. Primary data analysis was conducted by M.Z.H. The data underlying this article will be shared on reasonable request to the corresponding author.

Reporting follows STROBE guidelines.

Supplemental digital contents are available for this article. Direct URL citation appears in the printed text and is provided in the HTML and PDF versions of this article on the journal's Web site ([www.joem.org](http://www.joem.org)).

Address correspondence to: Curtis W. Noonan, PhD, Center for Population Health Research, University of Montana, 32 Campus Drive, Missoula, MT 59812 ([curtis.noonan@umontana.edu](mailto:curtis.noonan@umontana.edu)).

Copyright © 2024 American College of Occupational and Environmental Medicine  
DOI: 10.1097/JOM.0000000000000341

## LEARNING OUTCOMES

After reading this article, the audience will be able

- Describe a way to compare cardiovascular health among WFFs and a sample from the US general population.
- Evaluate the prevalence of hypertension among WFFs compared with the US general population.
- Understand that subclinical measures can be used to better describe acute and long-term cardiovascular disease risk among this worker population.

Wildland firefighting is a physically demanding, and young and healthy individuals tend to self-select into this arduous duty occupation. Nevertheless, WFFs are at risk of stroke, myocardial infarction, and thermal stress.<sup>13</sup> The National Wildfire Coordinating Group<sup>14</sup> reports that cardiovascular disease was the largest contributor to sudden deaths in WFFs from 2007 to 2016. Beyond these acute outcomes, WFFs may be increased long-term risk of cardiovascular disease. Applying estimated WFF PM<sub>2.5</sub> (particulate matter in the air, which is less than 2.5 micron in aerodynamic diameter) exposures to epidemiologically based risk estimates for PM<sub>2.5</sub> and lifetime cardiovascular mortality, WFFs may be at a 16%–30% increased risk of lifetime cardiovascular mortality.<sup>15</sup> Few studies have measured subclinical cardiovascular risk measures among WFFs, and underlying disease risk may better inform both acute and long-term cardiovascular disease risk estimates in this population.

We present here a cross-sectional study to test the hypothesis that cardiovascular risk measures (body mass index, total cholesterol, and systolic and diastolic blood pressure) among WFFs are different from those observed in a representative sample of the US general population. Body mass index, total cholesterol, and blood pressures are three important indicators of cardiovascular health.<sup>16</sup> The risks of cardiovascular diseases such as vascular damages, hypertension, coronary artery disease, congestive heart failure, stroke, arrhythmias, and sudden cardiac death have direct linkage with the high BMI or obesity.<sup>17</sup> Cholesterol is a crucial component for physiological function, but high cholesterol level is adverse to cardiovascular health and can lead to atherosclerosis.<sup>18</sup> Elevated blood pressure is related to increased risk for stroke, atrial fibrillation, and heart failure.<sup>19</sup> A better understanding of the status of subclinical indicators/risks of cardiovascular in WFFs may inform actionable, preventive health efforts to reduce risk of cardiovascular morbidity and mortality in this high-risk worker population. See Strengthening the Reporting of Observational studies in Epidemiology (STROBE) checklist in Supplementary Digital Content (SDC, <http://links.lww.com/JOM/B492>) file.

## METHODS

### Study Population

The study is based on clinical and survey data from a comprehensive medical screening program conducted by the Department of the Interior Wildland Fire Medical Standards Program (DOI MSP)

between 2014 and 2018.<sup>20</sup> The DOI WFFs are federal WFFs employed by Bureau of Indian Affairs, Bureau of Land Management, National Park Service, and US Fish and Wildlife Service.

The DOI MSP is conducted in various health facilities across the US by trained health personnel. Although DOI MSP is conducted at different health facilities across the US, a minimum standard protocol is maintained. The DOI WFFs complete DOI MSP screening examinations at entry to employment and at 3-year intervals thereafter to maintain status as an arduously qualified WFF. The purpose of the medical screening program is to verify that WFFs are free from medical conditions that would put them at risk for acute adverse health outcomes in the work setting. In addition, every DOI WFF must pass the work capacity test to meet a minimum fitness requirement for safe and efficient job performance. This dataset includes 11,051 WFFs between the ages of 17 and less than 65 years.<sup>20</sup>

To compare the cardiovascular health of WFFs with the US general population, we used the 2015–2016 National Health and Nutrition Examination Survey (NHANES). The NHANES is an extensive cross-sectional health and nutritional survey, conducted by the Centers for Disease Control and Prevention, National Center for Health Statistics. The NHANES is a complex probabilistic survey that is representative of the noninstitutionalized population of the United States. Notably, the NHANES collects data across the US without exclusions based on age, sex, or race/ethnicity. Similar to the DOI MSP screening program data, the NHANES includes data from a physical examination at a clinical examination center and augments clinical data with health history and health behavior information ascertained through interview. To align with the DOI MSP dataset, we eliminated children younger than 17 years from the NHANES dataset as well as adults 65 years and older. In total, the 2015–16 dataset included 4771 participants, ages of 17 to less than 65 years. To facilitate statistical comparison across populations, we created a new variable, “population,” with two categories, “WFF,” and “US general population.”

## Measures

To understand cardiovascular health status of WFFs and the US general population, we evaluated three clinical measures (body mass index [BMI], total cholesterol, and systolic and diastolic blood pressure) and two self-reported measures (smoking and diabetes status). Body mass index for WFFs was calculated from height and weight measures; BMI was reported directly in NHANES. We categorized BMI into obese ( $\geq 30$  kg/m<sup>2</sup>), overweight (25–29.9 kg/m<sup>2</sup>), and normal weight ( $< 25$  kg/m<sup>2</sup>).<sup>16</sup> Both datasets used a single measurement of total cholesterol (more information is added in additional materials section of this article). Cholesterol and blood pressure measures were also summarized by risk categories according to Lloyd-Jones et al.<sup>13</sup> We categorized total cholesterol into hypercholesterolemia ( $\geq 240$  mg/dL), borderline high (200–239 mg/dL), and normal level ( $< 200$  mg/dL). Single values for systolic blood pressure (SBP) and diastolic blood pressure (DBP) were reported for the MSP, whereas we averaged across three SBP and DBP measures for the NHANES population. We categorized blood pressure into hypertension (SBP  $\geq 140$  mm Hg OR DBP  $\geq 90$  mm Hg), prehypertension (SBP 120–139 mm Hg and/or DBP 80–89 mm Hg), normal blood pressure (SBP  $< 120$  mm Hg and DBP  $< 80$  mm Hg).<sup>16</sup> Data for these clinical measures were censored for out of range values according to Centers for Disease Control and Prevention guidelines and NHANES.<sup>21–23</sup> See also SDC for data cleaning procedures.

Smoking status among WFFs was based on self-reporting during their medical examinations, asking whether they smoke cigarettes, e-cigarettes, cigars, or any other tobacco products. Smoking status for NHANES participants was based on several questions related to smoking such as number of cigarettes they smoked within particular durations. The WFFs were asked whether they ever had diabetes or not, and NHANES participants were asked whether they were told by a doctor or health professional that they had diabetes other than during pregnancy.

## Covariates

To enable comparison of cardiovascular variables between WFFs and the US general population, we described both datasets (DOI MSP and NHANES) based on age, sex, and race. For age, we used the following categories: 17 to 29 years, 30 to 39 years, 40 to 49 years, and 50 to less than 65 years. For sex, we used male and female. Race was categorized as non-Hispanic White, non-Hispanic African, Hispanic, Asian, and others.

## Statistical Analysis

To compare the prevalence of cardiovascular health risk categories, we used age standardization based on four categories (17–29, 30–39, 40–49, 50–65 years). We also compared these two populations to understand whether they differed by sex or race using a chi-squared test. We presented the mean, median, minimum, 25th percentile, 75th percentile, and maximum value for continuous variables. For age-standardized continuous measures, we compared population means using *t* tests, and for categorical variables, we compared age-standardized populations using chi-squared tests. Finally, we used regression to compare measures between populations while accounting for other covariates such as sex and race. For all analyses with NHANES data, we employed survey weighting to account for the complex survey design. The goal of survey weighting is to increase the representativeness of NHANES dataset to the US general population using primary sampling unit (psu), 2 years sample weight (persWeight or WTINT2YR), and masked variance unit (strata). We used survey command (svy in STATA) to calculate the corresponding number of US general population for demographic and cardiovascular factors. We used logistic regression to assess differences between WFFs and the general population for dichotomous outcomes. Regression models were adjusted by age, sex, and race. Covariates (age, sex, and race) were selected based on their hypothesized relationships with both occupational history and cardiovascular health. For the analysis, we used R version 4.1.0 (The R Foundation for Statistical Computing, Austria) and STATA version 15.1 (StataCorp, College Station, TX).

## RESULTS

Table 1 summarizes the demographic characteristics of WFFs compared with the weighted NHANES sample. Wildland firefighters in the age groups 17 to less than 30 years and 30 to less than 40 years

**TABLE 1.** Distribution of WFFs and US General Population by their Demographic Characteristics

Demographic Characteristics	WFFs, n (%)	Weighted NHANES, %	P
Race			<0.001
Non-Hispanic African	81 (0.7%)	12.5%	
Asian	61 (0.6%)	6.2%	
Non-Hispanic White	6,810 (61.6%)	60.1%	
Hispanic	573 (5.2%)	17.6%	
Not available	632 (5.7%)	—	
Others	2,894 (26.2%)	3.7%	
Total	11,051 (100%)	100%	
Gender			<0.001
Female	652 (5.9%)	50.8%	
Male	9,425 (85.3%)	49.2%	
Not available	974 (8.8%)	—	
Total	11,051 (100%)	100%	
Age, yr			<0.001
17–29.99	4,214 (38.1%)	27.3%	
30–39.99	3,623 (32.8%)	20.9%	
40–49.99	2,049 (18.5%)	20.9%	
50–64.99	1,165 (10.5%)	30.9%	
Total	11,051 (100%)	100%	

NHANES, National Health and Nutrition Examination Survey; WFF, wildland firefighter.

TABLE 2. Summary Statistics for Cardiovascular Variables Among WFFs and NHANES Samples

Variables	Statistics	WFFs Between 2014–2018	NHANES US Population 2015–2016	Weighted NHANES US Population 2015–2016	P (T Test or $\chi^2$ )
BMI, kg/m <sup>2</sup>	n Mean, SD	10,382	4,535	187,698,660	<0.001
	Min, 25th percentile, median,	27.8 ± 5.1	29.3 ± 7.4	29.3 ± 0.3	
	75th percentile, max	8.5, 24.3, 27.2, 30.6, 85.9	14.5, 23.9, 28.1, 33.2, 67.3		
Total cholesterol, mg/dL	n Mean, SD	10,810	4,315	180,327,896	0.004
	Min, 25th percentile, median,	184.4 ± 37.9	188.9 ± 41.4	191.3 ± 1.6	
	75th percentile, max	38, 158, 181, 207, 486	77, 160, 185, 213, 545		
Systolic blood pressure, mm Hg	n Mean, SD	10,804	4,236	176,479,728	<0.001
	Min, 25th percentile, median,	123.0 ± 13.2	120.9 ± 16.0	120.0 ± 0.4	
	75th percentile, max	65, 114, 122, 130, 224	85, 110, 119, 129, 231		
Diastolic blood pressure, mm Hg	n Mean, SD	10,804	4,236	176,479,728	<0.001
	Min, 25th percentile, median,	77.1 ± 10.2	70.5 ± 11.3	70.9 ± 0.5	
	75th percentile, max	2, 70, 78, 84, 149	11, 63, 71, 77, 117		
Hypertension	Total n (%)	10,804 (6.2%)	4,236 (7.6%)	176,479,728 (5.8%)	<0.001*
Self-reported diabetes	Total n (%)	4,763 (0.9%)	4,768 (9.8%)	196,766,306 (7.8%)	<0.001*
Current smoking	Total n (%)	4,582 (16.2%)	4,618 (22.2%)	192,797,119 (21.9%)	<0.001*

Asterisk (\*) mark represents chi-squared test.  
NHANES, National Health and Nutrition Examination Survey; WFF, wildland firefighter.

were considerably overrepresented (38% and 33% in WFF compared with 27% and 21% in NHANES). The WFFs had few individuals (10%) aged 50 to less than 65, compared with approximately one third of the NHANES sample in this age range. Wildland firefighters were more likely to be male (85%–49%,  $P < 0.001$ ). The two populations had similar proportions reporting race/ethnicity as non-Hispanic White (61.6% and 60.1%). Compared with the NHANES sample, lower proportions of WFFs reported race/ethnicity as non-Hispanic African American, non-Hispanic Asian or Hispanic, but a large proportion of WFFs (31.9%) could not be classified into single race/ethnicity groups or were missing information on race/ethnicity. The

chi-squared test suggests that the composition of race and sex are different in WFFs than NHANES population.

Valid measures for BMI, cholesterol, and blood pressure were captured for 10,382 to 10,810 of the 11,051 MSP participants and for 4236 to 4535 of the 4771 NHANES participants (see SDC Figs. S1 and S2, <http://links.lww.com/JOM/B492>, respectively). Table 2 summarizes statistics for five measures for WFFs compared with the US general population. Mean BMI (29.3 vs 27.8 kg/m<sup>2</sup>,  $P < 0.001$ ) and total cholesterol (191.3 vs 184.4 mg/dL,  $P = 0.001$ ) were higher in the NHANES sample relative to WFFs. Wildland firefighters had a higher average SBP and DBP than the NHANES sample. The

TABLE 3. Percentage of People in the High-Risk Categories of Each Cardiovascular Risk Variable

Variables	WFFs		NHANES US Population 2015–2016
	No. WFFs, n	Overall % (95% CI); (Male %, Female %)	Overall % (95% CI); (Male %, Female %)
BMI, kg/m <sup>2</sup>	10,382	28.9% (27.9%–29.9%) (M = 29.9%, F = 17.3%)	38.24% (34.7%–42%) (M = 36.7, F = 39.7%)
Systolic blood pressure, mm Hg	10,804	7.4% (6.8%–8.1%)	5.8% (5.0%–6.8%)
Diastolic blood pressure, mm Hg	10,804	(M = 7.9%, F = 3.4%)	(M = 7.2%, F = 4.6%)
Total cholesterol, mg/dL	10,810	9.8% (9.1%–10.5%) (M = 10%, F = 7.4%)	12.4% (10.5%–14.5%) (M = 12.4%, F = 12.4%)
Current smoking	4,582	15.6% (14.3%–16.9%) (M = 16.7%, F = 5.5%)	21.8% (19.7%–24.1%) (M = 24.4%, F = 19.3%)
Self-reported diabetes	4,763	1.6% (1.1%–2.3%) (M = 1.7%, F = 0.7%)	7.8% (6.8%–8.9%) (M = 8.4%, F = 7.1%)
At least one high risk category among 3 risk categories (obesity, hypertension, hypercholesterolemia)	9,996	39% (37.8%–40.2%) (M = 40.4%, F = 27.6%)	47.7% (50.2%–45.2%) (M = 46.8%, F = 48.5%)

F, female; M, male; NHANES, National Health and Nutrition Examination Survey; WFF, wildland firefighter.  
\*High risk categories: BMI ≥ 30 kg/m<sup>2</sup>, obese; blood pressure: SBP ≥140 mm Hg or DBP ≥90 mm Hg, hypertension; total cholesterol: ≥240 mg/dL, hypercholesterolemia; smoking: current smoker. Percentages are age standardized.

**TABLE 4.** The Influence of Wildland Firefighter Occupational Status, Relative to the General Population, on Cardiovascular Risk Factors

Measure	Category	Population Sample, n (%)		OR*	
		WFF	NHANES	Crude OR	Adjusted OR (95% CI)
BMI	Normal weight	3,378	60,574,176	Ref.	Ref.
	Obesity + overweight	7,004	127,124,484	1.0	1.1 (0.87–1.30)
Cholesterol	Normal	7,299	109,363,603	Ref.	Ref.
	Hypercholesterolemia + moderately high cholesterol	3,511	70,964,293	0.7	1.0 (0.70–1.30)
Blood pressure	Normal blood pressure	3,457	92,554,912	Ref.	Ref.
	Hypertension + prehypertension	7,347	83,924,816	2.3	2.8 (2.3–3.5)

\*Logistic regression analyses adjusted for age, race/ethnicity, and sex.

\*Body mass index  $\geq 30$  kg/m<sup>2</sup>, obese and BMI: 25.0–29.9 kg/m<sup>2</sup>, overweight; total cholesterol:  $\geq 240$  mg/dL, hypercholesterolemia; total cholesterol: 200–239 mg/dL, borderline high cholesterol; blood pressure: SBP  $\geq 140$  mm Hg or DBP  $\geq 90$  mm Hg, hypertension; blood pressure: SBP 120–139 mm Hg and/or DBP 80–89 mm Hg, prehypertension.

BMI, body mass index; CI, confidence interval; NHANES, National Health and Nutrition Examination Survey; OR, odds ratio; WFF, wildland firefighter.

proportion with indication for hypertension was higher in WFFs than NHANES (6.2% vs 5.8%,  $P < 0.001$ ). Diabetes was reported among 8% of NHANES population compared with less than 1% of WFFs. Current smoking was reported among 22% of NHANES population compared with 16% of WFFs.

Table 3 presents percentages of WFFs in the high-risk categories of each cardiovascular risk variable. The percentage of BMI values greater than or equal to 30 kg/m<sup>2</sup> (indicating obesity) was higher (38.2% [95% CI: 34.7%–42%]) in the NHANES population compared with WFFs (28.9% [95% CI: 27.9%–29.9%]). The prevalence of hypercholesterolemia, cholesterol values greater than or equal to 240 mg/dL, was similar in the WFF and NHANES samples. Self-reported smoking and diabetes were more common among US general population compared with WFFs. In contrast with the other measures, high blood pressure was more common among WFFs. The proportion of WFFs with indication of hypertension (SBP  $\geq 140$  mm Hg or DBP  $\geq 90$  mm Hg) was higher among WFFs (7.4% [95% CI: 6.8%–8.1%]) compared with NHANES participants (5.8% [95% CI: 5.0%–6.8%]). The categorization of one or more variables as high risk indicates that at least one high-risk factor such as obesity, hypertension, and hypercholesterolemia affected 47.7% (95% CI: 45.2%–50.2%) of the US general population compared with 39% (95% CI: 37.8%–40.2%) of WFFs. Obesity, hypercholesterolemia, hypertension, current smoker, and self-reported diabetes are lower among female WFFs than US general population and male WFFs.


Table 4 shows the regression results for risk categories of BMI, cholesterol, and blood pressure adjusting for age, sex, and race. The logistic regression model for BMI (Table 4) shows that there is no significant difference of risks of having obesity or overweight between

the WFFs and NHANES groups. Similarly, the logistic regression model for total cholesterol (Table 4) shows that there is no significant difference of risks of having hypercholesterolemia or borderline high cholesterol between WFFs and US general population. The logistic regression model for blood pressure (Table 4) shows that WFFs have 2.8 times more risks of having hypertension or prehypertension (95% CI: 2.3, 3.5) than US general population.

Based on the three cardiovascular health risks (obesity/overweight, hypercholesterolemia/borderline cholesterol, hypertension/prehypertension), the prevalence of meeting more than 1 risk category is higher in WFFs than US general population, although this was primarily evident among males (Fig. 1).

## DISCUSSION

Given the fitness requirements for arduous duty firefighting, we expected that WFFs would be a particularly healthy subset of the general population. Wildland firefighters must annually pass an aerobic performance test, known as the arduous pack test, which includes a walking tolerability test with weights.<sup>9</sup> In addition to these tests, some WFFs, for example, smokejumpers and hotshots, need to pass more extensive physical tests to make sure they have the strength and endurance necessary to carry out their particular jobs.<sup>9</sup> However, in a large sample of federal WFFs, we found that, relative to the US general population, WFFs had a higher prevalence of hypertension and prehypertension. We observed no difference between WFF and NHANES with respect to overweight/obese and high cholesterol status. In addition, compared with the US male general population, a greater proportion of male WFFs had more than one cardiovascular risk factor.



**FIGURE 1.** Comparison of age-standardized prevalence (%) of three cardiovascular risks among WFFs and the survey-weighted NHANES sample (risk here consists of poor and intermediate categories of BMI, blood pressure and cholesterol).<sup>a</sup> \* mark represents statistically significant difference based on chi-squared test of risks (no risk group, 1 risk group, and more than 1 risk group) between NHANES sample and WFFs. <sup>a</sup> BMI  $\geq 30$  kg/m<sup>2</sup>, obese and BMI: 25.0–29.9 kg/m<sup>2</sup>, overweight; total cholesterol:  $\geq 240$  mg/dL, hypercholesterolemia; total cholesterol: 200–239 mg/dL, borderline high cholesterol; Blood Pressure: SBP  $\geq 140$  mm Hg or DBP  $\geq 90$  mm Hg, hypertension; blood pressure: SBP 120–139 mm Hg and/or DBP 80–89 mm Hg, prehypertension

A variety of job-related factors may put WFFs at greater risk of poor cardiovascular health. For example, the adverse conditions include fatigue, sleep deprivation, and increased mental and physical stress, all associated with decreased cardiovascular functions such as abnormal heart rate variability (HRV).<sup>11,24</sup> Wildland firefighters also experience multiple exposures, such as smoke which can contain numerous pollutants, including carbon monoxide, formaldehyde, nitrogen oxides, polycyclic aromatic hydrocarbons, volatile organic compounds, and secondary particulate matter with demonstrated links to cardiovascular diseases such as hypertension, dysrhythmia, atherosclerosis, peripheral thrombosis, and myocardial ischemia.<sup>9,25,26</sup> Apart from these, cardiovascular health of WFFs may also be affected by environmental factors such as ambient temperature in their workplace. The heavy physical work in the increased ambient temperature while suppressing wildfire may result in increased internal body temperature, which may cause heat stroke, cardiac failure, dehydration, and heat-related illness.<sup>27</sup> The subclinical measures reported here were from a medical standards program and not temporally related to the physical and environmental risks of arduous duty wildland firefighting work. Nevertheless, if further study confirms a greater baseline presence of hypertension among the WFF worker population, the work-related physical and environmental factors may translate to greater than anticipated cardiovascular risk.

These findings of our study agree with earlier findings where it was observed that self-reported elevated blood pressure or hypertension was associated with extended work history among WFFs.<sup>28</sup> One hypothesized mechanism whereby WFFs are at greater risk of CVD events is due to job task-related aberrations in HRV that is associated with risk of coronary heart disease, specifically acute myocardial infarction.<sup>24,29</sup> Job-specific factors related to WFF cardiovascular risk include dehydration, high intensity of physical activity, acclimatization, humidity, and ambient temperature. All of these factors increase the risk of developing heat-related disorders like heat exhaustion, heat stress, and heat stroke in WFFs.<sup>27</sup> Each of these factors can translate to risk of acute cardiovascular event, particularly among workers who may have underlying risk factors for cardiovascular disease.

We acknowledge several limitations to this work. First, WFFs perform a variety of tasks according to credentials and experience, and further investigation of how these subclinical measures vary across job titles is warranted. Second, we do not comprehensively define cardiovascular health; rather, we focus on select measures of BMI, total cholesterol, and blood pressure. These three objective measures were based on similar clinical procedures between the two populations. However, we do note some procedural discrepancy in blood pressure measures with one recorded measure for WFFs versus an average of three measures for the NHANES sample. The self-reported variables such as smoking and diabetes were less useful for comparison purposes as the corresponding questions were phrased in different ways for the two populations. We also do not have more detailed information on use of nicotine pouches and chew that can be common in this population. Furthermore, the self-reported information may have several biases, including the chance of WFF underreporting of diabetes or smoking status due to fear of the negative impact on job assignment or qualification. Third, the NHANES may not be an appropriate population to compare with WFFs, and we were unable to account for occupation among the comparison population. Finally, the MSP does not include data on medications the WFFs may have been using for control of hypertension, diabetes, or hypercholesterolemia nor on other potential confounders, such as education level or household income. The above data limitations are not surprising given that the agency's medical standards program is charged with determining medical qualifications for WFF arduous duty work and not primarily designed as a surveillance tool. Nevertheless, these data limitations indicate opportunities for improving data collection strategies should the agencies employing WFFs transition to a health surveillance paradigm.

This study is among the earliest attempt to explore the cardiovascular health in a large sample of federal WFFs. This study demonstrates

how routinely collected clinical measures as part of a medical standards program can be used to examine the health status of WFFs. These findings indicate the need to explore these and other cardiovascular health measures more deeply to understand the cardiovascular risk profile of WFFs. Future research should also assess how workplace factors and environmental exposures may be associated with cardiovascular risks.

## REFERENCES

1. National Interagency Fire Center (2023). Total wild and fires and acres. Available at: <https://www.nifc.gov/fire-information/statistics/wildfires>. Accessed July 19, 2023.
2. Hoover K, Hanson LA. *Wildfire statistics*. Congressional Research Service; 2021. Available at: <https://fas.org/spp/crs/misc/IF10244.pdf>. Accessed July 21, 2023.
3. Chikamoto Y, Timmermann A, Widlansky MJ, Balmaseda MA, Stott L. Multi-year predictability of climate, drought, and wildfire in southwestern North America. *Sci Rep* 2017;7:6568.
4. Aghakouchak A, Chiang F, Huning LS, et al. Climate extremes and compound hazards in a warming world. *Annu Rev Earth Planet Sci* 2020;48:519–548.
5. Ragno E, Aghakouchak A, Love CA, Cheng L, Vahedifard F, Lima CHR. Quantifying changes in future intensity-duration-frequency curves using multimodel ensemble simulations. *Water Resour Res* 2018;54:1751–1764.
6. Burke M, Driscoll A, Hefl-Neal S, Xue J, Burney J, Wara M. The changing risk and burden of wildfire in the United States. *Proc Natl Acad Sci U S A* 2021;118:e2011048118.
7. Jaffe DA, O'Neill SM, Larkin NK, et al. Wildfire and prescribed burning impacts on air quality in the United States. *J Air Waste Manag Assoc* 2020;70:583–615.
8. Navarro KM, Butler CR, Fent K, et al. The Wildland Firefighter Exposure and Health Effect (WFFEHE) study: rationale, design, and methods of a repeated-measures study. *Ann Work Exposures Health* 2022;66:714–727.
9. Ruby BC, Coker RH, Sol JA, Quindry JC, Montain SJ. Physiology of the wildland firefighter: managing extreme energy demands in hostile, smoky, mountainous environments. *Compr Physiol* 2023;13:4587–4615.
10. Gurney SC, Christison KS, Williamson-reisdorph CM, et al. Alterations in metabolic and cardiovascular risk factors during critical training in wildland firefighters. *J Occup Environ Med* 2021, 2021;63:594–599.
11. Vincent GE, Aisbett B, Wolkow A, Jay SM, Ridgers ND, Ferguson SA. Sleep in wildland firefighters: what do we know and why does it matter? *Int J Wildland Fire* 2018;27:73–84.
12. US Government Accountability Office (2022). Wildland fire: barriers to recruitment and retention of Federal Wildland Firefighters. GAO-23-105517 November 2022. Available at: <https://www.gao.gov/products/gao-23-105517>. Accessed July 21, 2023.
13. Rahn M, Broyles G, Mchale T, Brown C, Edwards T, Hale K. Wildland firefighter exposure study: evaluating core temperature, heart rate & hydration. *Wildfire research report 5 – summer 2016* 2016. Available at: <https://wildfirescience.org/wp-content/uploads/2018/09/2017WildlandResearchReport.pdf>. Accessed July 21, 2023.
14. National Wildfire Coordinating Group. (2017). NWCG report on wildland firefighter fatalities in the United States: 2007-2016. Report, December, 18. Available at: <https://www.nwcg.gov/sites/default/files/publications/pms841.pdf>. Accessed July 21, 2023.
15. Navarro KM, Kleinman MT, Mackay CE, et al. Wildland firefighter smoke exposure and risk of lung cancer and cardiovascular disease mortality. *Environ Res* 2019;173:462–468.
16. Lloyd-Jones DM, Hong Y, Labarthe D, et al, American Heart Association Strategic Planning Task Force and Statistics Committee. Defining and setting national goals for cardiovascular health promotion and disease reduction: the American Heart Association's strategic impact goal through 2020 and beyond. *American Heart Association* 2010;121:586–613.
17. Chrostowska M, Ph D, Szyndler A, et al. Impact of obesity on cardiovascular health. *Best Pract Res Clin Endocrinol Metab* 2013;27:147–156.
18. Daniels TF, Killinger KM, Michal JJ, Wright RW Jr., Jiang Z. Lipoproteins, cholesterol homeostasis and cardiac health. *Int J Biol Sci* 2009;5:474–488.
19. Fuchs FD, Whelton PK. High blood pressure and cardiovascular disease. *Hypertension* 2020;75:285–292.
20. U.S. Department of the Interior (2023). Wildland firefighter medical standards. Available at: <https://www.doi.gov/wildlandfire/medical-standards>. Accessed June 19, 2023.
21. Center for Disease Control and Prevention (2017). Laboratory procedure manual: cholesterol. Available at: [https://wwwn.cdc.gov/nchs/data/nhanes/2015-2016/labmethods/BIOPRO\\_1\\_MET\\_CHOLESTEROL\\_DX800and660i.pdf](https://wwwn.cdc.gov/nchs/data/nhanes/2015-2016/labmethods/BIOPRO_1_MET_CHOLESTEROL_DX800and660i.pdf). Accessed July 21, 2023.
22. Center for Disease Control and Prevention (2016). National Health and Nutrition Examination Survey (NHANES): anthropometry procedures manual. Available at: [https://wwwn.cdc.gov/nchs/data/nhanes/2015-2016/manuals/2016\\_Anthropometry\\_Procedures\\_Manual.pdf](https://wwwn.cdc.gov/nchs/data/nhanes/2015-2016/manuals/2016_Anthropometry_Procedures_Manual.pdf). Accessed July 21, 2023.

23. Center for Disease Control and Prevention (2019). National Health and Nutrition Examination Survey (NHANES): blood pressure procedures manual. June 2019. Available at: <https://wwwn.cdc.gov/nchs/data/nhanes/2019-2020/manuals/2019-Blood-Pressure-Procedures-Manual-508.pdf>. Accessed July 21, 2023.
24. Jeklin AT, Perrotta AS, Davies HW, Bredin SSD, Paul DA, Warburton DER. The association between heart rate variability, reaction time, and indicators of workplace fatigue in wildland firefighters. *Int Arch Occup Environ Health* 2021;94:823–831.
25. Brook RD, Rajagopalan S, Pope CA 3rd, et al. Particulate matter air pollution and cardiovascular disease: an update to the scientific statement from the American Heart Association. *Circulation* 2010;121:2331–2378.
26. Pope CA 3rd, Dockery DW. Health effects of fine particulate air pollution: lines that connect. *J Air Waste Manage Assoc* 2006;56:709–742.
27. West MR, Costello S, Sol JA, Domitrovich JW. Risk for heat-related illness among wildland firefighters: job tasks and core body temperature change. *Occup Environ Med* 2020;77:433–438.
28. Semmens EO, Domitrovich J, Conway K, Noonan CW. A cross-sectional survey of occupational history as a wildland firefighter and health. *Am J Ind Med* 2016;335:330–335.
29. Liao D, Cai J, Rosamond WD, et al. Cardiac autonomic function and incident coronary heart disease: a population-based case-cohort study: the ARIC study. *Am J Epidemiol* 1997;145:696–706.