

## Perception of whole day workload as a mediator between activity engagement and stress in workers with type 1 diabetes

Raymond Hernandez, Haomiao Jin, Elizabeth A. Pyatak, Shawn C. Roll, Jeffrey S. Gonzalez & Stefan Schneider

**To cite this article:** Raymond Hernandez, Haomiao Jin, Elizabeth A. Pyatak, Shawn C. Roll, Jeffrey S. Gonzalez & Stefan Schneider (2024) Perception of whole day workload as a mediator between activity engagement and stress in workers with type 1 diabetes, *Theoretical Issues in Ergonomics Science*, 25:1, 67-85, DOI: [10.1080/1463922X.2022.2149878](https://doi.org/10.1080/1463922X.2022.2149878)

**To link to this article:** <https://doi.org/10.1080/1463922X.2022.2149878>



Published online: 28 Nov 2022.



Submit your article to this journal [↗](#)



Article views: 177



View related articles [↗](#)



View Crossmark data [↗](#)



# Perception of whole day workload as a mediator between activity engagement and stress in workers with type 1 diabetes

Raymond Hernandez<sup>a,b</sup> , Haomiao Jin<sup>a,c</sup>, Elizabeth A. Pyatak<sup>b</sup> ,  
Shawn C. Roll<sup>b</sup> , Jeffrey S. Gonzalez<sup>d,e</sup> and Stefan Schneider<sup>a</sup>

<sup>a</sup>Dornsife Center for Economic & Social Research, University of Southern California, Los Angeles, CA, USA; <sup>b</sup>Chan Division of Occupational Science and Occupational Therapy, University of Southern California, Los Angeles, CA, USA; <sup>c</sup>School of Health Sciences, University of Surrey, Guildford, UK; <sup>d</sup>Department of Medicine (Endocrinology), Albert Einstein College of Medicine, Bronx, NY, USA; <sup>e</sup>Ferkauf Graduate School of Psychology, Yeshiva University, Bronx, NY, USA

## ABSTRACT

Associations between various forms of activity engagement (e.g. work, leisure) and the experience of stress in workers have been widely documented. The mechanisms underlying these effects, however, are not fully understood. Our goal was to investigate if perceived whole day workload accounted for the relationships between daily frequencies of activities (i.e. work hours and leisure/rest) and daily stress. We analysed data from 56 workers with type 1 diabetes (T1D) who completed approximately two weeks of intensive longitudinal assessments. Daily whole day workload was measured with an adapted version of the National Aeronautics and Space Administration Task Load Index (NASA-TLX). A variety of occupations were reported, including lawyer, housekeeper and teacher. In multilevel path analyses, day-to-day changes in whole day workload mediated 67% ( $p < .001$ ), 61% ( $p < .001$ ), 38% ( $p < .001$ ), and 55% ( $p < .001$ ) of the within-person relationships between stress and work hours, rest frequency, active leisure frequency, and day of week, respectively. Our results provided evidence that whole day workload perception may contribute to the processes linking daily activities with daily stress in workers with T1D. Perceived whole day workload may deserve greater attention as a possible stress intervention target, ones that perhaps ergonomists would be especially suited to address.

## ARTICLE HISTORY

Received 25 August 2022  
Accepted 11 November 2022

## KEYWORDS

Workload; stress; rest; work hours; type 1 diabetes

## Relevance to human factors/ergonomics theory

Day-to-day changes in whole day workload accounted for a substantial portion of the relationship between daily activity engagement and daily stress in workers with type 1 diabetes, thus suggesting that the workload dimensions of the whole day NASA-TLX (e.g. mental demands, physical demands, etc.) may warrant greater consideration as potential

**CONTACT** Raymond Hernandez [hlay57024@gmail.com](mailto:hlay57024@gmail.com) USC Center for Economic & Social Research, 635 Downey Way, VPD 405, Los Angeles, CA 90089-3332, USA.

© 2022 Informa UK Limited, trading as Taylor & Francis Group

stress intervention targets. Ergonomists may be particularly suited to address stress by targeting the dimensions of whole day workload. For instance, they could help ensure that the systems people interact with are optimised to minimise the associated workload.

## Introduction

Stress continues to be an issue that needs to be addressed in the U.S. According to the 2015 American Psychological Association report “Stress in America”, thirty-four percent of adults reported that their stress increased over their past year, with approximately 24% of adults experiencing extreme stress (Anderson et al. 2016). The COVID pandemic has worsened matters, with a 2021 Gallup poll finding that 57% of U.S. and Canadian workers felt stressed on a daily basis (Gallup 2021). There has been consistent evidence that exposure to excessive stress can contribute to the development of conditions such as cardiovascular disease and depression (Ganster and Rosen 2013), as well as decrements in job performance (Jamal 2011; Rosen et al. 2010).

Stress has different conceptualizations, but here we use its Transactional Model of Stress definition as the judgement that particular environmental demands may tax an individual's resources and threatens his/her well-being (Holroyd and Lazarus 1982). According to the Transactional Model, the appraisal of a potential threat is a transaction between person and environment (Lazarus 2001). Primary appraisal involves the judgement that something of value (e.g. well-being) is being threatened by the environment (Lazarus 2001). If a threat is perceived, then a secondary appraisal is made of whether coping resources are available, after which varying degrees of the stress experience may arise (Lazarus 2001).

## *Stress and activity engagement*

Longer engagement in work activities may be associated with both higher perception of threat to well-being (primary appraisal), and use of more resources (e.g. physical and mental energy) to meet the demands of often taxing job tasks (secondary appraisal), resulting in a higher magnitude of stress experienced as per the Transactional Model of Stress. Consistent with this line of reasoning, the activities workers engage in are associated with the amount of stress experienced. For instance, longer work hours have often been associated with higher stress. In a review of 21 studies, Sparks et al. (2001) found small but significant positive correlations between work hours and various psychological symptoms, including stress. Compared to those working shorter hours, workers with longer working hours have been found to perceive more stress (Lee et al. 2017; Maruyama and Morimoto 1996).

Apart from engagement in work activities, greater engagement in recovery activities may be associated with a decrease in perceived threats (primary appraisal) and an increase in the perception of personal resources (secondary appraisal), lowering experienced stress. Recovery has been defined as return of psychobiological systems to baseline after changes they had undergone due to the experience of stress (Meijman and Mulder 1998). It is important to distinguish between different types of recovery, as their relationships with stress may differ. Both rest and active forms of casual leisure (e.g. play and socializing) (Stebbins 1997) are frequently examined forms of recovery from work (Sonnentag and Fritz

2007; Trenberth and Dewe 2002). Rest is defined here as tasks accompanied by increases in parasympathetic activity (Tindle and Tadi 2021). Active casual leisure is casual leisure activities characterised by a greater degree of action and effort (mental or physical) exerted compared to passive casual leisure pursuits, such as relaxation and passive entertainment (Stebbins 1997). Examples of active casual leisure are play, active entertainment, and socializing (Stebbins 1997). Casual leisure can be distinguished from serious leisure, which is leisure requiring special training and thus greater effort for its enjoyment (Stebbins 1997). Relative to casual leisure, serious leisure may have a greater chance of at times being associated with greater stress, especially forms with high pressure for performance due to associated social evaluation (e.g. sports) (Kimball and Freysinger 2003). Decreased stress has been associated with more time resting (Esch, Fricchione, and Stefano 2003; Sianoja et al. 2018) and in active forms of casual leisure (Winwood, Bakker, and Winefield 2007; Zawadzki, Smyth, and Costigan 2015).

### ***Workload as a mediator between activity engagement and stress at the daily level***

Perception of workload should theoretically inform the primary appraisal of environmental threats (i.e. demands) that precede the experience of stress, making workload a potential mediator between activity engagement and stress. Results of a few prior studies support the workload as a mediator. In one cross-sectional study, a significant association between overtime work and stress was much reduced and became nonsignificant after statistical adjustment for self-reported workload and sleeping time (Sato, Miyake, and Thériault 2009). The authors concluded that engagement in overtime work may have in part impacted stress through workers' perceptions of the overall workload of their positions. Another study with repeated daily measures found that engagement in recovery activities was moderately correlated with both lower stress and lower perceived whole day workload (Hernandez et al. 2021), suggesting the possibility that workload may also in part account for the relationship between recovery activity frequency and stress at the daily level.

Examination of whether whole day workload acts as a mediator between activity engagement and stress, with ambulatory assessments completed over multiple days, may lead to insights potentially useful for ergonomics. Ambulatory assessments completed daily allow for examining phenomena in people's natural environments, and account for daily fluctuations in the phenomena being measured (Trull and Ebner-Priemer, 2013). Thus, ambulatory assessment data for activity engagement, workload, and stress, allows us to investigate the dynamic within-person relationships among these variables across time, and in real world contexts. There has been evidence supporting the use of the NASA-TLX to assess workload experienced *over whole days*, inclusive of both work and non-work periods (Hernandez et al. 2022). In a two-week longitudinal study, daily workload ratings as captured by the NASA-TLX were found to demonstrate convergent validity as evidenced by theoretically expected associations with a variety of daily activity engagement and well-being measures (Hernandez et al. 2022). We build on this prior work to examine whether perceptions of whole day workload may account for the relationships between day-to-day changes in work and recovery activities and corresponding changes in workers' daily stress levels.

## The present study

We hypothesised that engagement in work, rest, and active casual leisure may impact daily stress in part *through* their effects on whole day workload (Figure 1). From here on, we will refer to active casual leisure as “leisure” or “active leisure”. Four different operationalizations of activity engagement were used. The first three are daily frequencies of work hours, rest, and leisure, which could differ in the degree to which they impact stress through whole day workload. Weekends versus weekdays may also influence daily stress through whole day workload given how highly they are related to work hours, rest, and leisure activity frequency (Sonnentag, Venz, and Casper 2017). Thus, we also examined weekend versus weekday as an activity engagement variable.

Our hypotheses align with the literature described prior. We anticipated that greater work hours would increase the perceived whole day workload, which in turn could increase experienced stress. That is, we expected the perception of whole day workload to act as a partial mediator between work hours and same day stress.


*Hypothesis 1:* Perception of whole day workload acts as a partial mediator between work hours and same day stress.

Furthermore, we also anticipated that rest and active leisure (both during and after work hours) may *reduce* stress, also in some part through their effects on whole day workload. We anticipated that greater frequency of rest in a day would decrease stress experienced indirectly via less experienced whole day workload. Additionally, we hypothesised that a greater frequency of active leisure would also decrease stress, but that only a small portion of this relationship would be attributable to a decrease in same day workload. That is, we expected active leisure to decrease stress only in small part due to associated workload, because of their higher demands relative to restful engagements, and more so through other pathways such as the intrinsic pleasure of play (Stebbins 1997).

*Hypothesis 2a:* Perception of whole day workload acts as a partial mediator between recovery activities (rest and leisure) and same day stress.

*Hypothesis 2b:* Whole day workload mediates a greater portion of the relationship between rest and stress, as compared to the proportion mediated between leisure and stress.

Finally, we hypothesised that weekends were less stressful than weekdays, in part through their relationship with whole day workload. The weekday/weekend distinction often brings with it corresponding frequencies in work/rest/leisure engagement, with work hours typically lower and rest/leisure participation higher on weekends for workers with the traditional five-day work week.



**Figure 1.** Whole day workload as mediator between engagement in work hours/rest/leisure and stress.

*Hypothesis 3:* Perception of whole day workload partially mediates the decreased stress typically associated with weekends compared to weekdays.

We examine our hypotheses in a population for whom whole day workload may be particularly relevant, workers with type 1 diabetes. In addition to the daily demands of their jobs, these workers also contend with persistent demands from the management of their diabetes (Hansen et al. 2018). Their self-management responsibilities include managing insulin intake, monitoring their blood glucose, problem-solving issues that may arise with diabetes management equipment, and planning ways of continuing diabetes management in various contexts (Beck et al. 2017). Thus, workload experienced over whole days may be especially appropriate to consider for workers with diabetes as a mediator between activity engagement and daily stress.

## Methods

### *Study overview*

We analysed data collected as part of a multisite study investigating the relationship between function, emotion, and blood glucose in adults with type 1 diabetes (T1D), described elsewhere in greater detail (Pyatak et al. 2021). Participants aged 18 and older with T1D were recruited from three clinical sites through mailings, phone calls, email invitations and health provider referrals. Only participants that identified as workers were included in analyses. Participants completed the following: electronic informed consent remote through the REDCap e-consent framework (Harris et al. 2009), baseline surveys, 14 days of ecological momentary assessment (EMA) and end-of-day survey data collection with 5–6 surveys per day, and follow-up surveys. EMA is repeated measurement of participant's momentary experiences in their real-world environment (Shiffman, Stone, and Hufford 2008). The data collection procedures were approved by the [authors'] Institutional Review Board.

### *Measures*

Participants were asked to complete 5–6 surveys per day on smartphones, for approximately 2 weeks. The Mobile EMA application (mEMA: ilumivu.com) was the platform through which surveys were completed, and was also the source of alarm notifications every three hours for when to complete surveys. Table 1 lists all the EMA questions utilised in this paper's analyses. At each survey, participants reported the type of activity they were presently engaging in and momentary feelings of stress. At the last assessment of each day, work hours and a subjective assessment of whole day overall workload were obtained. For a full list of the items administered, please refer to (Pyatak et al. 2021).

The frequency of rest and active leisure engagements in a day were calculated from the activity type item using an approach with preliminary evidence of validity (Hernandez et al. 2021). Reports of engagement in “relaxing/chilling” and “sleeping/napping” contributed to the daily frequency count for rest, and “socializing” and “fun/play/leisure” increased the count for active leisure. Daily activity frequencies were calculated by dividing activity counts for a day, by the number of surveys taken in the same day, only on days with four or more surveys completed. As an example, if a participant took six surveys in a day, reporting

**Table 1.** EMA and end of day measures to investigate relationship between activity engagement, workload, and stress.

Construct	Item(s)	Response Option(s)	Time
Activity type	What were you doing right before starting this survey?	<ul style="list-style-type: none"><li>• Work/school activities (e.g. paid labour, volunteer work, and studying)<sup>a</sup></li><li>• Traveling (e.g. driving, riding in a car, walking)</li><li>• Relaxing/chilling (e.g. passive leisure like watching Netflix, listening to music)</li><li>• Sleeping/napping</li><li>• Socializing (e.g. talking with friends/family)</li><li>• Caring for myself (e.g. eating, dressing, bathing, toileting, personal grooming)</li><li>• Caring for others (e.g. caring for your children and pets, if you're caring for others as part of work this counts as "work")</li><li>• Doing housework/errands (e.g. paying bills, washing dishes and clothes, exercising for health)</li><li>• Fun/play/leisure activities (e.g. active leisure like exercising for fun, video games, reading for fun)</li><li>• Other (If chosen, please specify)</li></ul>	All survey times
Work Hours	About how many hours did you work?	Hours, whole number, 0 to 24	End of day
Whole Day Workload	Six NASA-TLX items ask about mental demand, physical demand, time pressure, effort, performance satisfaction, and frustration for activities over the whole day. The sum of the ratings indicates total workload.	0 to 100 sliding scale for each item	End of day
Stress	How stressed are you right now?	0 (Not at all stressed) to 100 (Extremely stressed)	All survey times

<sup>a</sup>Activity examples were not in the actual item, but were explained during training and listed in a manual provided to participants.

engaging in “fun/play/leisure” during one and “socializing” in one, then the rest activity frequency for that day would be  $2/6 = .33$ .

We could have assessed the daily frequency of work using the approach applied for rest and leisure engagements, but instead opted to use the report of work hours. Self-report of work hours is a frequently used measure with evidence supporting its accuracy (Imai et al. 2016; Todd et al. 2010), while calculating work hours from an EMA activity item is a more novel approach (Hernandez et al. 2021). Thus, we opted to use self-report of work hours, as it was the more established measure.

Whole day workload was assessed with an adapted version of the National Aeronautics and Space Administration Task Load Index (NASA-TLX) with some evidence supporting its validity (Hernandez et al. 2022). Like the original TLX, six items addressing different contributors to task load were used (Hart and Staveland 1988), but the period being asked about was changed to a whole day rather than a particular task (e.g. How much mental activity was required for your whole day?). An overall workload score was calculated based



on the summation of the six items addressing an individual's perception of mental demand, physical demand, temporal demand, performance, effort, and frustration level (Hart 2006). The raw, as opposed to weighted, sum is used because of prior research supporting this approach (Hart, 2006; Hernandez et al. 2022). Response options for all items were a 0 to 100 slider scale.


Our item for stress was derived from a prior EMA study (Dunton et al. 2019), and asked in all EMA surveys ("How stressed are you right now?"). Single-item measures are often used in EMA contexts (Dunton et al. 2019; Tung et al. 2022) to avoid burdening participants with repeatedly answering long sets of questions, and some evidence supports the validity of a single stress item for capturing momentary stress (King et al. 2019). We assumed that our participants' understanding of the term stress would align with its conceptualisation in the Transtheoretical Model of Stress, in part based on prior qualitative research finding that lay interpretations of stress often involved a focus on stressors, and/or strain reactions to the stressors (Mark and Smith 2018; Mazzola, Schonfeld, and Spector 2011). The ideas of stressors (i.e. threat) and strain reactions (i.e. stress experience after the evaluation of a threat) align well with the Transtheoretical Model of Stress. The average of the stress ratings in a day was used to indicate stress experienced during that day. A previous study found a high correlation between the average of momentary stress ratings, and an end of day overall stress question (Lourenco et al. 2022).

### **Statistical analyses**

To help characterise the participants in our sample and possibly aid in the interpretation of results of the primary analyses, we calculated a variety of descriptive statistics: demographic characteristics, frequency various activities were reported, mean values of study measures by day of week, and between and within-person correlations between study measures. Between and within-person correlations were calculated using the "statsBy" function in the "psych" package, available for the statistical software R (Revelle and Revelle 2015). The focus of our analyses is on the within-person level, but between-person correlations were calculated to comprehensively represent the relationships between variables.

Multilevel structural equation modeling (MSEM), specifically multilevel path analysis, was used to investigate if whole day workload mediated the relationship between activity engagement variables (work hours, rest frequency, active leisure frequency, weekday vs. weekend) and stress (Figures 2 and 3). Note that, unlike the other activity variables, the "weekend versus weekday" distinction is binary, and the model coefficients for the weekend model will be comparisons to weekdays. Analyses were run with the software *Mplus* (Muthén and Muthén 1998) and R package *Mplus* automation (Hallquist and Wiley 2018). Multi-level path analysis is a special case of MSEM where only measured variables are used and no latent variables are modelled (Stoelting 2002). Direct effects ( $c'$ ), indirect effects ( $a*b$ ), and total effects ( $c' + a*b$ ) were examined. MSEM is able to accommodate nested data structures (Huang 2017), as was necessary for our data as it had multiple surveys observations nested in individuals. Had we used single-level SEM, resultant parameter estimates would have been biased as the non-independence of survey observations from the same person would not have been accounted for (Huang 2017). To account for potential non-independence in observations stemming from proximity in time (i.e., autocorrelated






**Figure 2.** Hypothesised mediation model relating work hours with stress via whole day workload. Mediation occurs at the within-person level (i.e. for an individual, daily stress is higher on days with more working hours, and part of this effect stems from the perception of whole day workload for the same day). All other aspects of the model account for factors that can impact the accuracy of the estimated mediation effects in the current multilevel (days nested in individuals) study design. Random intercepts shown on the between-person level account for between person variation (i.e. individual differences in work hours, whole day workload, and stress typically experienced). The arrows between these intercepts indicate that they are allowed to correlate with one another. Study day is controlled for at the within-person level. Finally, the residuals of work hours, whole day workload, and daily stress ( $e_t$ ) are regressed on residuals of the day prior ( $e_{t-1}$ ) to account for dependencies of observations that are more proximal in time.

residuals), a residual dynamic structural equation modeling (RDSEM) structure was added in all models, where the residuals of the main variables of interest were regressed on their respective residuals from the day prior, at the within-person level (Asparouhov, Hamaker, and Muthén 2018). Mediation in Figures 2 and 3 occurs within-person, or at level 1. The primary variables at level 1 are also regressed on study day to account for potential effects of time since data collection commencement. At the between-person level (level 2), only correlations between person specific averages (intercepts) of variables measured at level 1 were modelled, which accounts for dependencies of observations nested within individuals that may affect the standard errors of the within-person mediation (Preacher, Zyphur, and Zhang 2010).

MSEMs were tested with Bayesian parameter estimation using the *Mplus* default non-informative priors. Like bootstrapping, this procedure appropriately takes the non-normal distribution of indirect (mediated) effects into account (Muthén and Asparouhov 2012). Statistical significance of mediated effects was determined using the product of coefficients method and inspecting 95% credible intervals.



**Figure 3.** Theorised within-person mediation model between rest/active leisure frequency/weekend and stress, through whole day workload. Solid lines indicate positive pathways, and dashed ones negative.

One notable benefit in our use of MSEM was that it allowed us to take full advantage of our longitudinal dataset to analyse the within person relationships between activity engagement, workload, and stress. MSEM allows for disaggregation of within-person and between-person effects (Curran and Bauer 2011). Without such disaggregation, one could not disentangle, for instance, whether experiencing greater leisure than average decreases stress (within-person effect), or if people that on average have greater participation in leisure activities experience less stress (between-person effect).

### Statistical power

To determine the magnitude of between- and within-person correlations that could be detected, power analyses were conducted using Monte Carlo simulations in *Mplus* version 8.8 (Muthén and Muthén 1998). Based on prior research, we assumed that study measures had intraclass correlation coefficients (ICCs) of 0.5 (Podsakoff et al. 2019). Given a sample size of 56 people and assuming 12 observations per person, the results of the simulation indicated 80% power to detect within-person correlations of at least 0.12 and between-person correlations of at least 0.39.

The MSEMs appeared sufficiently powered to detect the extent to which whole day workload mediated the relationship between activity engagement and stress. There is no consensus yet on sample size requirements for different types of MSEM, and research in this area is ongoing (Sadikaj et al. 2021), but we may be able to follow recommendations for SEM. In

MSEM with paths at both the between- and within-person levels, at least 50 clusters has often been suggested as the minimum required (Muthén 1989). The MSEMs in this paper, however, are like single-level mediation models, because only intercepts and no paths are modelled at the between-person level. Thus, we may be able to follow sample size recommendations for single-level mediation modeling (SEM). In SEM, simulations found that a sample size of 562 observations is required to be able to detect a small indirect effect (with  $a$  and  $b$  paths jointly as small as .14) with 80% power (Fritz and MacKinnon 2007). At the within-person level, with approximately 60 observations from each participant, we calculated that  $n = 50$  would yield  $50 \times 60 = 3000$  observations. Because this figure is higher than 562, our MSEM analyses should have sufficient power to detect small within-person mediation effects.

## Results

Analyses were conducted on 56 workers who primarily worked full-time and had an average age of 39.4 (SD = 12.8) years. Participants were diverse with regards to gender, ethnicity, education, and annual income (Table 2). Reporting of vocations was not required, but participants that responded to this question disclosed a variety of occupations, including lawyer, engineer, housekeeper, teacher, and security guard. In terms of EMA compliance, the median EMA completion percentage was 92%, and four or more EMA surveys were completed on 83% of all data collection days across the participants. On average, we received 12.5 days of survey data from participants with at least four EMA surveys completed. In total, 4,051 EMA surveys were completed by participants across 843 days. Among these, the distribution of responses within each activity type is provided in Table 3.

**Table 2.** Demographic characteristics for study on activity engagement, workload, and stress.

Characteristic	<i>n</i>	Mean (SD) or Percentage (%)
Age (years)	56	39.4 (12.8)
Gender		
Male	25	45%
Female	31	55%
Ethnicity		
White	23	41%
Latino/x	15	27%
African American	8	14%
Multi-ethnic	4	7%
Other	6	11%
Employment status		
Full-time	44	79%
Part-time	12	21%
Education		
High school grad or less	7	13%
Some college, no degree	10	18%
Associate's degree	4	7%
Bachelor's degree	21	38%
Graduate degree	14	25%
Annual household income		
<\$50,000	14	25%
\$50,000–\$99,999	12	21%
≥\$100,000	17	30%
Don't wish to provide	10	18%
Don't know	3	5%

Full between and within-person correlation matrices are shown in [Tables 4a and 4b](#), respectively. At the within-person level, where mediation was tested in our models, intercorrelations between all study measures were small or moderate (Cohen 2013), except for the relationship between rest frequency and active leisure frequency ( $r = -.05$ ,  $p = .173$ ).

The mean values of work hours, daily rest/active leisure, and stress/workload by day of week appeared to be in line with what would be expected for the typical five-day work week ([Figure 4](#)). Mean work hours were higher on weekdays as compared to weekends, and the opposite was seen for rest/active leisure. Across all days, rest had a higher mean frequency as compared to active leisure. Both mean stress and whole day workload were also lower on weekends as compared to weekdays.

**Primary analyses: MSEM to test mediation**

[Table 5](#) shows the results of MSEM testing the models in [Figures 2 and 3](#). More work hours in a day was associated with higher stress, as indicated by the positive total effect on stress (total effect = .91, 95% C.I. 0.67 to 1.16, standardised estimate = .26). Whole day workload was found to be a partial mediator between work hours and stress, with 67% of work hours’ effect on stress attributable to whole day workload. Conversely, a higher frequency of both rest and active leisure was associated with less stress, as evidenced by their total effects on stress of  $-14.78$  (95% C.I.  $-19.45$  to  $-10.03$ , standardised estimate =  $-0.22$ ) and  $-16.67$  (95% C.I.  $-23.31$  to  $-9.99$ , standardised estimate =  $-0.17$ ), respectively. Whole day workload again acted as a significant partial mediator for these relationships, with workload

**Table 3.** Frequency distribution of activity types across all datapoints ( $n=4,051$ ) for study on activity engagement, workload, and stress.


Activity type	Frequency (%)
Work/school activities	1095 (27.0%)
Relaxing/chilling	852 (21.0%)
Sleeping/napping	535 (13.2%)
Doing housework/errands	448 (11.1%)
Caring for myself	432 (10.7%)
Fun/play/leisure activities	249 (6.1%)
Traveling	202 (4.8%)
Socializing	119 (2.9%)
Caring for others	65 (1.6%)
Other	54 (1.3%)

**Table 4a.** Between-person intercorrelation matrix.

	Work Hours	Rest Freq.	Active Leisure Freq.	Whole Day Workload	Stress
Work Hours	1, $p < .001$				
Rest Freq.	$-0.08$ , $p = .577$	1, $p < .001$			
Active Leisure Freq.	$-0.14$ , $p = .288$	$-0.34$ , $p = .011$	1, $p < .001$		
Whole Day Workload	$.30$ , $p = .026$	$-.28$ , $p = .034$	$-.04$ , $p = .748$	1, $p < .001$	
Stress	$.28$ , $p = .039$	$-.17$ , $p = .215$	$-.12$ , $p = .396$	$.74$ , $p < .001$	1, $p < .001$

**Table 4b.** Within-person intercorrelation matrix.

	Work Hours	Rest Freq.	Active Leisure Freq.	Whole Day Workload	Stress
Work Hours	1, $p < .001$				
Rest Freq.	−0.38, $p < .001$	1, $p < .001$			
Active Leisure Freq.	−0.30, $p < .001$	−0.05, $p = .173$	1, $p < .001$		
Whole Day Workload	.54, $p < .001$	−.40, $p < .001$	−.22, $p < .001$	1, $p < .001$	
Stress	.31, $p < .001$	−.26, $p < .001$	−.21, $p < .001$	.45, $p < .001$	1, $p < .001$



**Figure 4.** Day of the week and a) mean work hours, b) rest and leisure activity frequency, and c) stress and workload.

accounting for 61% of rest frequency’s negative effect on stress, and 38% of active leisure frequency’s negative effect on stress. Relative to weekdays, weekends were associated with lower stress (total effect =  $-6.78$ , 95% C.I.  $-8.75$  to  $-4.75$ , standardised estimate =  $-0.23$ ). Perception of whole day workload mediated 55% of the effect (compared to weekdays) of weekends on stress. Study day was not significantly associated with any of the primary variables. Autocorrelated residuals were statistically significant and positive for all variables (with the exception of leisure,  $p = .06$ ), and their estimates ranged from 0.16 to 0.42.

**Discussion**

The correlations between study measures were consistent with our model assumptions. Whole day workload was theorised to be a precursor to stress, consistent with the large correlation between the two at both the between-person ( $r = .74$ ) and within-person ( $r = .45$ ) levels. Furthermore, at the within-person level, a significant positive association was seen between stress and work hours ( $r = .31$ ), and negative associations were observed between stress and both rest ( $r = -.26$ ) and active leisure ( $r = -.21$ ).

Our study results supported our hypothesised models in [Figures 2 and 3](#). More than two-thirds of the effect of work hours on stress was attributable to whole day workload (67%), suggesting the possibility that the perception of whole day workload is an integral aspect of how stress arises from work hours (hypothesis 1). The high proportion of mediation by whole day workload on the relationship between work hours and stress made theoretical sense, given that much of the stress from work hours is expected to come from associated workload, in contrast to other sources of stress such as workplace

**Table 5.** Whole day workload as a mediator between work hours/rest/leisure frequency and stress.

	EFFECT	EST.	EST. 95% C.I.	STD. EST.
<b>WORK HOURS</b> $a = 1.78$ ( $p < .001$ ) $b = 0.35$ ( $p < .001$ )	Indirect	0.61	(0.46,0.77)*	0.18
	Direct	0.30	(0.04,0.56)*	0.09
	Total	0.91	(0.67,1.16)*	0.26
	Indirect/Total	0.67	(0.48,0.953)*	0.67
<b>REST</b> $a = -25.42$ ( $p < .001$ ) $b = 0.35$ ( $p < .001$ )	Indirect	-9.00	(-11.52,-6.60)*	-0.13
	Direct	-5.73	(-10.51,-1.15)*	-0.09
	Total	-14.78	(-19.45,-10.03)*	-0.22
	Indirect/Total	0.61	(0.43,0.89)*	0.61
<b>ACTIVE LEISURE</b> $a = -16.86$ ( $p < .001$ ) $b = 0.37$ ( $p < .001$ )	Indirect	-6.21	(-9.23,-3.41)*	-0.06
	Direct	-10.43	(-16.56,-4.34)*	-0.11
	Total	-16.67	(-23.31,-9.99)*	-0.17
	Indirect/Total	0.38	(0.21,0.61)*	0.38
<b>WEEKEND</b> $a = -10.70$ ( $p < .001$ ) $b = 0.35$ ( $p < .001$ )	Indirect	-3.74	(-4.81,-2.75)*	-0.13
	Direct	-3.03	(-5.03,-0.98)*	-0.10
	Total	-6.78	(-8.75,-4.75)*	-0.23
	Indirect/Total	0.55	(0.38,0.80)*	0.55

Note: \*CIs don't contain 0, indicating statistical significance; EST. = Estimate; C.I. = Credible Interval; STD. EST. = Standardized Estimate.

harassment (Bowling and Beehr 2006) and role ambiguity (Fisher and Gitelson 1983). More than half of the effect of rest frequency on stress was also attributable to whole day workload (61%) (hypothesis 2a). Given that rest (i.e. relaxation experiences) constitutes recovery through low activation activities (Sonnentag and Fritz 2007) with low associated workload, this high proportion again made sense. Active leisure had a higher proportion of its effect on stress mediated by workload than we had expected (38%) (hypothesis 2b). We hypothesised that active leisure activities would be associated with greater demands relative to rest because of its “active” component, and thus that it would decrease stress through other pathways, such as increased self-efficacy from mastery experiences in learning a new hobby (Bandura, Freeman, and Lightsey 1999). While active leisure does have an active component, perhaps the necessary demands are still much lower compared to those experienced from work, and thus much of its effect on stress still acts through decreased whole day workload. Finally, weekends were found to have a large portion (55%) of their effect on stress attributable to whole day workload (hypothesis 3), which is consistent with their association with less work hours and higher rest/leisure activity engagement frequency compared to weekdays.

## Implications

Our results suggest that perceived whole day workload may be useful to research further as a potential stress intervention target in workers with T1D. Firstly, study results provide preliminary evidence that whole day workload is a strong precursor to stress. The two were highly correlated at the between and within-person levels, and constructs with strong relationships to stress (i.e. work hours, rest, and active leisure) were found to impact stress largely through whole day workload. Secondly, whole day workload as it was conceptualised here, may lend itself more readily to intervention efforts as compared to other constructs. Some variables, such as number of work hours, may be difficult to address in interventions because they often are often not readily adjustable by organisations. The six dimensions of

the adapted TLX, however, may each serve as potentially malleable intervention targets. For instance, mental demands may be reducible in work and non-work contexts by automating cognitively demanding tasks where feasible, or helping to streamline particular steps to minimise cognitive load.

Future research examining if whole day workload also acts as a mediator between activity engagement and stress in the general working population may be beneficial. For workers with T1D, whole day workload may have been particularly impactful on stress levels because of the combined demands from work and health management responsibilities. For the general working population, whole day workload may also be an influential precursor to stress, but not to the extent observed in this study. Replication of the study in a general worker population may help provide a clearer idea of the extent to which whole day workload appears to be a viable stress intervention target for all workers.

Assessment of *both* work specific and whole day workload may be useful in future work stress research. Work specific workload assessment may have particular utility in research or interventions focussed purely on the work context. Assessment of whole day workload may be especially useful for holistic research/interventions that consider both work and non-work contexts, as partially evidenced by our study results. When both are measured, additional burden may be placed on participants, but they would provide different information that may provide a more comprehensive understanding of workload exposure.

## Limitations

Study analyses were conducted on a sample of workers with T1D experiencing the COVID-19 pandemic. Our sample may be limited to individuals with T1D, but we expect our sample to be similar to populations with chronic condition(s) generally. Living with chronic conditions is often accompanied by self-care duties such as following complex treatment regimens, self-monitoring one's condition(s), and problem solving when/how to access healthcare services (Dixon, Hibbard, and Tusler 2009). This burden shared by many with chronic conditions is why programs targeting that population generally have been created, such as the Chronic Disease Self-Management program (Lorig et al. 1999). The commonality from the shared burden may also be why research is often conducted on populations with any chronic condition (Keles et al. 2007; Schokker et al. 2010). Nearly 45% of all Americans are diagnosed with at least one chronic condition (Raghupathi and Raghupathi 2018). Regardless, as described prior, replication within other worker populations and settings is needed to confirm and strengthen our findings.

From the mediation models, we cannot determine if work hours, rest, active leisure, and weekends caused stress through whole day workload, given that daily activity engagement, whole day workload, and daily stress were contemporaneous. A variety of other relationships were possible, such as stress acting as a mediator between activity engagement and stress. Even though activity engagement may act as a causal agent of stress, we also cannot exclude that stress experiences throughout the day modulate people's decisions to engage in or disengage from certain types of activities. Stronger conclusions about the direction of mediation effects may be derived from the analysis of temporally lagged effects. However, activity engagement, the perception of demands, and the experience of stress theoretically co-evolve over time, rather than occurring in discrete time frames. The nature of these effects would not have been adequately captured by examining mediated pathways with lagged effects in



which activity engagement on one day affects experiences on the subsequent day, and it would have been unclear what an appropriate time lag between the study measures would have been. One possible method of investigating if whole day workload causes stress is a randomised control trial, where an intervention targets perception of whole day workload, and stress is assessed as one of the primary outcome variables.

## Conclusion

We found that whole day workload, as measured by an adapted version of the NASA-TLX, mediated the relationship between different daily activity engagement measures (work hours, rest frequency, active leisure frequency, type of day) and same day stress in workers with T1D. A large proportion of each measure's effect on stress was mediated by whole day workload, providing preliminary evidence that the perception of workload over a whole day may be an influential precursor to stress. Active leisure frequency's effect on stress was not as strongly mediated by whole day workload relative to the effect seen for rest, and this weaker mediation was expected given its "active" component. Our results provide preliminary evidence that whole day workload may be worth greater consideration as a potential stress intervention target.

## Data availability

The data that support the findings of this study are available from the corresponding author, R.H., upon reasonable request.

## Disclosure statement

No potential conflict of interest was reported by the authors.

## Notes on Contributors

**Raymond Hernandez** is a postdoctoral scholar- research associate in the Center for Self-Report Science at the University of Southern California (USC). He is an occupational therapist, and earned his PhD in occupational science from the USC Chan Division of Occupational Science and Occupational Therapy in 2021. His research interests include recovery from work, whole day workload, momentary assessment of cognitive performance, and ecological momentary assessment.

**Haomiao Jin** is a health data scientist in the School of Health Sciences at the University of Surrey. He earned his PhD in industrial and systems engineering (focus: health data sciences and information technology assessment) from USC in 2016. He is interested in developing and applying machine learning, computational statistics, and systems engineering methods to model complex biomedical data for understanding and improving mental and behavioral health, chronic disease care, and healthy aging.

**Elizabeth A. Pyatak** is an associate professor in the USC Chan Division of Occupational Science and Occupational Therapy. She is an occupational therapist, certified diabetes educator, and earned her PhD in occupational science from USC in 2010. Her current research interests are focused on three primary aims: evaluating innovative care delivery models for adolescents and young adults with chronic conditions; developing strategies to implement occupational therapy lifestyle interventions

in primary and specialty care settings; and examining the lived experience of type 1 diabetes through innovative and emerging technologies.

**Shawn C. Roll** is an associate professor in the USC Chan Division of Occupational Science and Occupational Therapy. He is an occupational therapist, registered musculoskeletal sonographer, and earned his PhD in Health and Rehabilitation Sciences from The Ohio State University in 2011. Dr. Roll has developed an interdisciplinary research career combining his clinical occupational therapy skills with specialization in musculoskeletal sonography and industrial engineering/ergonomics. His research interests include understanding how individual behaviors, physical exposures, stress, psychosocial factors, and personality traits relate to the development of musculoskeletal disorders or other work-related injuries, and optimizing interactions among environments (built, organizational, social) and performance to support the health and well-being of workers.

**Jeffrey S. Gonzalez** is a professor in the Ferkauf Graduate School of Psychology at Yeshiva University, and professor in the Department of Medicine (Endocrinology) at the Albert Einstein College of Medicine. He is a licensed psychologist and earned his degree in clinical psychology from the University of Miami in 2005. His research focuses on identifying psychological and cognitive factors involved in treatment adherence in chronic illness. He has a particular interest in understanding 1) how beliefs about illness and treatment relate to illness self-management and 2) how depression and distress are associated with worse outcomes in chronic illness.

**Stefan Schneider** is a senior research scientist in the Center for Self-Report Science at the University of Southern California (USC). He is a social/health psychologist and earned his PhD in Social/Health Psychology from Stony Brook University in 2010. He is interested in the measurement of self-reported health and wellbeing. In particular, his research seeks to understand the information contained in self-reports, and to develop methods that help to improve their usefulness for characterizing people's health and wellbeing experiences.

## Funding

This work was supported by the National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, under Grant NIH/NIDDK 1R01DK121298-01; Pilot Project Research Training Program of the Southern California NIOSH Education and Research Center under Grant T42 OH008412.

## ORCID

Raymond Hernandez  <http://orcid.org/0000-0002-6761-9276>

Elizabeth A. Pyatak  <http://orcid.org/0000-0003-3280-8912>

Shawn C. Roll  <http://orcid.org/0000-0002-4202-396X>

Stefan Schneider  <http://orcid.org/0000-0002-4562-0524>

## References

- Anderson, N., K. Nordan, D. Ballard, L. Bufka, J. Diaz-Granados, G. Kelta, C. Wright, L. Bossolo, and S. Bethune. 2016. *Stress in America: The Impact of Discrimination. Stress in America Survey*. American Psychological Association, Washington DC.
- Asparouhov, T., E. L. Hamaker, and B. Muthén. 2018. "Dynamic Structural Equation Models." *Structural Equation Modeling: A Multidisciplinary Journal* 25 (3): 359–388. doi:10.1080/10705511.2017.1406803
- Bandura, A., W. H. Freeman, and R. Lightsey. 1999. *Self-Efficacy: The Exercise of Control*. Springer.
- Beck, J., D. A. Greenwood, L. Blanton, S. T. Bollinger, M. K. Butcher, J. E. Condon, M. Cypress, 2017 Standards Revision Task Force, et al. 2017. "2017 National Standards for Diabetes Self-Management Education and Support." *Diabetes Care* 40 (10): 1409–1419. doi:10.2337/dci17-0025

- Bowling, N. A., and T. A. Beehr. 2006. "Workplace Harassment from the Victim's Perspective: A Theoretical Model and Meta-Analysis." *The Journal of Applied Psychology* 91 (5): 998–1012.
- Cohen, J. 2013. *Statistical Power Analysis for the Behavioral Sciences*. Academic Press.
- Curran, P. J., and D. J. Bauer. 2011. "The Disaggregation of within-Person and between-Person Effects in Longitudinal Models of Change." *Annual Review of Psychology* 62: 583–619. doi:10.1146/annurev.psych.093008.100356
- Dixon, A., J. Hibbard, and M. Tusler. 2009. "How Do People with Different Levels of Activation Self-Manage Their Chronic Conditions?" *The Patient: Patient-Centered Outcomes Research* 2 (4): 257–268. doi:10.2165/11313790-000000000-00000
- Dunton, G. F., W. Ke, E. Dzubur, S. G. O'Connor, N. V. Lopez, and G. Margolin. 2019. "Within-Subject Effects of Stress on Weight-Related Parenting Practices in Mothers: An Ecological Momentary Assessment Study." *Annals of Behavioral Medicine: a Publication of the Society of Behavioral Medicine* 53 (5): 415–425. doi:10.1093/abm/kay053
- Esch, T., G. L. Frichione, and G. B. Stefano. 2003. "The Therapeutic Use of the Relaxation Response in Stress-Related Diseases." *Medical Science Monitor* 9 (2): 34.
- Fisher, C. D., and R. Gitelson. 1983. "A Meta-Analysis of the Correlates of Role Conflict and Ambiguity." *Journal of Applied Psychology* 68 (2): 320.
- Fritz, M. S., and D. P. MacKinnon. 2007. "Required Sample Size to Detect the Mediated Effect." *Psychological Science* 18 (3): 233–239. doi:10.1111/j.1467-9280.2007.01882.x
- Gallup. 2021. *State of the Global Workplace Report*. Gallup.Com. <https://www.gallup.com/workplace/349484/state-of-the-global-workplace.aspx>
- Ganster, D. C., and C. C. Rosen. 2013. "Work Stress and Employee Health: A Multidisciplinary Review." *Journal of Management* 39 (5): 1085–1122. doi:10.1177/0149206313475815
- Hallquist, M. N., and J. F. Wiley. 2018. "MplusAutomation: An R Package for Facilitating Large-Scale Latent Variable Analyses in Mplus." *Structural Equation Modeling: A Multidisciplinary Journal* 25 (4): 621–638. doi:10.1080/10705511.2017.1402334
- Hansen, U. M., B. Cleal, I. Willaing, and T. Tjørnhøj-Thomsen. 2018. "Managing Type 1 Diabetes in the Context of Work Life: A Matter of Containment." *Social Science & Medicine* 219: 70–77. doi:10.1016/j.socscimed.2018.10.016
- Harris, P. A., R. Taylor, R. Thielke, J. Payne, N. Gonzalez, and J. G. Conde. 2009. "Research Electronic Data Capture (REDCap)—a Metadata-Driven Methodology and Workflow Process for Providing Translational Research Informatics Support." *J Biomed Inform* 42 (2): 377–381. doi:10.1016/j.jbi.2008.08.010
- Hart, S. G. 2006. "NASA-Task Load Index (NASA-TLX); 20 Years Later." *Proceedings of the Human Factors and Ergonomics Society Annual Meeting* 50: 904–908. doi:10.1177/154193120605000909
- Hart, S. G., and L. E. Staveland. 1988. Development of NASA-TLX (Task Load Index): Results of Empirical and Theoretical Research. In *Advances in Psychology* (Vol. 52, pp. 139–183). Elsevier. doi:10.1016/S0166-4115(08)62386-9
- Hernandez, R., Pyatak, E. A. Vigen, C. L. P. Jin, H. Schneider, S. Spruijt-Metz, D. and Roll, S. C. 2021. "Understanding Worker Well-Being Relative to High-Workload and Recovery Activities across a Whole Day: Pilot Testing an Ecological Momentary Assessment Technique." *International Journal of Environmental Research and Public Health* 18 (19): 10354. Article 19. doi:10.3390/ijerph181910354
- Hernandez, R., S. C. Roll, H. Jin, S. Schneider, and E. A. Pyatak. 2022. "Validation of the National Aeronautics and Space Administration Task Load Index (NASA-TLX) Adapted for the Whole Day Repeated Measures Context." *Ergonomics* 65 (7): 960–975. doi:10.1080/00140139.2021.2006317
- Holroyd, K. A., and R. S. Lazarus. 1982. "Stress, Coping and Somatic Adaptation." In L. Goldberger and S. Breznitz (Eds.), *Handbook of Stress: Theoretical and Clinical Aspects*: (pp. 21–35). New York: Free Press.
- Huang, F. L. 2017. *Conducting Multilevel Confirmatory Factor Analysis Using R*. 20.
- Imai, T., K. Kuwahara, T. Miyamoto, H. Okazaki, A. Nishihara, I. Kabe, T. Mizoue, and S. Dohi. & Group, J. E. C. on O. H. S 2016. "Validity and Reproducibility of Self-Reported Working Hours among Japanese Male Employees." *Journal of Occupational Health* 58: 340–346.

- Jamal, M. 2011. *Job Stress, Job Performance and Organizational Commitment in a Multinational Company: An Empirical Study in Two Countries*, International Journal of Business and Social Science. 2 (20): 10.
- Keles, H., A. Ekici, M. Ekici, E. Bulcun, and V. Altinkaya. 2007. "Effect of Chronic Diseases and Associated Psychological Distress on Health-Related Quality of Life." *Internal Medicine Journal* 37 (1): 6–11. doi:[10.1111/j.1445-5994.2006.01215.x](https://doi.org/10.1111/j.1445-5994.2006.01215.x)
- Kimball, A., and V. J. Freysinger. 2003. "Leisure, Stress, and Coping: The Sport Participation of Collegiate Student-Athletes." *Leisure Sciences* 25 (2-3): 115–141. doi:[10.1080/01490400306569](https://doi.org/10.1080/01490400306569)
- King, Z. D., J. Moskowitz, B. Egilmez, S. Zhang, L. Zhang, M. Bass, J. Rogers, R. Ghaffari, L. Wakschlag, and N. Alshurafa. 2019. "micro-Stress EMA: A Passive Sensing Framework for Detecting in-the-Wild Stress in Pregnant Mothers." *Proceedings of the ACM on Interactive, Mobile, Wearable and Ubiquitous Technologies* 3 (3): 91. doi:[10.1145/3351249](https://doi.org/10.1145/3351249)
- Lazarus, R. S. 2001. "Relational Meaning and Discrete Emotions." In *Appraisal Processes in Emotion: Theory, Methods, Research*, 37–67. Oxford University Press.
- Lee, K., C. Suh, J.-E. Kim, and J. O. Park. 2017. "The Impact of Long Working Hours on Psychosocial Stress Response among White-Collar Workers." *Industrial Health* 55 (1): 46–53. doi:[10.2486/ind-health.2015-0173](https://doi.org/10.2486/ind-health.2015-0173)
- Lorig, K. R., D. S. Sobel, A. L. Stewart, B. W. Brown, A. Bandura, P. Ritter, V. M. Gonzalez, D. D. Laurent, and H. R. Holman. 1999. "Evidence Suggesting That a Chronic Disease Self-Management Program Can Improve Health Status While Reducing Hospitalization: A Randomized Trial." *Medical Care* 37 (1): 5–14. doi:[10.1097/00005650-199901000-00003](https://doi.org/10.1097/00005650-199901000-00003)
- Lourenco, M. P. C. G., J. Simoes, J. W. S. Vlaeyen, and R. F. F. Cima. 2022. "The Daily Experience of Subjective Tinnitus: Ecological Momentary Assessment versus End-of-Day Diary." *Ear and Hearing* 43 (1): 45–52. doi:[10.1097/AUD.0000000000001054](https://doi.org/10.1097/AUD.0000000000001054)
- Mark, G., and A. Smith. 2018. "A Qualitative Study of Stress in University Staff." *Advances in Social Sciences Research Journal* 5 (2): 238–247. doi:[10.14738/assrj.52.4195](https://doi.org/10.14738/assrj.52.4195)
- Maruyama, S., and K. Morimoto. 1996. "Effects of Long Workhours on Life-Style, Stress and Quality of Life among Intermediate Japanese Managers." *Scandinavian Journal of Work, Environment & Health* 22: 353–359.
- Mazzola, J. J., I. S. Schonfeld, and P. E. Spector. 2011. "What Qualitative Research Has Taught us about Occupational Stress." *Stress and Health: Journal of the International Society for the Investigation of Stress* 27 (2): 93–110. doi:[10.1002/smi.1386](https://doi.org/10.1002/smi.1386)
- Meijman, T. F., and G. Mulder. 1998. "Psychological Aspects of Workload." In *Handbook of Work and Organizational: Work Psychology*, Vol. 2, 2nd ed, pp. 5–33. Psychology Press/Erlbaum (UK) Taylor & Francis.
- Muthén, B. O. 1989. "Latent Variable Modeling in Heterogeneous Populations." *Psychometrika* 54 (4): 557–585. doi:[10.1007/BF02296397](https://doi.org/10.1007/BF02296397)
- Muthén, L. K., and B. O. Muthén. 1998. *Mplus User's Guide*. Los Angeles, CA: Muthén & Muthén, 2017.
- Muthén, B., and T. Asparouhov. 2012. "Bayesian structural equation modeling: A more flexible representation of substantive theory." *Psychological Methods* 17 (3): 313.
- Podsakoff, N. P., T. M. Spoelma, N. Chawla, and A. S. Gabriel. 2019. "What Predicts within-Person Variance in Applied Psychology Constructs? An Empirical Examination." *The Journal of Applied Psychology* 104 (6): 727–754.
- Preacher, K. J., M. J. Zyphur, and Z. Zhang. 2010. "A General Multilevel SEM Framework for Assessing Multilevel Mediation." *Psychological Methods* 15 (3): 209–233. doi:[10.1037/a0020141](https://doi.org/10.1037/a0020141)
- Pyatak, E. A., R. Hernandez, L. Pham, K. Mehdiyeva, S. Schneider, A. Peters, V. Ruelas, et al. 2021. *Function and Emotion in Everyday Life with Type 1 Diabetes (FEEL-T1D): A fully remote intensive longitudinal study of blood glucose, function, and emotional well-being in adults with type 1 diabetes (Preprint)* [Preprint]. JMIR Research Protocols. doi:[10.2196/preprints.30901](https://doi.org/10.2196/preprints.30901)
- Raghupathi, W., and V. Raghupathi. 2018. "An Empirical Study of Chronic Diseases in the United States: A Visual Analytics Approach to Public Health." *International Journal of Environmental Research and Public Health* 15 (3): 431. doi:[10.3390/ijerph15030431](https://doi.org/10.3390/ijerph15030431)

- Revelle, W., and M. W. Revelle. 2015. Package 'psych.' *The Comprehensive R Archive Network* 337: 338.
- Rosen, C. C., C.-H. Chang, E. Djurdjevic, and E. Eatough. 2010. "Occupational Stressors and Job Performance: An Updated Review and Recommendations." In *New Developments in Theoretical and Conceptual Approaches to Job Stress*, edited by P. L. Perrewé and D. C. Ganster, Vol. 8, pp. 1–60. Emerald Group Publishing Limited. doi:[10.1108/S1479-3555\(2010\)000008004](https://doi.org/10.1108/S1479-3555(2010)000008004)
- Sadikaj, G., A. G. Wright, D. M. Dunkley, D. C. Zuroff, and D. S. Moskowitz. 2021. "Multilevel Structural Equation Modeling for Intensive Longitudinal Data: A Practical Guide for Personality Researchers." In *The Handbook of Personality Dynamics and Processes*, 855–885. Elsevier.
- Sato, Y., H. Miyake, and G. Thériault. 2009. "Overtime Work and Stress Response in a Group of Japanese Workers." *Occupational Medicine (Oxford, England)* 59 (1): 14–19. doi:[10.1093/occmed/kqn141](https://doi.org/10.1093/occmed/kqn141)
- Schokker, M. C., T. P. Links, M. L. Luttik, and M. Hagedoorn. 2010. "The Association between Regulatory Focus and Distress in Patients with a Chronic Disease: The Moderating Role of Partner Support." *British Journal of Health Psychology* 15 (Pt 1): 63–78. doi:[10.1348/135910709X429091](https://doi.org/10.1348/135910709X429091)
- Shiffman, S., A. A. Stone, and M. R. Hufford. 2008. "Ecological Momentary Assessment." *Annual Review of Clinical Psychology* 4: 1–32.
- Sianoja, Marjaana, Christine J. Syrek, Jessica de Bloom, Kalevi Korpela, and Ulla Kinnunen, Link to external site, this link will open in a new window, de Bloom, J., Korpela, K., Link to external site, this link will open in a new window, Kinnunen, U., & Link to external site, this link will open in a new window. 2018. "Enhancing Daily Well-Being at Work through Lunchtime Park Walks and Relaxation Exercises: Recovery Experiences as Mediators." *Journal of Occupational Health Psychology* 23 (3): 428–442. doi:[10.1037/ocp0000083](https://doi.org/10.1037/ocp0000083)
- Sonnentag, S., and C. Fritz. 2007. "The Recovery Experience Questionnaire: Development and Validation of a Measure for Assessing Recuperation and Unwinding from Work." *Journal of Occupational Health Psychology* 12 (3): 204–221. doi:[10.1037/1076-8998.12.3.204](https://doi.org/10.1037/1076-8998.12.3.204)
- Sonnentag, S., L. Venz, and A. Casper. 2017. "Advances in Recovery Research: What Have we Learned? What Should Be Done Next?" *Journal of Occupational Health Psychology* 22 (3): 365–380. doi:[10.1037/ocp0000079](https://doi.org/10.1037/ocp0000079)
- Sparks, K., C. Cooper, Y. Fried, and A. Shirom. 2001. "The Effects of Hours of Work on Health: A Meta-Analytic Review." In *Managerial, Occupational and Organizational Stress Research*. Routledge.
- Stebbins, R. A. 1997. "Casual Leisure: A Conceptual Statement." *Leisure Studies* 16 (1): 17–25. doi:[10.1080/026143697375485](https://doi.org/10.1080/026143697375485)
- Stoelting, R. 2002. Structural equation modeling/path analysis. Retrieved March, 16 2008.
- Tindle, J., and P. Tadi. 2021. Neuroanatomy, Parasympathetic Nervous System. In *StatPearls*. StatPearls Publishing. <http://www.ncbi.nlm.nih.gov/books/NBK553141/>
- Todd, S. R., B. N. Fahy, J. L. Paukert, D. Mersinger, M. L. Johnson, and B. L. Bass. 2010. "How Accurate Are Self-Reported Resident Duty Hours?" *Journal of Surgical Education* 67 (2): 103–107.
- Trenberth, L., and P. Dewe. 2002. "The Importance of Leisure as a Means of Coping with Work Related Stress: An Exploratory Study." *Counselling Psychology Quarterly* 15 (1): 59–72. doi:[10.1080/09515070110103999](https://doi.org/10.1080/09515070110103999)
- Trull, T. J., and U. Ebner-Priemer. 2013. "Ambulatory Assessment." *Annual Review of Clinical Psychology* 9: 151–176. doi:[10.1146/annurev-clinpsy-050212-185510](https://doi.org/10.1146/annurev-clinpsy-050212-185510)
- Tung, N. Y. C., Y. Yap, B. Bei, L. J. Luecken, and J. F. Wiley. 2022. "A 14-Day Ecological Momentary Assessment Study on Whether Resilience and Early Family Risk Moderate Daily Stress and Affect on Cortisol Diurnal Slope." *Scientific Reports* 12 (1). Article 1. doi:[10.1038/s41598-022-05277-w](https://doi.org/10.1038/s41598-022-05277-w)
- Winwood, P. C., A. B. Bakker, and A. H. Winefield. 2007. "An Investigation of the Role of Non-Work-Time Behavior in Buffering the Effects of Work Strain." *Journal of Occupational and Environmental Medicine* 49 (8): 862–871. doi:[10.1097/JOM.0b013e318124a8dc](https://doi.org/10.1097/JOM.0b013e318124a8dc)
- Zawadzki, M. J., J. M. Smyth, and H. J. Costigan. 2015. "Real-Time Associations between Engaging in Leisure and Daily Health and Well-Being." *Annals of Behavioral Medicine : a Publication of the Society of Behavioral Medicine* 49 (4): 605–615. doi:[10.1007/s12160-015-9694-3](https://doi.org/10.1007/s12160-015-9694-3)