

Employment Characteristics and Tobacco Product Use, U.S., 2021



Christine M. Kava, PhD,^{1,2} Girija Syamlal, MPH, MBBS,³ Brenna VanFrank, MD,⁴
David A. Siegel, MD,² S. Jane Henley, MSPH,² Jonathan Bryant-Genevier, PhD,² Jin Qin, ScD,²
Susan A. Sabatino, MD²

Introduction: Over 30 million U.S. working adults use tobacco, and tobacco use varies by occupation. Limited information is available on employment characteristics and tobacco use prevalence. The purpose of this study was to describe the prevalence of current tobacco use by employment characteristics and occupation group among U.S. working adults.

Methods: This cross-sectional study used 2021 National Health Interview Survey data for currently working adults (n=16,461) analyzed in 2023. Multivariable logistic regression was used to estimate adjusted odds of tobacco use by employment characteristics and occupation group.

Results: In 2021, 20.0% of working adults used tobacco. Any tobacco use was significantly lower among workers who were offered workplace health insurance (AOR=0.86, 95% CI=0.77–0.97), had paid sick leave (AOR=0.81, 95% CI=0.73–0.91), and government versus private employment (AOR=0.61, 95% CI=0.52–0.70). Any tobacco use was significantly higher among workers who usually worked ≥ 35 hours per week versus did not usually work ≥ 35 hours per week (AOR=1.21, 95% CI=1.06–1.39), worked a rotating or “some other” shift versus daytime shift (AOR=1.19, 95% CI=1.02–1.38), experienced schedule instability (AOR=1.17, 95% CI=1.03–1.31), and worked while physically ill in the past 3 months (AOR=1.25, 95% CI=1.11–1.41). Tobacco use by employment characteristics also varied by occupation group.

Conclusions: Current tobacco use varied according to employment characteristics and occupation group. Findings from this study could inform workplace tobacco cessation interventions and policies (e.g., access to paid sick leave or insurance coverage) to better support tobacco cessation and overall worker health.

Am J Prev Med 2024;67(3):423–433. Published by Elsevier Inc.

INTRODUCTION

Commercial tobacco product use (tobacco use) is the leading preventable cause of disease and death in the U.S.¹ While adult cigarette smoking has declined over the last decade, from 19.0% in 2011² to 11.5% in 2021,³ newer products, such as e-cigarettes, have entered the market and increased in prevalence.^{4,5} In October 2023, approximately 60% of the U.S. civilian, noninstitutionalized population was employed.⁶ The average annual prevalence of any tobacco use among working adults between 2014 and 2016 was 22%.⁷ Disparities existed by occupation, ranging from 9% in life, physical, and social science occupations to 37% in construction and extraction occupations.⁷

Cigarette smoking is higher among workers with higher psychological work demands,⁸ job insecurity,^{9,10} and temporary work status,¹⁰ and is associated with greater absence from work due to sickness.^{11–13} There

From the ¹Epidemic Intelligence Service, Centers for Disease Control and Prevention (CDC), Atlanta, Georgia; ²Division of Cancer Prevention and Control, National Center for Chronic Disease Prevention and Health Promotion, CDC, Atlanta, Georgia; ³Respiratory Health Division, National Institute for Occupational Safety and Health, Morgantown, West Virginia; and ⁴Office on Smoking and Health, CDC, Atlanta, Georgia

Address correspondence to: Christine M. Kava, PhD, MA, Division of Cancer Prevention and Control, Centers for Disease Control and Prevention, 4770 Buford Hwy NE, Atlanta GA 30341. E-mail: tqz1@cdc.gov.

0749-3797/\$36.00

<https://doi.org/10.1016/j.amepre.2024.05.001>

are limited data on how employment characteristics are associated with other forms of tobacco product use and how these associations might differ by occupation. Given changing patterns of tobacco use and disparities observed by occupation, further research in these areas could inform more focused efforts to increase cessation. This cross-sectional study describes the prevalence of current tobacco use by employment characteristics and occupation group among U.S. working adults.

METHODS

The National Health Interview Survey (NHIS) is a nationally representative, cross-sectional household interview survey.¹⁴ This study used 2021 NHIS data from employed adults (N=16,461), defined as respondents who: (1) worked for pay in the last week; (2) did not work for pay in the last week, but had a job or business in the last week and were temporarily absent due to illness, vacation, family or maternity leave, or some other reason; (3) performed seasonal or contract work in the past 12 months; or (4) worked at a family-owned job or business but not for pay.¹⁴ Institutional Review Board approval was exempted under the Common Rule (45 CFR §46) because the study used de-identified and publicly available data.

The main outcome was any tobacco use and was defined as current use of one or more of the following: cigarettes; regular cigars, cigarillos, or little filtered cigars (cigars); e-cigarettes; regular pipe, water pipe, or hookah (pipes); and smokeless tobacco. This study also examined use of each individual product, dual use of

cigarettes and e-cigarettes, and multiproduct use (i.e., use of 2 or more tobacco products).

The following employment characteristics were measured as dichotomous (yes versus no): workplace health insurance offered; paid sick leave; full-time status (i.e., usually worked ≥ 35 hours per week); experienced job insecurity; experienced work–life imbalance; experienced schedule instability; missed work in the past 12 months due to illness, injury, or disability; and worked while physically ill in the past 3 months. Employment type (private; government; self-employed or working at a family-owned job or business but not for pay [other type]) and work shift (daytime; evening/night; rotating or “some other” shift) were also examined.

Workers were grouped into 1 of 5 occupation groups based on the 2018 U.S. Bureau of Labor Statistics Standard Occupational Classification and Coding Structure¹⁵: management, business, science, and arts; service; sales and office; natural resources, construction, and maintenance; and production, transportation, and material moving (see [Appendix Table 1](#), available online). See table footnotes for detailed information on the measures.

Data analysis was conducted in SAS 9.4 (SAS Institute Inc.) and used the survey weights and design variables to account for the complex sample design and produce national estimates. The prevalence of current tobacco use by employment characteristics and occupation group was examined. 95% CIs were produced to allow for informal, conservative comparisons of prevalence estimates. Multivariable logistic regression was used to estimate adjusted odds of any tobacco use by

Table 1. Sample and Population Estimates, Working Adults, U.S., 2021

Variable	n ^a	N x 1,000 ^b	% (95% CI) ^b
Total	16,125	149,039	61.7 (61.0–62.5)
Occupation group ^c			
Management, business, science, and arts ^d	7,556	64,645	43.4 (42.3–44.5)
Service ^e	2,318	23,329	15.7 (14.9–16.4)
Sales and office ^f	3,143	29,239	19.6 (18.9–20.4)
Natural resources, construction, and maintenance ^g	1,270	12,946	8.7 (8.1–9.2)
Production, transportation, and material moving ^h	1,838	18,880	12.7 (11.9–13.4)
Employment characteristics ⁱ			
Workplace health insurance offered ^j			
No	4,500	43,905	29.6 (28.6–30.6)
Yes	11,576	104,489	70.4 (69.4–71.4)
Paid sick leave available ^k			
No	4,803	46,215	31.3 (30.4–32.3)
Yes	11,183	101,331	68.7 (67.7–69.6)
Usually worked ≥ 35 hours per week ^l			
No	3,069	29,312	19.7 (18.8–20.5)
Yes	13,042	119,619	80.3 (79.5–81.2)

(continued on next page)

Table 1. Sample and Population Estimates, Working Adults, U.S., 2021 (continued)

Variable	n ^a	N x 1,000 ^b	% (95% CI) ^b
Employment type ^m			
Private	11,546	109,018	73.5 (72.5–74.5)
Government	2,587	22,311	15.0 (14.3–15.8)
Other type	1,912	16,993	11.5 (10.8–12.1)
Work shift ⁿ			
Daytime	12,971	117,909	79.3 (78.5–80.1)
Evening or night	1,304	13,491	9.1 (8.5–9.7)
Rotating or “some other”	1,804	17,245	11.6 (11.0–12.2)
Experienced job insecurity ^o			
No	13,670	126,319	85.8 (85.0–86.5)
Yes	2,248	20,929	14.2 (13.5–15.0)
Experienced work–life imbalance ^p			
No	12,662	116,372	78.5 (77.7–79.3)
Yes	3,371	31,883	21.5 (20.7–22.3)
Experienced schedule instability ^q			
No	12,974	118,824	80.0 (79.2–80.8)
Yes	3,092	29,702	20.0 (19.2–20.8)
Missed work in past 12 months due to illness, injury, or disability ^r			
No	9,422	87,516	59.1 (58.2–60.0)
Yes	6,594	60,533	40.9 (40.0–41.8)
Worked while physically ill in past 3 months ^s			
No	12,946	119,614	81.5 (80.7–82.2)
Yes	2,934	27,190	18.5 (17.8–19.3)

Note: Frequencies may not add to totals due to missing data.

^aNumber of working adults in the 2021 National Health Interview Survey dataset.

^bNational estimates for the number and percentage of working adults in the U.S.

^cBased on the 2018 U.S. Bureau of Labor Statistics Standard Occupational Classification and Coding Structure. Workers that were missing data for occupation (n=328) and in the military specific occupation group (n=8) were excluded from analysis.

^dIncludes architecture and engineering; arts, design, entertainment, sports, and media; business and financial operations; community and social services; computer and mathematical; education, training, and library; healthcare practitioners and technical; legal; life, physical, and social science; and management occupations.

^eIncludes building and grounds cleaning and maintenance; food preparation and serving related; healthcare support; personal care and service; and protective service occupations.

^fIncludes office and administrative support; and sales and related occupations.

^gIncludes construction and extraction; farming, fishing, and forestry; and installation, maintenance, and repair occupations.

^hIncludes production; and transportation and material moving occupations.

ⁱEach employment characteristic was measured with one question (described in detail in the footnotes below). For all measures, “refused” and “don’t know” responses were coded as missing.

^jWorkers asked whether health insurance was offered to them through their workplace (yes or no). This measure is distinct from worker health insurance status; it is possible for the worker to have answered yes to this question without being covered by the insurance offered.

^kWorkers asked whether they had paid sick leave available if needed (yes or no).

^lWorkers asked whether they usually worked 35 hours or more per week in total at all jobs or businesses (yes or no).

^mWorkers asked what category best described their main job (employee of a private company for wages; a federal government employee; a state government employee; a local government employee; self-employed in own business, professional practice, or farm; working without pay in a family-owned business or farm). Federal, state, and local government workers were combined into one category. Workers who were self-employed or working without pay were coded as “other type.”

ⁿWorkers asked what category best described their usual hours of work at their main job (daytime shift; evening shift; night shift; rotating shift; some other shift). Workers with evening and night shifts were combined into one category. Workers with a rotating or “some other” shift were combined into one category.

^oWorkers asked how likely it was that they would lose their job or be laid off in the next 12 months (very likely; fairly likely; somewhat likely; not at all likely). Workers who responded very likely, fairly likely, or somewhat likely were coded as “yes” for job insecurity.

^pWorkers asked how easy or difficult it was for them to change their work schedule to do things that were important to them or their family (very easy; somewhat easy; somewhat difficult; very difficult). Workers who responded somewhat difficult or very difficult were coded as “yes” for work–life imbalance.

^qWorkers asked whether their work schedule at their main job changed on a regular basis (yes or no). Workers who responded yes were coded as “yes” for schedule instability.

^rWorkers asked how many days of work they missed in the past 12 months because of an illness, injury, or disability. Workers who missed ≥1 days were coded as “yes.”

^sWorkers asked how many days they worked while physically ill in the past 3 months. Workers who worked ≥1 days while ill were coded as “yes.”

Table 2. Current Commercial Tobacco Product Use^a by Workers' Employment Characteristics, U.S., 2021

Variable	Current tobacco use, % (95% CI) ^b							
	Any product ^c	Cigarettes	Cigars ^d	E-cigarettes	Pipes ^e	Smokeless	Dual use ^f	≥2 products ^g
Total	20.0 (19.2–20.8)	11.0 (10.4–11.6)	4.1 (3.8–4.5)	5.4 (4.9–5.8)	1.0 (0.8–1.2)	2.7 (2.4–3.0)	1.4 (1.2–1.6)	3.7 (3.3–4.0)
Occupation group ^h								
Management, business, science, and arts ⁱ	14.2 (13.2–15.1)	6.4 (5.7–7.1)	3.4 (2.9–3.8)	4.0 (3.5–4.5)	1.1 (0.8–1.3)	1.7 (1.4–2.1)	0.8 (0.6–1.0)	2.1 (1.7–2.4)
Service ^j	22.2 (20.1–24.2)	14.5 (12.8–16.1)	3.9 (2.9–4.8)	6.7 (5.4–8.0)	0.8 (0.4–1.3)	1.8 (1.2–2.4)	2.2 (1.4–3.0)	4.9 (3.8–5.9)
Sales and office ^k	18.6 (17.0–20.2)	9.5 (8.3–10.6)	4.1 (3.4–4.9)	6.0 (5.0–7.1)	1.2 (0.7–1.7)	1.7 (1.1–2.2)	1.3 (0.8–1.8)	3.6 (2.7–4.4)
Natural resources, construction, and maintenance ^l	36.0 (32.9–39.1)	21.9 (19.2–24.5)	6.9 (5.2–8.6)	6.0 (4.4–7.5)	1.1 (0.5–1.6)	8.9 (7.1–10.7)	2.1 (1.2–3.1)	7.5 (5.8–9.1)
Production, transportation, and material moving ^m	28.6 (26.3–30.9)	17.4 (15.5–19.3)	5.2 (4.1–6.4)	7.0 (5.6–8.5)	ES	4.5 (3.4–5.6)	2.2 (1.4–3.0)	5.4 (4.2–6.5)
Employment characteristic ⁿ								
Workplace health insurance offered ^o								
No	21.7 (20.3–23.2)	12.6 (11.5–13.7)	3.7 (3.1–4.4)	6.1 (5.3–7.0)	1.2 (0.8–1.6)	2.2 (1.7–2.7)	1.5 (1.1–1.9)	3.5 (2.9–4.2)
Yes	19.3 (18.4–20.2)	10.3 (9.6–11.0)	4.3 (3.9–4.8)	5.0 (4.6–5.5)	1.0 (0.8–1.2)	3.0 (2.6–3.3)	1.4 (1.1–1.6)	3.7 (3.3–4.1)
Paid sick leave available ^p								
No	23.1 (21.7–24.6)	13.9 (12.8–15.0)	4.1 (3.4–4.8)	6.1 (5.3–6.9)	1.1 (0.7–1.6)	2.5 (2.0–3.0)	1.8 (1.3–2.2)	4.2 (3.5–4.8)
Yes	18.6 (17.7–19.5)	9.7 (9.0–10.4)	4.2 (3.7–4.6)	5.0 (4.5–5.5)	1.0 (0.8–1.2)	2.8 (2.4–3.2)	1.3 (1.0–1.5)	3.5 (3.1–3.9)
Usually worked ≥35 hours per week ^q								
No	16.3 (14.8–17.8)	9.7 (8.5–10.8)	2.6 (2.0–3.2)	5.5 (4.6–6.5)	1.0 (0.5–1.5)	0.7 (0.4–1.0)	1.1 (0.7–1.5)	2.7 (2.0–3.3)
Yes	20.9 (20.0–21.8)	11.3 (10.6–12.0)	4.5 (4.1–5.0)	5.3 (4.8–5.8)	1.0 (0.8–1.2)	3.2 (2.8–3.6)	1.5 (1.2–1.8)	3.9 (3.5–4.3)
Employment type ^r								
Private	21.1 (20.1–22.1)	11.8 (11.1–12.5)	4.1 (3.7–4.6)	6.0 (5.4–6.6)	1.1 (0.9–1.4)	2.6 (2.3–3.0)	1.7 (1.4–1.9)	3.9 (3.5–4.4)
Government	13.4 (11.9–15.0)	6.8 (5.8–7.8)	3.2 (2.4–4.0)	2.7 (2.0–3.4)	ES	3.2 (2.4–4.0)	0.6 (0.3–0.9)	2.6 (1.8–3.3)
Other type	21.4 (19.3–23.6)	11.5 (9.8–13.2)	5.4 (4.2–6.5)	4.5 (3.4–5.6)	ES	2.8 (2.0–3.7)	0.9 (0.5–1.4)	3.3 (2.4–4.2)
Work shift ^s								
Daytime	19.1 (18.3–20.0)	10.3 (9.7–11.0)	4.0 (3.5–4.4)	4.8 (4.3–5.2)	0.9 (0.7–1.0)	2.8 (2.5–3.2)	1.2 (0.9–1.4)	3.1 (2.8–3.5)
Evening or night	23.5 (20.8–26.2)	14.9 (12.6–17.2)	4.0 (2.8–5.3)	8.0 (6.2–9.8)	1.6 (0.7–2.5)	2.0 (1.1–2.9)	3.1 (1.8–4.3)	6.3 (4.6–8.0)
Rotating or “some other”	23.2 (20.8–25.6)	12.5 (10.7–14.4)	5.6 (4.4–6.7)	7.5 (5.8–9.1)	1.7 (1.0–2.5)	2.7 (1.8–3.5)	1.9 (1.2–2.6)	5.6 (4.4–6.8)
Experienced job insecurity ^t								
No	19.6 (18.8–20.5)	10.5 (9.8–11.1)	4.2 (3.8–4.6)	5.2 (4.7–5.7)	0.9 (0.7–1.1)	2.8 (2.5–3.2)	1.3 (1.1–1.6)	3.5 (3.1–3.9)
Yes	22.2 (20.1–24.4)	13.7 (12.1–15.4)	3.8 (2.9–4.8)	6.5 (5.2–7.8)	1.7 (1.0–2.5)	2.0 (1.4–2.6)	1.9 (1.2–2.7)	4.7 (3.6–5.8)
Experienced work–life imbalance ^u								
No	19.6 (18.7–20.4)	10.5 (9.9–11.2)	4.3 (3.8–4.7)	5.1 (4.6–5.6)	1.0 (0.8–1.2)	2.7 (2.3–3.0)	1.3 (1.1–1.5)	3.5 (3.1–3.8)
Yes	21.4 (19.8–23.1)	12.5 (11.2–13.8)	3.6 (2.9–4.4)	6.4 (5.4–7.5)	1.1 (0.7–1.4)	2.9 (2.2–3.5)	1.9 (1.3–2.5)	4.5 (3.6–5.3)

(continued on next page)

Table 2. Current Commercial Tobacco Product Use^a by Workers' Employment Characteristics, U.S., 2021 (*continued*)

Variable	Current tobacco use, % (95% CI) ^b							
	Any product ^c	Cigarettes	Cigars ^d	E-cigarettes	Pipes ^e	Smokeless	Dual use ^f	≥2 products ^g
Experienced schedule instability ^v								
No	19.3 (18.5–20.1)	10.6 (10.0–11.3)	3.9 (3.5–4.3)	5.0 (4.6–5.5)	1.0 (0.7–1.2)	2.6 (2.3–3.0)	1.3 (1.0–1.5)	3.4 (3.0–3.8)
Yes	22.8 (20.9–24.7)	12.4 (11.0–13.8)	5.3 (4.4–6.2)	6.7 (5.6–7.8)	1.3 (0.8–1.7)	3.0 (2.3–3.7)	1.9 (1.3–2.5)	4.8 (3.9–5.7)
Missed work in past 12 months due to illness, injury, or disability ^w								
No	19.0 (18.0–20.1)	10.6 (9.9–11.4)	4.2 (3.7–4.7)	4.7 (4.2–5.2)	1.0 (0.8–1.2)	2.5 (2.1–2.9)	1.3 (1.0–1.6)	3.4 (2.9–3.8)
Yes	21.3 (20.1–22.5)	11.5 (10.6–12.3)	4.1 (3.5–4.6)	6.3 (5.6–7.1)	1.1 (0.8–1.4)	3.1 (2.6–3.6)	1.5 (1.2–1.9)	4.1 (3.6–4.7)
Worked while physically ill in past 3 months ^x								
No	18.8 (18.0–19.7)	10.2 (9.6–10.9)	4.0 (3.6–4.3)	4.8 (4.4–5.3)	1.0 (0.8–1.2)	2.6 (2.3–2.9)	1.2 (1.0–1.4)	3.3 (2.9–3.6)
Yes	25.1 (23.1–27.1)	14.1 (12.7–15.5)	4.9 (3.9–6.0)	7.8 (6.6–9.1)	1.2 (0.7–1.7)	3.1 (2.4–3.9)	2.2 (1.6–2.9)	5.4 (4.4–6.5)

^aDefined as having ever used the tobacco product (for cigarettes, ever smoked at least 100 cigarettes in lifetime) and current use every day or some days.

^bEstimates are weighted to represent the national population. Estimates with relative standard errors ≥30% were suppressed.

^cCurrent use of ≥1 of the following products: cigarettes; regular cigars, cigarillos, or little filtered cigars (cigars); e-cigarettes; regular pipe, water pipe, or hookah (pipes); smokeless tobacco.

^dIncludes regular cigars, cigarillos, and little filtered cigars.

^eIncludes regular pipe, water pipe, and hookah.

^fCurrent use of cigarettes and e-cigarettes.

^gCurrent use of ≥2 of the following products: cigarettes; regular cigars, cigarillos, or little filtered cigars (cigars); e-cigarettes; regular pipe, water pipe, or hookah (pipes); smokeless tobacco.

^hBased on the 2018 U.S. Bureau of Labor Statistics Standard Occupational Classification and Coding Structure. Workers that were missing data for occupation (n=328) and in the military specific occupation group (n=8) were excluded from analysis.

ⁱIncludes architecture and engineering; arts, design, entertainment, sports, and media; business and financial operations; community and social services; computer and mathematical; education, training, and library; healthcare practitioners and technical; legal; life, physical, and social science; and management occupations.

^jIncludes building and grounds cleaning and maintenance; food preparation and serving related; healthcare support; personal care and service; and protective service occupations.

^kIncludes office and administrative support; and sales and related occupations.

^lIncludes construction and extraction; farming, fishing, and forestry; and installation, maintenance, and repair occupations.

^mIncludes production; and transportation and material moving occupations.

ⁿEach employment characteristic was measured with one question (described in detail in the footnotes below). For all measures, "refused" and "don't know" responses were coded as missing.

^oWorkers asked whether health insurance was offered to them through their workplace (yes or no). This measure is distinct from worker health insurance status; it is possible for the worker to have answered yes to this question without being covered by the insurance offered.

^pWorkers asked whether they had paid sick leave available if needed (yes or no).

^qWorkers asked whether they usually worked 35 hours or more per week in total at all jobs or businesses (yes or no).

^rWorkers asked what category best described their main job (employee of a private company for wages; a federal government employee; a state government employee; a local government employee; self-employed in own business, professional practice, or farm; working without pay in a family-owned business or farm). Federal, state, and local government workers were combined into one category. Workers who were self-employed or working without pay were coded as "other type."

^sWorkers asked what category best described their usual hours of work at their main job (daytime shift; evening shift; night shift; rotating shift; some other shift). Workers with evening and night shifts were combined into one category. Workers with a rotating or "some other" shift were combined into one category.

^tWorkers asked how likely it was that they would lose their job or be laid off in the next 12 months (very likely; fairly likely; somewhat likely; not at all likely). Workers who responded very likely, fairly likely, or somewhat likely were coded as "yes" for job insecurity.

^uWorkers asked how easy or difficult it was for them to change their work schedule to do things that were important to them or their family (very easy; somewhat easy; somewhat difficult; very difficult). Workers who responded somewhat difficult or very difficult were coded as "yes" for work-life imbalance.

^vWorkers asked whether their work schedule at their main job changed on a regular basis (yes or no). Workers who responded yes were coded as "yes" for schedule instability.

^wWorkers asked how many days of work they missed in the past 12 months because of an illness, injury, or disability. Workers who missed ≥1 days were coded as "yes."

^xWorkers asked how many days they worked while physically ill in the past 3 months. Workers who worked ≥ 1days while ill were coded as "yes."

ES, estimate suppressed.

employment characteristics; separate models were run for each employment characteristic.

RESULTS

Of the estimated 149 million working adults in the U.S. (Table 1), 20.0% (95% CI=19.2–20.8) currently used tobacco (Table 2). By occupation group and employment characteristics, any tobacco use was highest among workers in the natural resources, construction, and maintenance occupation group and who worked while physically ill in the past 3 months (46.3%; 95% CI=38.1–54.5) (Table 3).

Adjusted odds of any tobacco use (Table 4) were significantly lower among workers who were offered health insurance at their workplace (AOR=0.86; 95% CI=0.77–0.97); had paid sick leave (AOR=0.81; 95% CI=0.73–0.91); and government versus private employment

(AOR=0.61; 95% CI=0.52–0.70). Odds were significantly higher among workers who usually worked ≥ 35 hours per week (AOR=1.21; 95% CI=1.06–1.39); worked a rotating or “some other” shift versus daytime shift (AOR=1.19; 95% CI=1.02–1.38); experienced schedule instability (AOR=1.17; 95% CI=1.03–1.31); and worked while physically ill in past 3 months (AOR=1.25; 95% CI=1.11–1.41).

In the management, business, science, and arts group, higher odds of any tobacco use were observed among workers who usually worked ≥ 35 hours per week (AOR=1.32; 95% CI=1.02–1.70); worked a rotating or “some other” shift (AOR=1.41; 95% CI=1.10–1.81); experienced schedule instability (AOR=1.29; 95% CI=1.04–1.60); and worked while physically ill in the past 3 months (AOR=1.30; 95% CI=1.09–1.56). Odds were significantly lower among workers in government employment (AOR=0.64; 95% CI=0.52–0.79).

Table 3. Prevalence of Any Current Commercial Tobacco Product Use,^a Employment Characteristics,^b and Occupation,^c U.S., 2021

Variable	Occupation group, % (95% CI) ^d				
	Management, business, science, and arts ^e	Service ^f	Sales and office ^g	Natural resources, construction, and maintenance ^h	Production, transportation, and material moving ⁱ
Workplace health insurance offered ^j					
No	15.8 (13.5–18.1)	22.8 (19.8–25.8)	19.2 (16.1–22.2)	33.4 (28.8–38.1)	25.0 (20.2–29.7)
Yes	13.8 (12.7–14.8)	21.8 (19.0–24.6)	18.3 (16.3–20.2)	38.1 (34.4–41.9)	30.0 (27.2–32.8)
Paid sick leave available ^k					
No	15.9 (13.6–18.2)	24.6 (21.5–27.6)	20.1 (17.0–23.1)	35.6 (31.1–40.0)	26.9 (22.9–30.8)
Yes	13.8 (12.7–14.8)	19.8 (17.1–22.6)	17.8 (15.9–19.7)	36.5 (32.3–40.7)	29.9 (26.9–33.0)
Usually worked ≥ 35 hours per week ^l					
No	11.0 (8.8–13.2)	19.5 (16.5–22.6)	14.5 (11.6–17.5)	32.3 (23.4–41.1)	19.0 (13.8–24.2)
Yes	14.7 (13.6–15.7)	23.6 (20.9–26.3)	19.8 (17.9–21.7)	36.4 (33.1–39.7)	30.7 (28.2–33.3)
Employment type ^m					
Private	15.2 (14.0–16.4)	24.2 (21.6–26.8)	18.8 (17.0–20.6)	35.8 (32.4–39.3)	28.7 (26.2–31.3)
Government	9.9 (8.4–11.5)	16.7 (12.9–20.6)	14.3 (9.8–18.8)	35.7 (24.5–46.9)	24.1 (14.7–33.5)
Other type	15.4 (12.6–18.2)	20.0 (14.9–25.1)	19.5 (14.9–24.1)	36.9 (30.0–43.8)	29.7 (20.5–38.8)
Work shift ⁿ					
Daytime	13.5 (12.5–14.5)	20.9 (18.3–23.4)	18.6 (16.8–20.5)	36.0 (32.6–39.3)	27.7 (25.0–30.5)
Evening or night	18.3 (13.0–23.7)	25.6 (20.4–30.7)	15.1 (9.5–20.6)	36.9 (21.9–51.9)	29.7 (23.9–35.5)
Rotating or “some other”	18.4 (15.0–21.8)	23.4 (18.5–28.3)	20.7 (15.8–25.6)	36.3 (25.7–46.9)	31.4 (24.8–38.1)
Experienced job insecurity ^o					
No	13.8 (12.8–14.8)	21.8 (19.5–24.0)	18.4 (16.7–20.1)	35.4 (32.0–38.8)	29.2 (26.7–31.8)
Yes	16.1 (13.2–19.0)	24.6 (19.2–30.0)	20.5 (15.8–25.2)	37.7 (30.6–44.9)	25.5 (20.0–31.1)

(continued on next page)

Table 3. Prevalence of Any Current Commercial Tobacco Product Use,^a Employment Characteristics,^b and Occupation,^c U.S., 2021 (continued)

Variable	Occupation group, % (95% CI) ^d				
	Management, business, science, and arts ^e	Service ^f	Sales and office ^g	Natural resources, construction, and maintenance ^h	Production, transportation, and material moving ⁱ
Experienced work–life imbalance ^p					
No	14.0 (13.0–15.0)	21.4 (19.1–23.7)	18.1 (16.4–19.7)	37.6 (34.0–41.2)	27.4 (24.7–30.0)
Yes	14.5 (12.5–16.6)	24.7 (20.2–29.2)	20.9 (16.7–25.1)	30.0 (23.7–36.4)	32.2 (27.4–37.0)
Experienced schedule instability ^q					
No	13.5 (12.5–14.5)	20.9 (18.5–23.2)	18.6 (16.8–20.4)	35.6 (32.0–39.2)	28.8 (26.2–31.3)
Yes	17.3 (14.6–20.0)	26.2 (21.9–30.5)	18.7 (14.9–22.6)	37.2 (30.8–43.5)	27.6 (22.6–32.6)
Missed work in past 12 months due to illness, injury, or disability ^r					
No	13.6 (12.4–14.7)	20.4 (17.8–23.1)	17.5 (15.6–19.5)	32.8 (28.8–36.7)	27.8 (24.6–31.1)
Yes	14.8 (13.4–16.3)	24.6 (21.3–28.0)	19.9 (17.4–22.5)	41.9 (37.1–46.8)	30.0 (26.2–33.7)
Worked while physically ill in past 3 months ^s					
No	13.2 (12.2–14.2)	21.1 (18.8–23.3)	17.5 (15.9–19.1)	34.0 (30.8–37.3)	26.5 (24.1–29.0)
Yes	17.9 (15.7–20.2)	27.9 (22.8–32.9)	23.4 (19.2–27.7)	46.3 (38.1–54.5)	38.7 (32.5–44.9)

^aCurrent use of ≥ 1 of the following products: cigarettes; regular cigars, cigarillos, or little filtered cigars (cigars); e-cigarettes; regular pipe, water pipe, or hookah (pipes); smokeless tobacco.

^bEach employment characteristic was measured with one question (described in detail in the footnotes below). For all measures, “refused” and “don’t know” responses were coded as missing.

^cBased on the 2018 U.S. Bureau of Labor Statistics Standard Occupational Classification and Coding Structure. Workers missing data for occupation (n=328) and in the military specific occupation group (n=8) were excluded.

^dEstimates are weighted to represent the national population.

^eIncludes architecture and engineering; arts, design, entertainment, sports, and media; business and financial operations; community and social services; computer and mathematical; education, training, and library; healthcare practitioners and technical; legal; life, physical, and social science; and management occupations.

^fIncludes building and grounds cleaning and maintenance; food preparation and serving related; healthcare support; personal care and service; and protective service occupations.

^gIncludes office and administrative support; and sales and related occupations.

^hIncludes construction and extraction; farming, fishing, and forestry; and installation, maintenance, and repair occupations.

ⁱIncludes production; and transportation and material moving occupations.

^jWorkers asked whether health insurance was offered to them through their workplace (yes or no). This measure is distinct from worker health insurance status; it is possible for the worker to have answered yes to this question without being covered by the insurance offered.

^kWorkers asked whether they had paid sick leave available if needed (yes or no).

^lWorkers asked whether they usually worked 35 hours or more per week in total at all jobs or businesses (yes or no).

^mWorkers asked what category best described their main job (employee of a private company for wages; a federal government employee; a state government employee; a local government employee; self-employed in own business, professional practice, or farm; working without pay in a family-owned business or farm). Federal, state, and local government workers were combined into one category. Workers who were self-employed or working without pay were coded as “other type.”

ⁿWorkers asked what category best described their usual hours of work at their main job (daytime shift; evening shift; night shift; rotating shift; some other shift). Workers with evening and night shifts were combined into one category. Workers with a rotating or “some other” shift were combined into one category.

^oWorkers asked how likely it was that they would lose their job or be laid off in the next 12 months (very likely; fairly likely; somewhat likely; not at all likely). Workers who responded very likely, fairly likely, or somewhat likely were coded as “yes” for job insecurity.

^pWorkers asked how easy or difficult it was for them to change their work schedule to do things that were important to them or their family (very easy; somewhat easy; somewhat difficult; very difficult). Workers who responded somewhat difficult or very difficult were coded as “yes” for work–life imbalance.

^qWorkers asked whether their work schedule at their main job changed on a regular basis (yes or no). Workers who responded yes were coded as “yes” for schedule instability.

^rWorkers asked how many days of work they missed in the past 12 months because of an illness, injury, or disability. Workers who missed ≥ 1 days were coded as “yes.”

^sWorkers asked how many days they worked while physically ill in the past 3 months. Workers who worked ≥ 1 days while ill were coded as “yes.”

Table 4. Odds of Any Current Commercial Tobacco Product Use,^a Employment Characteristics,^b and Occupation,^c U.S., 2021

Variable	Total	Occupation group, AOR (95% CI) ^d				
		Management, business, science, and arts ^e	Service ^f	Sales and office ^g	Natural resources, construction, and maintenance ^h	Production, transportation, and material moving ⁱ
Workplace health insurance offered ^j						
No	Ref	Ref	Ref	Ref	Ref	Ref
Yes	0.86 (0.77–0.97)	0.88 (0.72–1.09)	0.82 (0.63–1.07)	0.88 (0.68–1.15)	0.98 (0.73–1.32)	1.13 (0.82–1.56)
Paid sick leave available ^k						
No	Ref	Ref	Ref	Ref	Ref	Ref
Yes	0.81 (0.73–0.91)	0.92 (0.76–1.13)	0.68 (0.52–0.89)	0.86 (0.67–1.11)	0.89 (0.66–1.21)	1.21 (0.93–1.59)
Usually worked ≥35 hours per week ^l						
No	Ref	Ref	Ref	Ref	Ref	Ref
Yes	1.21 (1.06–1.39)	1.32 (1.02–1.70)	1.10 (0.83–1.47)	1.30 (0.94–1.78)	0.99 (0.64–1.55)	1.69 (1.15–2.47)
Employment type ^m						
Private	Ref	Ref	Ref	Ref	Ref	Ref
Government	0.61 (0.52–0.70)	0.64 (0.52–0.79)	0.50 (0.37–0.68)	0.71 (0.47–1.06)	0.78 (0.45–1.35)	0.78 (0.44–1.40)
Other type	0.98 (0.85–1.13)	0.96 (0.75–1.22)	0.82 (0.56–1.18)	1.06 (0.76–1.48)	1.05 (0.75–1.48)	1.15 (0.70–1.86)
Work shift ⁿ						
Daytime	Ref	Ref	Ref	Ref	Ref	Ref
Evening or night	1.16 (0.98–1.36)	1.28 (0.89–1.86)	1.10 (0.79–1.54)	0.65 (0.40–1.04)	0.82 (0.38–1.75)	1.19 (0.86–1.64)
Rotating or “some other”	1.19 (1.02–1.38)	1.41 (1.10–1.81)	1.00 (0.70–1.41)	1.10 (0.78–1.56)	1.00 (0.60–1.67)	1.18 (0.82–1.69)
Experienced job insecurity ^o						
No	Ref	Ref	Ref	Ref	Ref	Ref
Yes	1.11 (0.97–1.28)	1.09 (0.86–1.39)	1.25 (0.89–1.75)	1.12 (0.82–1.52)	1.26 (0.90–1.78)	0.88 (0.63–1.21)
Experienced work–life imbalance ^p						
No	Ref	Ref	Ref	Ref	Ref	Ref
Yes	1.04 (0.93–1.16)	1.02 (0.85–1.24)	1.05 (0.79–1.38)	1.18 (0.89–1.56)	0.74 (0.52–1.04)	1.13 (0.86–1.48)
Experienced schedule instability ^q						
No	Ref	Ref	Ref	Ref	Ref	Ref
Yes	1.17 (1.03–1.31)	1.29 (1.04–1.60)	1.31 (0.99–1.74)	1.00 (0.74–1.35)	0.98 (0.70–1.36)	0.92 (0.69–1.22)
Missed work in past 12 months due to illness, injury, or disability ^r						
No	Ref	Ref	Ref	Ref	Ref	Ref
Yes	1.04 (0.94–1.14)	1.04 (0.89–1.21)	1.01 (0.78–1.31)	1.07 (0.85–1.34)	1.17 (0.88–1.56)	0.94 (0.71–1.25)
(continued on next page)						

(continued on next page)

Table 4. Odds of Any Current Commercial Tobacco Product Use,^a Employment Characteristics,^b and Occupation,^c U.S., 2021 (*continued*)

Variable	Total	Occupation group, AOR (95% CI) ^d				
		Management, business, science, and arts ^e	Service ^f	Sales and office ^g	Natural resources, construction, and maintenance ^h	Production, transportation, and material moving ⁱ
Worked while physically ill in past 3 months ^s						
No	Ref	Ref	Ref	Ref	Ref	Ref
Yes	1.25 (1.11–1.41)	1.30 (1.09–1.56)	1.21 (0.89–1.66)	1.18 (0.89–1.57)	1.35 (0.91–1.99)	1.44 (1.08–1.92)

Note: Boldface indicates statistical significance ($p < 0.05$).

^aCurrent use of ≥ 1 of the following products: cigarettes; regular cigars, cigarillos, or little filtered cigars (cigars); e-cigarettes; regular pipe, water pipe, or hookah (pipes); smokeless tobacco.

^bEach employment characteristic was measured with one question (described in detail in the footnotes below). For all measures, “refused” and “don’t know” responses were coded as missing.

^cBased on the 2018 U.S. Bureau of Labor Statistics Standard Occupational Classification and Coding Structure. Workers missing data for occupation ($n=328$) and in the military specific occupation group ($n=8$) were excluded.

^dLogistic regression was used to calculate AORs, defined as the adjusted odds of any tobacco use ($=1$) versus no tobacco use ($=0$). Separate models were run for each employment characteristic while adjusting for age group (18–24; 25–34; 35–44; 45–54; 55–64; ≥ 65 years), sex (female; male), race and ethnicity (non-Hispanic Asian; non-Hispanic Black or African American; non-Hispanic White; non-Hispanic American Indian or Alaska Native, alone or with other racial identities, and other single and multiple racial identities [other race]; Hispanic or Latino), sexual orientation (bisexual; gay or lesbian; straight; other sexual orientation), family income as a percentage of the federal poverty level ($<100\%$; ≥ 100 to $<200\%$; $\geq 200\%$ to $<400\%$; $\geq 400\%$), county classification (rural; urban), U.S. census region (Northwest; Midwest; South; West), self-rated health (excellent/very good/good; fair/poor), and whether the worker had ever been diagnosed with any type of depression (yes; no). County classification was defined based on the 2013 National Center for Health Statistics Urban-Rural Classification Scheme for Counties. Counties designated as large central metropolitan, large fringe metropolitan, medium metropolitan, and small metropolitan were coded as urban, while counties designated as micropolitan or noncore were coded as rural.

^eIncludes architecture and engineering; arts, design, entertainment, sports, and media; business and financial operations; community and social services; computer and mathematical; education, training, and library; healthcare practitioners and technical; legal; life, physical, and social science; and management occupations.

^fIncludes building and grounds cleaning and maintenance; food preparation and serving related; healthcare support; personal care and service; and protective service occupations.

^gIncludes office and administrative support; and sales and related occupations.

^hIncludes construction and extraction; farming, fishing, and forestry; and installation, maintenance, and repair occupations.

ⁱIncludes production; and transportation and material moving occupations.

^jWorkers asked whether health insurance was offered to them through their workplace (yes or no). This measure is distinct from worker health insurance status; it is possible for the worker to have answered yes to this question without being covered by the insurance offered.

^kWorkers asked whether they had paid sick leave available if needed (yes or no).

^lWorkers asked whether they usually worked 35 hours or more per week in total at all jobs or businesses (yes or no).

^mWorkers asked what category best described their main job (employee of a private company for wages; a federal government employee; a state government employee; a local government employee; self-employed in own business, professional practice, or farm; working without pay in a family-owned business or farm). Federal, state, and local government workers were combined into one category. Workers who were self-employed or working without pay were coded as “other type.”

ⁿWorkers asked what category best described their usual hours of work at their main job (daytime shift; evening shift; night shift; rotating shift; some other shift). Workers with evening and night shifts were combined into one category. Workers with a rotating or “some other” shift were combined into one category.

^oWorkers asked how likely it was that they would lose their job or be laid off in the next 12 months (very likely; fairly likely; somewhat likely; not at all likely). Workers who responded very likely, fairly likely, or somewhat likely were coded as “yes” for job insecurity.

^pWorkers asked how easy or difficult it was for them to change their work schedule to do things that were important to them or their family (very easy; somewhat easy; somewhat difficult; very difficult). Workers who responded somewhat difficult or very difficult were coded as “yes” for work–life imbalance.

^qWorkers asked whether their work schedule at their main job changed on a regular basis (yes or no). Workers who responded yes were coded as “yes” for schedule instability.

^rWorkers asked how many days of work they missed in the past 12 months because of an illness, injury, or disability. Workers who missed ≥ 1 days were coded as “yes.”

^sWorkers asked how many days they worked while physically ill in the past 3 months. Workers who worked ≥ 1 days while ill were coded as “yes.”

In the service group, lower odds were observed among workers who had paid sick leave (AOR=0.68; 95% CI=0.52–0.89) and government employment (AOR=0.50; 95% CI=0.37–0.68). In the production, transportation, and material moving group, odds were significantly higher among workers who usually worked ≥ 35 hours per week (AOR=1.69; 95% CI=1.15–2.47) and worked while physically ill in the past 3 months (AOR=1.44; 95% CI=1.08–1.92). There were no significant associations between employment characteristics and any tobacco use in the sales and office group and natural resources, construction, and maintenance group.

DISCUSSION

In 2021, an estimated 20% of U.S. working adults used commercial tobacco products, and prevalence was higher among some employment characteristics and occupation groups. Current study findings add to the limited evidence on how tobacco use varies by employment characteristics and occupation, including how the workplace environment might impact workers' tobacco use. The Guide to Community Preventive Services recommends several tobacco cessation interventions that can be implemented in the workplace, including smoke-free policies and reducing out-of-pocket costs for evidence-based cessation treatments.¹⁶ Group therapy programs have also proven effective.¹⁷ Workplaces that have a higher prevalence of tobacco use could consider focused implementation of these strategies to reduce tobacco-related disparities and chronic disease.

Higher odds of any tobacco use were found among rotating or "some other" shift workers and workers who experienced schedule instability. These findings are also consistent with previous studies, which have found higher levels of smoking among workers with rotating shifts¹⁸ and low job control.¹⁹ Providing greater stability of shift work when possible could reduce workers' tobacco use. For example, Hurtado et al²⁰ found reductions in cigarette consumption at workplaces implementing an intervention focused in part on increasing schedule control.

Missing work due to illness, injury, or disability was not associated with any tobacco use. This finding differs from previous studies, which have found tobacco use to be associated with higher illness-related absence from work and subsequent annual excess costs to employers.^{11–13,21,22} One explanation for this finding is the increased role of telework since the COVID-19 pandemic. Telework may increase working while ill,²³ so it is possible that some workers who used tobacco who might have otherwise had an illness-related absence worked from home instead.

Odds of any tobacco use were also lower among government versus private workers. These findings might be

partially explained by availability of smoke-free policies and comprehensive insurance coverage for cessation in some government workspaces.^{24–27} For example, 40 states have laws that prohibit cigarette smoking within government workplaces.²⁵ The Federal Employees Health Benefits Program is required to provide comprehensive cessation support to federal employees, including FDA-approved cessation medication and counseling, with no cost sharing (e.g., no copayments).²⁶ In contrast, there is substantial variability in state and local smoke-free laws for private-sector workplaces,²⁸ as well as insurance coverage for tobacco cessation medications offered in private and public workplaces.²⁹

Limitations

Data collection procedures in 2021 differed from previous years due to challenges resulting from the COVID-19 pandemic¹⁴; findings may be less generalizable and comparable to other years. Due to small sample sizes, this study was unable to examine disparities in tobacco use by individual occupations or tobacco products by occupation group. The NHIS is a cross-sectional survey, limiting our ability to examine causal directions and pathways between employment characteristics and tobacco use.

CONCLUSIONS

One in 5 U.S. workers used some form of tobacco product and differences existed by occupation group, with as many as 1 in 3 workers using tobacco in the national resources, construction, and maintenance occupation group. Tobacco use was also higher among workers employed at workplaces that did not offer health insurance, had no paid sick leave, usually worked ≥ 35 hours per week, were employed with a private company, worked a rotating or "some other" shift, experienced schedule instability, and worked while physically ill in the past 3 months. These findings could inform workplace interventions and policies to better support worker health, including tobacco cessation, among occupation groups with higher tobacco-use prevalence.

ACKNOWLEDGMENTS

The authors would like to thank members of the Centers for Disease Control and Prevention's (CDC's) tobacco-related cancers workgroup for providing their input during paper development. The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the CDC.

There was no funding for this study.

No conflicts of interest were reported by the authors of this paper.

No financial disclosures were reported by the authors of this paper.

CREDIT AUTHOR STATEMENT

Christine M. Kava: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Resources, Software, Validation, Visualization, Writing—original draft, Writing—review and editing. Girija Syamlal: Conceptualization; Data curation; Formal analysis; Methodology; Software; Validation; Writing—review and editing. Brenna VanFrank: Conceptualization, Writing—review and editing. David A. Siegel: Conceptualization, Supervision, Writing—review and editing. S. Jane Henley: Conceptualization, Writing—review and editing. Jonathan Bryant-Genevier: Conceptualization, Writing—review and editing. Jin Qin: Conceptualization, Supervision, Writing—review and editing. Susan A. Sabatino: Conceptualization, Supervision, Writing—review and editing.

SUPPLEMENTAL MATERIAL

Supplemental materials associated with this article can be found in the online version at <https://doi.org/10.1016/j.amepre.2024.05.001>.

REFERENCES

- Centers for Disease Control and Prevention. Tobacco use. National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention; <https://www.cdc.gov/chronicdisease/resources/publications/factsheets/tobacco.htm>. Accessed January 2, 2024.
- Agaku I, King B, Dube SR. Current cigarette smoking among adults—United States, 2011. *MMWR Morb Mortal Wkly Rep*. 2012;61(44):889–894.
- Cornelius ME, Loretan CG, Jamal A, et al. Tobacco product use among adults—United States, 2021. *MMWR Morb Mortal Wkly Rep*. 2023;72(18):475–483. <https://doi.org/10.15585/mmwr.mm7218a1>.
- Jordt SE. Synthetic nicotine has arrived. *Tob Control*. 2023;32(e1):e113–e117. <https://doi.org/10.1136/tobaccocontrol-2021-056626>.
- Boakye E, Osuji N, Erhabor J, et al. Assessment of patterns in e-cigarette use among adults in the U.S., 2017–2020. *JAMA Netw Open*. 2022;5(7):e2223266. <https://doi.org/10.1001/jamanetworkopen.2022.23266>.
- U.S. Bureau of Labor Statistics. Table A-1. Employment status of the civilian population by sex and age. Division of Current Employment Statistics, U.S. Bureau of Labor Statistics; <https://www.bls.gov/news.release/empstat.t01.htm>. Accessed January 2, 2024.
- Syamlal G, King BA, Mazurek JM. Tobacco use among working adults—United States, 2014–2016. *MMWR Morb Mortal Wkly Rep*. 2017;66(42):1130–1135. <https://doi.org/10.15585/mmwr.mm6642a2>.
- Dobson KG, Gilbert-Ouimet M, Mustard CA, Smith PM. Association between dimensions of the psychosocial and physical work environment and latent smoking trajectories: a 16-year cohort study of the Canadian workforce. *Occup Environ Med*. 2018;75(11):814–821. <https://doi.org/10.1136/oemed-2018-105138>.
- Khubchandani J, Price JH. Association of job insecurity with health risk factors and poorer health in American workers. *J Community Health*. 2017;42(2):242–251. <https://doi.org/10.1007/s10900-016-0249-8>.
- Syamlal G, King BA, Mazurek JM. Tobacco product use among workers in the construction industry, United States, 2014–2016. *Am J Ind Med*. 2018;61(11):939–951. <https://doi.org/10.1002/ajim.22907>.
- Lund I, Moan IS, Edvardsen HME. The relative impact of smoking, alcohol use and drug use on general sickness absence among Norwegian employees. *BMC Public Health*. 2019;19(1):500. <https://doi.org/10.1186/s12889-019-6891-1>.
- van den Berg S, Burdorf A, Robroek SJW. Associations between common diseases and work ability and sick leave among health care workers. *Int Arch Occup Environ Health*. 2017;90(7):685–693. <https://doi.org/10.1007/s00420-017-1231-1>.
- Weng SF, Ali S, Leonardi-Bee J. Smoking and absence from work: systematic review and meta-analysis of occupational studies. *Addiction*. 2013;108(2):307–319. <https://doi.org/10.1111/add.12015>.
- National Center for Health Statistics. National Health Interview Survey, 2021 survey description. Division of Health Interview Statistics, National Center for Health Statistics, Centers for Disease Control and Prevention; https://ftp.cdc.gov/pub/Health_Statistics/NCHS/Dataset_Documentation/NHIS/2021/srvydesc-508.pdf. Accessed January 2, 2024.
- U.S. Bureau of Labor Statistics. Standard occupational classification manual, United States, 2018. Executive Office of the President, Office of Management and Budget; 2018. Accessed January 2, 2024.
- Guide to Community Preventive Services. Tobacco. U.S. Department of Health and Human Services; <https://www.thecommunityguide.org/topics/tobacco.html>. Accessed January 2, 2024.
- Cahill K, Lancaster T. Workplace interventions for smoking cessation. *Cochrane Database Syst Rev*. 2014(2):CD003440. <https://doi.org/10.1002/14651858.CD003440.pub4>.
- Tucker P, Harma M, Ojajarvi A, et al. Association of rotating shift work schedules and the use of prescribed sleep medication: a prospective cohort study. *J Sleep Res*. 2021;30(6):e13349. <https://doi.org/10.1111/jsr.13349>.
- Khorrami Z, Zolala F, Haghdoost A, et al. Job-related stress and tobacco smoking: a systematic review. *J Workplace Behav Health*. 2021;36(4):259–277. <https://doi.org/10.1080/15555240.2021.1960854>.
- Hurtado DA, Okechukwu CA, Buxton OM, et al. Effects on cigarette consumption of a work-family supportive organisational intervention: 6-month results from the work, family and health network study. *J Epidemiol Community Health*. 2016;70(12):1155–1161. <https://doi.org/10.1136/jech-2015-206953>.
- Halpern MT, Shikhar R, Rentz AM, Khan ZM. Impact of smoking status on workplace absenteeism and productivity. *Tob Control*. 2001;10(3):233–238. <https://doi.org/10.1136/tc.10.3.233>.
- Berman M, Crane R, Seiber E, Munur M. Estimating the cost of a smoking employee. *Tob Control*. 2014;23(5):428–433. <https://doi.org/10.1136/tobaccocontrol-2012-050888>.
- Steidelmuller C, Meyer SC, Muller G. Home-based telework and presenteeism across Europe. *J Occup Environ Med*. 2020;62(12):998–1005. <https://doi.org/10.1097/JOM.0000000000001992>.
- Clinton WJ. Executive Order 13058 of August 9, 1997: Protecting federal employees and the public from exposure to tobacco smoke in the federal workplace. *Federal Register*. 1997;62:43451–43452.
- Centers for Disease Control and Prevention. State Tobacco Activities Tracking and Evaluation (STATE) system. Office on Smoking and Health, National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention; <https://www.cdc.gov/statesystem/index.html>. Accessed January 2, 2024.
- U.S. Office of Personnel Management. Quit smoking. U.S. Office of Personnel Management; <https://www.opm.gov/healthcare-insurance/special-initiatives/quit-smoking/>. Accessed January 2, 2024.
- American Lung Association. State tobacco cessation coverage database. American Lung Association; <https://www.lung.org/policy-advocacy/tobacco/cessation/state-tobacco-cessation-coverage-database>. Accessed January 2, 2024.
- Tynan MA, Holmes CB, Promoff G, Hallett C, Hopkins M, Frick B. State and local comprehensive smoke-free laws for worksites, restaurants, and bars—United States, 2015. *MMWR Morb Mortal Wkly Rep*. 2016;65(24):623–626. <https://doi.org/10.15585/mmwr.mm6524a4>.
- Linnan LA, Cluff L, Lang JE, Penne M, Leff MS. Results of the workplace health in America survey. *Am J Health Promot*. 2019;33(5):652–665. <https://doi.org/10.1177/0890117119842047>.